

WILLIAM J. BACHA, JR. AND LINDA M. BACHA

WILEY-BLACKWELL



COLOR ATLAS OF VETERINARY HISTOLOGY Third Edition

COMPANION CD-ROM

This book is accompanied by a companion CD providing the more than 900 photographs digitally in an atlas interface with zoom capability. There are also interactive multiple choice study questions available for each chapter.

Note on access

Wiley publishes in a variety of print and electronic formats and by print-on-demand. Some material included with standard print versions of this book may not be included in e-books or in print-on-demand. If this book refers to media such as a CD or DVD that is not included in the version you purchased, please check for the material at http://booksupport.wiley.com. For more information about Wiley products, visit www. wiley.com

VETERINARY HISTOLOGY

Third Edition

William J. Bacha, Jr., PhD

Professor Emeritus
Department of Biology
Rutgers University
Camden College of Arts and Sciences
Camden, New Jersey

Linda M. Bacha, MS, VMD

Assistant Professor of Biology Department of Biology Camden County College Blackwood, New Jersey This edition first published 2012 © 2012 by John Wiley & Sons, Ltd. First and Second editions © 1990 and 2000, Lippencott Williams & Wilkins

Wiley-Blackwell is an imprint of John Wiley & Sons, formed by the merger of Wiley's global Scientific, Technical and Medical business with Blackwell Publishing.

Registered office: John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK

Editorial offices: 2121 State Avenue, Ames, Iowa 50014-8300, USA The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK 9600 Garsington Road, Oxford, OX4 2DQ, UK

For details of our global editorial offices, for customer services, and for information about how to apply for permission to reuse the copyright material in this book please see our website at www.wiley.com/wiley-blackwell.

Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by Blackwell Publishing, provided that the base fee is paid directly to the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923. For those organizations that have been granted a photocopy license by CCC, a separate system of payments has been arranged. The fee codes for users of the Transactional Reporting Service are ISBN-13: 978-0-4709-5851-3/2012.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks, or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. This publication is designed to provide accurate and authoritative information in regard to the subject matter covered. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

Library of Congress Cataloging-in-Publication Data

Bacha, William J.

Color atlas of veterinary histology / William J. Bacha Jr., Linda M. Bacha. – 3rd ed.

p.; cm.

Includes bibliographical references and index.

ISBN-13: 978-0-470-95851-3 (hard cover : alk. paper)

ISBN-10: 0-470-95851-0 (hard cover : alk. paper)

1. Veterinary histology-Atlases. I. Bacha, Linda M. II. Title.

[DNLM: 1. Animals, Domestic-anatomy & histology-Atlases.

2. Anatomy, Veterinary–Atlases. SF 757.3]

SF757.3.B33 2011

636.089'1018-dc23

2011020601

A catalogue record for this book is available from the British Library.

Set in 10/11 pt Sabon by Toppan Best-set Premedia Limited

Disclaimer

The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation warranties of fitness for a particular purpose. No warranty may be created or extended by sales or promotional materials. The advice and strategies contained herein may not be suitable for every situation. This work is sold with the understanding that the publisher is not engaged in rendering legal, accounting, or other professional services. If professional assistance is required, the services of a competent professional person should be sought. Neither the publisher nor the author shall be liable for damages arising herefrom. The fact that an organization or website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or website may provide or recommendations it may make. Further, readers should be aware that Internet websites listed in this work may have changed or disappeared between when this work was written and when it is read.

TO OUR PARENTS, ANNA and BILL LOLA and CHIP

PREFACE TO THE THIRD EDITION

t is our hope that students and researchers will continue to find this atlas a useful resource. In this edition, we have expanded many of the beginning chapters that introduce the basic types of tissues to provide the user with a stronger foundation in histology. The glossary has been expanded, and other extras have been included that we hope will be valuable.

Once again, many thanks to all of those who have made the first and second edition of our atlas possible! We were able to prepare our page layouts for the third edition thanks to the scanner given to us

by Aunt Lila and Uncle Jack McKean and Lola and Chip Wood. Thanks to Jessie Bacha for proofreading our work, and to Tristan Bacha for keeping busy while we worked! We greatly appreciate the helpful suggestions from Professor Nancy Gartland and the students at the University Of Pennsylvania School of Veterinary Medicine, and from reviewers and users of the previous edition. Finally, thanks to Nancy Turner, Erica Judisch, Tracy Petersen, Erin Magnani, and all of the other people at Wiley-Blackwell for their role in the production of this edition.

PREFACE TO THE SECOND EDITION

e wish to thank those who have used the first edition for their suggestions. We believe the incorporation of many of these recommendations will make this edition even more helpful to the user.

To this end, we have updated the material for the second edition by scanning all of the original kodachromes and relabeling the art. We have added thirteen new photographs and have enlarged over one

hundred others. Four of the original black and white line drawings have also been redrawn. Also, a glossary of nearly 750 words has been added.

The style, format, and purpose of this edition remain essentially unchanged from the first edition. We continue to view the atlas as a useful, benchside reference for those interested in understanding and interpreting histologic and cytologic preparations.

PREFACE TO THE FIRST EDITION

Ithough we have written this atlas primarily to fulfill a need of the student of veterinary medicine, we believe that clinicians, private practitioners, and researchers will find it a useful reference for normal tissues and organs. Currently, students rely heavily, if not exclusively, on atlases of human histology for guidance in the laboratory. There are, of course, similarities between organs and tissues of domestic animals and those of humans. There are also differences, however, and these are rarely encountered in atlases dealing specifically with human histology.

Our aim has been to compare the histologic structure of organs in a variety of domestic animals. We have used representative examples in instances where tissues and organs from different animals share a common structure. Wherever differences exist, we have tried to provide examples that are characteristic of a particular group of animals. Our selection of animals includes the dog, cat, horse, cow, sheep, goat, pig, and chicken because they are most frequently referenced in veterinary school curricula.

All photomicrographs and drawings are original. Some drawings were done freehand, while others were made with the aid of a camera lucida. Light microscopy and colored photomicrographs have been used exclusively. We have chosen color rather than black and white because of its correspondence to stained preparations. With the exception of the few histologic preparations loaned to us by generous donors or purchased from a dealer, slides were prepared by the authors. Fresh organ samples were obtained from a slaughterhouse or from animals that were euthanized for various reasons. With the exception of smear preparations (blood, bone marrow, and vaginal), mesenteric spreads, ground bone, and a single plastic section, slides were prepared using the paraffin method. All slides were stained with hematoxylin and eosin unless otherwise noted. Magnifications of photomicrographs are total magnifications (enlargement of photograph × objective × projector lens). Throughout the atlas, hollow structures, for example, blood vessels, kidney tubules, and alveoli, are usually identified by labeling the lumen of the structure.

ACKNOWLEDGMENTS FROM THE FIRST EDITION

elp is often just around the corner. Dr. Henry Stempen, whose office was down the hall from ours at Rutgers University in Camden, New Jersey, stopped by one day and volunteered his artistic talents. We'd like to thank him for his excellent pen and ink drawings of various animal parts, which are somewhat removed from the fungi he usually draws. Our gratitude also to Ms. Kathleen Carr for her secretarial services. Special thanks are extended to Dr. Edward Zambraski. Ms. Kathleen O'Hagan, and Ms. Gail Thomas of Cook College, Rutgers University, for making fresh porcine material available to us, and to Dr. Barry Jesse and Dr. James Harner for supplying us with sheep parts.

Without the unqualified use of the facilities and equipment of the Biology Department of Rutgers, our tissue processing and photomicrography could not have been accomplished. Our special thanks to the department for this courtesy.

This book would never have had a beginning were it not for the generosity of Dr. Leon Weiss, Department of Animal Biology, University of Pennsylvania School of Veterinary Medicine, who invited us to teach in the veterinary histology laboratory and kindly allowed us access to the slide collection and facilities of the depart-

ment. We would also like to express appreciation to the following individuals from the University of Pennsylvania School of Veterinary Medicine: Mr. Richard Aucamp and Mrs. Kathy Aucamp, who provided us with specimens, slides, advice, and assistance in a variety of other ways; Dr. Mark Haskins for kindly making available fresh canine and feline material; Dr. John Fyfe and Dr. Vicki Meyers-Wallen for supplying us with canine vaginal smears: Dr. and Mrs. Loren Evans and Dr. David McDevitt for lending us reference material; Dr. Peter Hand and Ms. Graziella Mann for providing material on the nervous system; and Dr. Helen Acland, Dr. Linda Bachin, Mr. James Bruce, Dr. Sherrill Davison, Ms. Dawn Dowling, Dr. Robert Dyer, Dr. Robert Eckroade, Dr. George Farnbach, Dr. David Freeman, Dr. Wendy Freeman, Dr. Alan Kelly, Mr. Joseph McGrane, and Dr. Mary Sommer for their time and consideration in helping us to obtain tissue specimens.

We are grateful to Dr. Carol Jacobson and the Department of Anatomy of the Iowa State University College of Veterinary Medicine for providing valuable slide preparations and text material.

Our gratitude is also extended to Hill's Pet Products, Topeka, Kansas, and

Pitman-Moore, Inc., Washington Crossing, New Jersey, for their generous financial assistance.

Many thanks also to: Dr. Caroline Czarnecki of the University of Minnesota, College of Veterinary Medicine, for providing copies of her informative laboratory guide; Dr. Deborah Ganster, Dr. James Lawhead, Dr. Virginia Pierce, Dr. Maria Salvaggio, Dr. Barbara Strock, and Dr. Cindi Ward for assisting us in obtaining tissue samples; Mr. Jeff Bringhurst, Bringhurst Brothers, Tansboro, New Jersey, for allowing us access to fresh large material; the Longenecker animal Hatchery, Elizabethtown, Pennsylvania, for providing chicken specimens; Ms. Susan Ulrich, Cornell University Press, for lending us a difficult-to-obtain reference; the helpful people at Optical Apparatus Company Inc., Ardmore, Pennsylvania, for supplies and for assistance with equipment for the microscope; and Mr. Charles Behl and Mr. James Durso of Webb and

Company Inc., Cherry Hill, New Jersey, for their courteous service and helpful advice.

We are indebted to Mr. William J. Bacha, Sr., for building a super light box for us, and to Mr. Thomas H. Wood, Jr., for providing black and white prints of our photomicrographs, which saved us countless hours of drudgery in the darkroom. Thanks to Barbara Frasco, Esq., for her helpful advice. Our hats are off also to Snuff, Chew, Chapter Seat, Angel, Clyde, and all the other animals for their participation.

We also wish to extend our gratitude to all at Lippincott Williams & Wilkins whose efforts helped bring this second edition into being. We are especially grateful to Carroll C. Cann and Jennifer D. Weir for their professional advice, courtesy, and assistance.

William J. Bacha, Jr. Linda M. Bacha

CONTENTS

| 1 | General Principles of Histology | 3 |
|------------|--|------------|
| 2 | Epithelium | 11 |
| 3 | Connective Tissue Proper and Embryonic Connective Tissue | 19 |
| 4 | Cartilage | 27 |
| 5 | Bone Tissue | 31 |
| 6 | Blood | 41 |
| 7 | Bone Marrow | 53 |
| 8 | Muscle Tissue | 57 |
| 9 | Nervous System | 65 |
| 10 | Cardiovascular System | 77 |
| 11 | Lymphatic System | 89 |
| 12 | Integument | 105 |
| 13 | Digestive System | 139 |
| 14 | Urinary System | 183 |
| 15 | Respiratory System | 195 |
| 16 | Endocrine System | 211 |
| 1 <i>7</i> | Male Reproductive System | 225 |
| 18 | Female Reproductive System | 243 |
| 19 | The Eye | 267 |
| 20 | The Ear | 283 |
| | Glossary | 291 |
| | Bibliography | 309 |
| | Index | 311 |

COLOR ATLAS OF VETERINARY HISTOLOGY Third Edition

GENERAL PRINCIPLES OF HISTOLOGY

PREPARATION OF HISTOLOGIC SECTIONS

histologic section is a thin slice of tissue, varying, usually, from 0.5 to 10 or more micrometers (μ) thick. In preparing such a section, a piece of tissue is either infiltrated with a supporting medium or frozen and is then cut with an instrument called a microtome. Sections obtained from tissue infiltrated with plastic can be as thin as 0.5 μ and show superior detail. Excellent preparations as thin as 2 or 3 μ also can be made from tissue infiltrated with paraffin-based embedding media. Sections are affixed to microscope slides and colored with one or more stains to increase the visibility of various cellular and intercellular components.

Schematically, Figure 1.1 outlines various steps involved in producing a stained histologic slide using the paraffin procedure. After being removed from an animal, a tissue or organ is cut into pieces. These pieces are placed into a fixative such as buffered formalin or Bouin's, which, ideally, preserves normal morphology and facilitates further processing. After fixation, the specimen is dehydrated by transferring it through a series of alcohols of increasing concentrations to 100% alcohol. Next, it is placed into a substance such as xylene or xylene substitute, which is miscible with both 100% alcohol and paraffin. This intermediate step (called clearing) is essential before infiltrating the dehydrated tissue with paraffin because alcohol and paraffin do not mix. During infiltration, melted paraffin completely replaces the xylene. This procedure is done in an oven at a temperature just above the melting point of the paraffin. When infiltration is complete, the specimen is transferred to an embedding mold of fresh paraffin, which is allowed to harden. Then the mold is removed and excess paraffin is trimmed away.

The block of paraffin is then secured to the microtome and oriented appropriately with respect to the knife. With each revolution of the microtome handle, the specimen moves through the blade and a section of the desired thickness is produced. Each successive section adheres to the preceding one, forming a continuous ribbon. Subsequently, one or more sections are carefully separated from the ribbon and transferred to the




Figure 1.1. The various steps involved in producing a histologic slide using the paraffin method.

surface of warm water in a waterbath. This softens the paraffin and flattens the section, eliminating wrinkles. The flattened section is floated onto a slide, which is then placed on a warming table. As the preparation dries, the section adheres to the surface of the slide.

Next, the paraffin is removed with xylene or another appropriate solvent and the specimen is rehydrated. It is then stained, dehydrated, cleared (made transparent) with xylene, covered with a resinous mounting medium, and topped with a cover-slip.

Various stains are available to the histologist. Hematoxylin and eosin (H&E) is a frequently used combination of stains. Hematoxylin imparts a purple color to substances, but must be linked to a metallic salt called a mordant before it can function effectively. This combination, called a lake, carries a positive charge and behaves as a basic (cationic) stain. The lake combines electrostatically with negatively charged radicals such as phosphate groups of nucleoproteins. Substances that become colored by a basic stain are said to be basophilic. Methylene blue, toluidine blue, and basic fuchsin are basic stains. Unlike hematoxylin, these stains have molecules that carry a positive charge of their own and do not require a mordant. Acidic (anionic) stains carry a negative charge and color cell or tissue components that bear positive charges. Eosin is an acid stain. It imparts an orange or red color to acidophilic substances. Other commonly used acid stains are orange G, phloxine, and aniline blue.

In addition to the widely used H&E staining procedure, numerous other stain combinations and techniques are available. Some are especially useful for identifying certain tissue elements. For example, trichrome procedures such as Mallory's and Masson's specifically stain collagenous fibers within connective tissue. Orcein and Weigert's resorcin fuchsin are stains used to color elastic fibers, providing a means of distinguishing them from other fibrous elements. Reticular fibers and nervous tissue components such as neurons, myelin, and cells of the neuroglia can be stained by procedures employing the use of silver. There are also special histochemical and immunohistochemical procedures that make possible the localization of various carbohydrates, lipids, and proteins found in tissue. Lastly, stains such as Wright's and Giemsa's (Romanovsky stains) are available for differentiating the various cells found in blood and bone marrow.

INTERPRETING SECTIONS

One must know the gross structure of an organ before a histologic section from it can be comprehended. It is also helpful to know how the section was cut, that is, whether it was a cross section (x.s.), a longitudinal section (l.s.), or an oblique slice through the organ. Was the cut made through the entire organ or only through a portion of it? Frequently, prepared slides are labeled indicating the particular orientation of the section. This is not important in an asymmetric organ such as the spleen or liver because their appearance would be unaffected by the direction of the cut. Conversely, the small intestine is radially symmet-

ric and its appearance is affected by the direction of the

The three-dimensional structure of organs and their components also must be considered when examining a histologic preparation. Cells are three-dimensional objects differing in size and shape. For example, some are long and thin, some cuboidal, and others ovoid. They may have a random or specific arrangement within an organ. How they appear depends on their shape as well as how they were cut. Imagine how the spindle-shaped and tall columnar cells shown in Figure 1.2A would look if sectioned in various planes. Note that the nucleus may or may not be included in a particular cut through a cell.

The histologist examines multicellular structures having a wide variety of shapes. Some are hollow, some branch repeatedly, some open onto surfaces, etc. Figure 1.2, B and C, and Figure 1.3 show a variety of three-dimensional structures and how they would appear if cut at different levels. Examine these carefully. They will help you to understand situations you will encounter on actual slides.

HELPFUL HINTS

Be sure that the lenses of your microscope are clean before you begin examining slides. Use a piece of lens paper or a soft, clean cloth such as an old (but clean) linen handkerchief. If the lenses have been coated with oil or another substance, remove it using lens tissue moistened sparingly with a glass cleaner such as Windex. Slides also should be cleaned using a soft, lint-free cloth or tissue moistened with glass cleaner.

Every microscope should have a pointer in the ocular. This is usually supplied by the manufacturer, but can be made from a short piece of hair. The latter is cemented into place inside the ocular with a dab of quick-drying glue or nail polish. Without a pointer, it is not possible to accurately indicate an object in the microscope field for another observer.

Before beginning a session at the microscope, make sure that the fine-adjustment knob is near the middle of its range of rotation. If you do not, you may find that the knob is at the limit of its excursion when you are busily making observations. At that point, you must stop everything and correct it.

It is also a good habit to examine your slide with the unaided eye before placing it on the stage of your microscope. By doing so you will gain information about the gross aspects of the specimen and be more likely to center it properly over the light source. Centering is especially important for small specimens that might otherwise be difficult to locate. Also, make sure that you put the slide on the stage with the cover glass uppermost. If the slide is upside down, you will not be able to focus on it with the high-power lenses. Do not snicker. We have seen this happen often in the teaching laboratory!

It is always a good idea to start your observations using the lowest power objective available on your microscope. This is usually the 4x lens. The field of view will be




Figure 1.2. A. Slices, indicated by numbered planes, taken through two different types of cells would appear as identified by the prime numbers. Only if the plane of the cut passes through the nucleus will the latter be seen. B and C. Planes of section taken from different levels in four separate multicellular objects are illustrated. Note how the appearance of sections varies with the level of the cut.




Figure 1.3. The prime numbers illustrate sections resulting from transverse (4), oblique (1), and longitudinal (2, 3, 5, 6) cuts made through a plate of cells bearing hollow projections (above plate) and invaginations (below plate). Plane 3 differs from the others because it passes only through the cellular wall of a projection, and not the lumen; therefore, section 3' appears as a plate of cells rather than a hollow structure. You should also be aware that structures may often appear unrelated to a surface or another object, when in fact they are. Compare planes 5 and 6 with sections 5' and 6', where continuity of the invagination with the surface is evident only in 6 and 6'. While not apparent from a single section, such continuity would be evident if an uninterrupted series of sections through the entire invagination were made and examined.

large, enabling you to locate regions of special interest more easily. When you locate something you wish to examine at a higher magnification, center the object in the middle of the field of view. Then, when you change to a stronger lens, the object should be somewhere in the field.

Binocular microscopes often have at least one ocular that can be adjusted to accommodate your vision. It is important that you adjust this properly if you want to have a comfortable, headache-free session at the microscope. Assuming that your microscope is of the binocular type and that it has at least one adjustable ocular, you should first bring the specimen into focus with the ocular that is not adjustable by using the fine-adjustment knob. When you have done this, focus the other eye using the adjustable ocular. Use of this procedure will ensure a proper focus for both eyes and prevent eye strain.

Yikes! After you turn on the microscope, you do not see any light as you look through the oculars! Before you replace the bulb, try the following:

- Check the light control lever or knob to make sure the light intensity is not set too low or turned down to zero.
- Check that the objective lens is clicked fully into position.
- Make sure that the plug is fitted properly into the electrical outlet and that the outlet is working.

Bright, even lighting is absolutely essential to effective microscopy. The best way to achieve this is to use Köhler illumination. This can be obtained with any microscope that is equipped with both a condenser aperture diaphragm (the one in the condenser) and a field diaphragm (the one in the light source). If you have such an instrument, proceed as follows:

- 1. Center the light source, using the directions you received with the microscope.
- 2. Open both the field and aperture diaphragms fully.
- 3. Raise the condenser to its uppermost position.
- 4. Place a specimen on the stage and focus on it using the $10 \times$ objective.
- Close the field diaphragm so that its leaves are clearly imaged in the field of view.
- 6. Center the image of the diaphragm by manipulating the condenser centering screws, then open the field diaphragm until its leaves disappear just beyond the edge of the field of view.
- 7. Remove an ocular and, while looking into the back aperture of the objective, close the aperture diaphragm completely and then open it until it is about 75% of being fully open.

You now have Köhler illumination. If you want to increase or decrease the light intensity, use the rheostat or neutral-density filters, but do not adjust the condenser aperture diaphragm or field diaphragm. If the aperture diaphragm is open to excess, the image will lack some contrast and be flooded with light. If it is closed too far,

there will be a loss of resolution and increase in contrast. This increase in contrast is often confused with sharpness or high resolution; this is a common error in microscopy. All of the above adjustments (except for centering the light source) must be made each time a different objective is used.

If your microscope lacks a field diaphragm, you will not be able to obtain Köhler illumination. You can still acquire good and useful lighting, however. Place a specimen on the stage, open the aperture diaphragm fully, and adjust the light intensity with the rheostat so that it is comfortable for your eyes. Be sure that the condenser is raised to its highest position, or close to it, when you do this. Now, remove an ocular and look at the back aperture of the objective. Close the aperture diaphragm fully and then open it until it is about 75% of being fully open. This will provide proper lighting for most purposes. If you should need more or less illumination, make adjustments only with the rheostat or neutral density filter; do not use the aperture diaphragm.

To get the most from a specimen, you must avoid being a passive microscopist, that is, one who finds an object and then stares at it admiringly without making further adjustments of the focus. Get into the habit of focusing continuously with the fine adjustment as you peruse a slide, because even though a tissue section may be only a few micrometers thick, the depth of field of the higher power objectives may be less than the thickness of the specimen. Therefore, if you do not focus repeatedly as you examine a preparation, you will certainly miss seeing structural detail that might be important to your work.

You might like to return to a particular location on your slide preparation at a future time. Remembering landmarks in the vicinity of the object of interest will aid you in locating the object later. A more expedient way of relocating structures is by using verniers, which are mounted on both the X and Y axes of the mechanical stage. A vernier consists of two parallel, graduated, sliding scales, one long and one short. The smaller scale is 9 millimeters (mm) long and is divided into 10 subdivisions (0 to 10). The larger scale is several centimeters (cm) long and is graduated in millimeters, for example, 0 to 80 or 100 to 160. To relocate an object on a slide, you must first center it in the microscope field. Once this has been done, you establish its location by reading each of the verniers (X and Y). For example, the 0 point on the small scale of the vernier on the X axis might be located somewhere between lines 42 and 43 on the larger scale (Figure 1.4). To determine its specific location, find the line on the small scale that coincides exactly with a line on the longer scale. Then count, on the smaller scale, the number of spaces between 0 and the point of coincidence. This number is your decimal point. In the example given (Figure 1.4), the




Figure 1.4. Small and large vernier scales.

decimal is 0.6 and you should read 42.6 as the vernier value. Do the same for the other vernier (Y) and record the numbers for both. In the future, if you want to return to the same location, simply secure the slide to the mechanical stage and move the stage controls until the verniers are adjusted to the numbers you previously recorded. These manipulations will have returned the slide to its former position, and the object you are looking for should be somewhere within the microscope field.

By knowing the approximate diameter of a red blood cell in a section, you can estimate the size of other tissue components. Therefore, it is useful to know that in tissue sections prepared by the paraffin method the average size of erythrocytes for each of the following animals is as follows:

Goat: 2.4μ diameter (smallest erythrocytes of the domestic mammals)

Dog: $4.9 \,\mu$ diameter (largest erythrocytes of the domestic mammals)

Chicken: 9.4 µ long

Each average value is based on a total of 20 to 30 cells that were measured from five different slide preparations of tissues embedded in Paraplast X-TRA (Monoject Scientific, Division of Sherwood Medical, St. Louis, MO 63103).

ARTIFACTS

Folds, knife marks, stain precipitate, spaces (where none belong), shrinkage, and air bubbles are examples of common imperfections seen in slide preparations. They were introduced during processing and are called artifacts. Figures 1.5 through 1.9 are examples of such artifacts.

Troubleshooting a blurred or cloudy image:

- If an image is cloudy or blurry, the oculars and/or objective lenses may need to be cleaned. To determine if an ocular needs to be cleaned, turn it as you look through the microscope. A dirty mark or smear will rotate if the ocular is not clean
- If the image is still blurred, clean the objective lenses.
- Make sure the slide is resting on the stage properly and is right side up! Sometimes an image cannot be focused clearly because the slide is on the stage upside down!
- Are you using the oil immersion lens without a drop of oil on the slide?
- Be sure that the light source is not partially blocked by something, such as the electric cord or a filter holder below the stage.




Figure 1.5 × 62.5




Figure 1.6 × 62.5




Figure 1.7 × 62.5




Figure 1.8 × 25




Figure 1.9 × 62.5

| | KEY | |
|--------------------------------------|---|--|
| 1. Dermis 2. Epidermis 3. Fold | 4. Knife mark5. Separation artifact6. Stain precipitate | |
| | | |

Figure 1.5. Stain Precipitate, Cartilage, Dog. Occasionally, solutions accumulate precipitate that may stick to the surface of tissue sections during the staining procedure.

Figure 1.6. Separation (Space) Artifact, Skin, Dog. Tissues may be subjected to excessive pressures, tensions, or shrinkage during processing, resulting in separations within otherwise intact tissue.

Figure 1.7. Crackling Artifact, Thymus, Horse. Highly cellular tissues, for example, thymus, liver, pancreas, and spleen, often show numerous tiny cracks throughout. Also note that this specimen is not in sharp focus.

Figure 1.8. Knife Marks and Folds, Esophagus, Horse (Masson's). Knife marks (scratches) in the tissue section may be caused by defects in the microtome knife or by accumulations of debris on the knife edge. Folds occur when the tissue sections fail to spread properly on the surface of the slide.

Figure 1.9. Fold, Aorta, Pig. In a tissue section, folds are raised areas that frequently overlap. Note that portions of this picture are not in sharp focus.

EPITHELIUM

pithelium is one of the four main types of tissue in the body along with connective tissue, muscle tissue, and nervous tissue.

GENERAL FEATURES

Epithelium is a ubiquitous tissue that covers surfaces, lines cavities, and forms glands of the body. It is comprised entirely of cells separated only by a thin layer of intercellular substance that helps hold them together. The cells are supported by a thin extracellular basement membrane that separates them from the underlying connective tissue.

The free (apical) surfaces of epithelial cells may possess cilia (motile processes), microvilli (short finger-like projections of the cell membrane), or stereocilia (long microvilli).

When the epithelium consists of a single layer of cells, it is called a **simple epithelium**. If it is formed from two or more layers of cells it is said to be a **stratified epithelium**. The individual cells of either type of epithelium may be **squamous**, **cuboidal**, or **columnar**. In profile view, squamous cells are flat, cuboidal cells are short cells that are approximately as tall as they are wide, and columnar cells are rectangular cells that are taller than they are wide.

SIMPLE EPITHELIAL TISSUE

Simple epithelia, formed by a single layer of cells, are found in many different locations throughout an animal's body. They line the heart; blood and lymphatic vessels; and

pleural, pericardial, and abdominal cavities; and they form the secretory units of many glands. They also line the stomach, intestines, and portions of the respiratory system. Types of simple epithelia, simple squamous, simple cuboidal, and simple columnar, are named according to the shape of the cells. Pseudostratified columnar is a special form of simple epithelium. Although it appears in profile to be multilayered, it is not. Its apparent stratification is an illusion that results because the nuclei of cells of different heights occur at different levels. All of its cells, though, are in contact with the basement membrane and not layered on top of each other.


STRATIFIED EPITHELIAL TISSUE

Stratified epithelia consist of two or more layers of cells, with only the bottommost layer in contact with the basement membrane. Each additional layer is added on top of the previous one and does not contact the basement membrane. The total number of layers can vary from two to a dozen or more. Classification of stratified epithelia depends upon the shape of the cells in the outermost (surface) layer of the epithelium. If the outermost layer is squamous, the epithelium is stratified squamous; if it is cuboidal, the epithelium is stratified cuboidal; if it is columnar, the epithelium is stratified columnar. Stratified epithelia can be found in such places as the mouth, esophagus, larynx, epidermis, vagina, and anal canal. Transitional epithelium is the term applied to a special category of stratified epithelium found only in the urinary system. The shape of its cells changes with the amount of fluid pressure applied against them. For example, its surface cells have a domed shape when the urinary bladder is empty but become flattened as the bladder fills with urine.

Summary of Types of Epithelial Tissue

Simple Epithelium
Simple squamous
Simple cuboidal
Simple columnar
Pseudostratified columnar
Stratified Epithelium
Stratified squamous
Nonkeratinized
Keratinized
Stratified cuboidal
Stratified columnar
Transitional

| Word Roots | | | |
|------------|--------------|---|--|
| ROOT | MEANING | EXAMPLE SENTENCE | |
| Column | A pillar | A <i>columnar</i> cell is tall and slender like a pillar. | |
| Epi | Upon or over | Epithelial tissue covers surfaces and lines cavities and is above deeper tissues. The term epithelium is a noun, and epithelial is an adjective. For example, one can speak of a simple squamous epithelium , or simple squamous epithelial tissue. Plural = epithelia . | |
| Pseudo | False | Pseudostratified epithelium appears to be stratified, but it is not. | |
| Squam | A scale | Squamous cells, being thin and flat, resemble scales that cover the body of a fish. | |
| Strat | A layer | A stratified epithelium is layered. | |





Pseudostratified Columnar






Figure 2.1. Types of Epithelia.








Figure 2.4 × 250




Figure 2.5 × 250

KEY

- 1. Blood vessel, lumen
- 2. Collagenous fiber
- 3. Connective tissue
- 4. Cuboidal cell, nucleus
- 5. Fibroblast, nucleus
- 6. Hepatocytes

- 7. Lymphatic vessel, lumen
- 8. Smooth muscle cell, nucleus
- 9. Squamous cell, nucleus
- 10. Tubule, lumen
- 11. Urinary space

Figure 2.2. Simple Squamous Epithelium of Blood Vessels and Lymphatic Vessels, Submucosa of the Stomach, Pig. Blood vessels and lymphatic vessels (and the heart) are lined by a simple squamous epithelium that is specifically called an endothelium in those locations. The cytoplasm of the squamous cells is sparse, and generally only the nucleus is visible. Characteristically, the nuclei appear flat in histologic section. If the squamous cells are bunched up, such as those lining a vessel that is contracted, their nuclei appear round.


Figure 2.3. Simple Squamous Epithelium, Kidney, Sheep. The outer layer of Bowman's capsule of a renal corpuscle is called the parietal layer. It is formed by a single layer of squamous cells. A magnified example of the parietal layer of Bowman's capsule is shown in Figure 2.4. Tubules lined by a simple cuboidal epithelium are evident.

Figure 2.4. Simple Squamous Epithelium, Kidney, Sheep. The flat nuclei of the squamous cells of the parietal layer of Bowman's capsule are visible, forming the outer lining of the urinary space of the renal corpuscle.

Figure 2.5. Simple Squamous Epithelium, Mesothelium, Liver, Cat. The surface of the liver is covered by a single layer of squamous cells. Mesothelium is a specific term for the epithelium of the serous membranes of the body, namely the peritoneum, pleura, and pericardium.

Helpful Hint

The nuclei of a squamous epithelium will not always appear regularly spaced. When a layer of squamous cells of an organ is sectioned, the slice may or may not pass through the nucleus of each cell, resulting in uneven spacing.






Figure 2.6 × 250




Figure 2.7 \times 360

| | KEY | |
|---|------------------|--|
| Cuboidal cell, nucleus Squamous cell, nucleus | 3. Tubule, lumen | |

Figure 2.6. Simple Cuboidal Epithelium, Kidney, Cow (Trichrome). The lining of the tubules shown here consists of a single layer of cuboidal cells.

Figure 2.7. Simple Cuboidal and Simple Squamous Epithelium, Medulla of Kidney, Sheep. The boundaries between the cuboidal cells are not distinct in the tubules lined by simple cuboidal epithelium in this image. To determine the shape of the cells, imagine drawing lines between the nuclei of adjacent cells to approximate their borders. Note the simple squamous epithelium that lines the capillaries and the thin segments of the loop of Henle.

Helpful Hints

Organs are formed by various types of tissues, so you may feel overwhelmed when you begin to look at a histologic section of an organ to study epithelial tissue.

- Remember that epithelial tissue is found lining a cavity or covering a surface, so begin to examine an organ at low magnification and look for epithelium to be bordering the white space of a cavity or surface.
- Epithelial tissue is formed by cells that are close together, with little or no extracellular matrix. Look for closely arranged basophilic nuclei of the epithelial cells to help locate an epithelium.
- The shape of nuclei of the cells will help you name the type
 of epithelium. The nuclei of squamous cells are typically flat
 (sometimes round if the cells are bunched up, as in those
 lining a contracted vessel) and the cytoplasm is usually not
 visible. The nuclei are round in cuboidal cells and typically
 elongated in columnar cells.






Figure 2.9 \times 250




Figure 2.10 × 250





Figure 2.11 \times 250



| rigure 2.12 | × 230 |
|---|---|
| | KEY |
| Basement membrane Cilia Collagenous fiber Columnar cell, nucleus Gastric gland Gastric pit, lumen Goblet cell Jejunum, lumen Lamina propria | Lymphocyte Mucus precursor region Pseudostratified columnar epithelium Simple columnar epithelium Smooth muscle tissue Stomach, lumen Striated border Trachea, lumen Similar area magnified in Figure 2.9 |

Figure 2.8. Simple Columnar Epithelium, Pyloric Stomach, Cat. This low power view shows the location of the simple columnar epithelium that lines the lumen of the stomach and invaginates to form gastric pits. Figure 2.9 is a magnified view of the epithelium of the stomach similar to the area outlined by the rectangle in this

Figure 2.9. Simple Columnar Epithelium, Pyloric Stomach, Dog. The columnar cells that line the lumen of the stomach and the gastric pits have apical cup-shaped regions that contain mucus precursor in living tissue. Note the elongated nuclei that are typical of columnar cells.

Figure 2.10. Simple Columnar Epithelium, Jejunum, Dog. The jejunum is lined by a simple columnar epithelium. A striated border that consists of numerous microvilli is evident. Goblet cells and migrating lymphocytes are present among the columnar cells. In the upper part of this image, the simple columnar epithelium was cut obliquely, making it appear atypical.

Figure 2.11. Ciliated Pseudostratified Columnar Epithelium, Trachea, Sheep. A pseudostratified columnar epithelium is formed by a single layer of cells of different sizes whose nuclei are at different levels, so that the epitheliums appear stratified. However, it is a simple epithelium because all of the epithelial cells contact the basement membrane, although this relationship is not visible.

Figure 2.12. Ciliated Pseudostratified Columnar Epithelium, Trachea, Cow. Goblet cells are evident in this example of a pseudostratified columnar epithelium.

For additional examples of simple epithelia, see these figures:

Simple squamous: 8.2, 10.1 to 10.10 (showing endothelium of blood vessels), 14.9, 14.10
Simple cuboidal: 9.9, 16.15, 17.10, 18.21, 18.30
Simple columnar: 13.83, 13.88, 13.115, 13.139, 18.23
Pseudostratified columnar: 15.3, 15.4, 15.14, 15.15,

15.17, 15.40, 17.10, 18.85




Figure 2.13 × 125




Figure 2.14 × 125




Figure 2.15 $\times 125$




Figure 2.16 × 250




Figure 2.17 \times 250

KEY

- 1. Bistratified cuboidal epithelium
- 2. Collagenous fiber
- 3. Dermis
- 4. Esophagus, lumen
- 5. Keratinized cells
- 6. Lamina propria
- 7. Lymphatic vessel, lumen 8. Pharynx, lumen
- 9. Separation artifact
- 10. Smooth muscle tissue
- 11. Squamous cell, nucleus
- 12. Stratified squamous epithelium

Figure 2.13. Stratified Squamous Epithelium, Nonkeratinized, Epiglottis, Goat. Only cells of the basal layer of a stratified epithelium contact the basement membrane. In a nonkeratinized epithelium, the nuclei of the cells are visible throughout the epithelium, even those of the most superficial cells. Note the simple squamous epithelium that lines the lymphatic vessel in the underlying connective tissue.

Figure 2.14. Stratified Squamous Epithelium, Nonkeratinized, Esophagus, Cat. A nonkeratinized stratified squamous epithelium is visible lining the collapsed lumen of the esophagus.

Figure 2.15. Stratified Squamous Epithelium, Keratinized, Skin of Scrotum, Sheep. The superficial layer of the keratinized stratified squamous epithelium is formed by dead, keratinized cells that appear as thin strands, without visible nuclei, separated from one another. Compare the keratinized cells here with the superficial cells in the nonkeratinized epithelium in Figures 2.13 and 2.14.

Figure 2.16. Stratified Squamous Epithelium, Keratinized, Wattle, Pig. The wattle is covered by a keratinized stratified squamous epithelium.

Figure 2.17. Bistratified Cuboidal Epithelium, Esophagus, Dog. This duct of an esophageal gland is lined by a bistratified cuboidal epithelium.

Helpful Hints:

- The cells of a stratified epithelium will be different shapes at different levels. The specific type of stratified epithelium is named according to the shape of the most superficial (apical or surface) cells. For example, the surface cells of a stratified squamous epithelium are flat.
- The basal layers of a stratified squamous epithelium look darker (especially noticeable at lower magnification) because the deeper cells are smaller with less cytoplasm and, therefore, their dark basophilic nuclei are closely




Figure 2.18 × 250




Figure 2.19 × 250




Figure **2.20** × 125




Figure 2.21 × 125




Figure 2.22 × 125

KEY

- 1. Bistratified columnar epithelium
- 2. Capillary, lumen
- 3. Collagenous fibers
- 4. Columnar cell, nucleus
- 5. Duct, lumen
- 6. Smooth muscle tissue
- 7. Squamous cell, nucleus
 - 8. Stratified columnar epithelium
 - 9. Transitional epithelium
 - 10. Urethra, lumen
 - 11. Urinary bladder, lumen

Figure 2.18. Bistratified Columnar Epithelium, Duct of Carpal Gland, Pig. This duct is lined by a bistratified columnar epithelium. Note the columnar-shaped apical cells (the cells bordering the lumen) with elongated nuclei. A simple squamous epithelium lines the longitudinal section of a capillary that is visible in this field

Figure 2.19. Stratified Columnar Epithelium, Urethra, Goat. This portion of the urethra is lined by a stratified columnar epithelium. The surface cells of this stratified epithelium are columnar.

Figure 2.20. Transitional Epithelium, Unstretched, Urinary Bladder, Pig. A transitional epithelium lines the lumen of the bladder as well as other parts of the urinary system. Surface cells of the transitional epithelium are either balloon-shaped or broadly cuboidal when not under tension.

Figure 2.21. Transitional Epithelium, Unstretched, Urinary Bladder, Cat. Note the large rounded surface cells in this unstretched transitional epithelium.

Figure 2.22. Transitional Epithelium, Stretched, Urinary Bladder, Cat. Surface cells of this epithelium are flattened and elongated when the bladder is stretched with urine.

For additional examples of stratified epithelia, see these figures:

Bistratified cuboidal: 12.58 **Bistratified columnar**: 12.58

Stratified squamous, nonkeratinized: 15.9, 15.47,

19.15

Stratified squamous, keratinized: 12.3, 12.14,

12 16 12 24

Transitional: 14.20, 14.24, 14.26

CONNECTIVE TISSUE PROPER AND EMBRYONIC CONNECTIVE TISSUE

f the four main types of tissues (epithelial, connective, muscle, and nervous), connective tissue is the most abundant and includes numerous varieties. Embryonic connective tissue, connective tissue proper, and the special kinds of connective tissue are distinguished from one another by the types and numbers of cells present and the nature of the extracellular matrix. Certain cells, such as fibroblasts, are common in connective tissue proper and may also be present in some of the special types of connective tissue. Other cells are unique to one kind, such as osteocytes in bone, chondrocytes in cartilage, and erythrocytes in blood. The matrix is semifluid in connective tissue proper; it is gel-like in cartilage, hard in bone, and fluid in blood.

Embryonic connective tissue and connective tissue proper are presented in this chapter. The special types of connective tissue, cartilage, bone, blood, and bone marrow, will be covered in subsequent chapters.

COMPONENTS OF CONNECTIVE TISSUE PROPER AND EMBRYONIC CONNECTIVE TISSUE

Connective tissue binds together and supports other tissues. It is a composite of various cells and fibers in an amorphous ground substance. The latter two components comprise the extracellular matrix, which typically predominates over the cellular elements.

Extracellular Matrix

Ground Substance

The ground substance, composed largely of glycoproteins and glycosaminoglycans, forms a well-hydrated gel that fills the spaces between cells, fibers, and vessels of

connective tissue. It acts as a reservoir for interstitial fluid, providing a medium through which oxygen, nutrients, and metabolic by-products diffuse to and from cells of various tissues and the vascular system.

Fibers

Collagenous, reticular, and elastic fibers occur in connective tissue. Collagenous fibers, comprised of the fibrous protein collagen, are generally the most abundant. They are strong and flexible, yet able to resist stretch. They may be fine or coarse, and they are characteristically unbranched and somewhat wavy. In tissues stained with H&E, they appear pink and shiny.

Reticular fibers are also formed from the protein collagen. They are delicate, branching fibers that possess a coat of glycoproteins and proteoglycans. They are argyrophilic (silver-loving) and can be stained with silver to distinguish them from other fibers of the connective tissue. They may also be selectively stained with Schiff's reagent.

Elastic fibers, formed from the protein elastin, range in diameter from fine to coarse and ordinarily cannot easily be distinguished from collagenous fibers without the use of special stains such as orcein or Weigert's resorcin fuchsin. In some H&E preparations, however, they become colored more intensely by eosin than the collagenous fibers.

Cells

The types and numbers of cells vary with the specific type of connective tissue. Fibroblasts are generally the most numerous of the cells found in connective tissue proper. They are responsible for the formation of fibers and ground substance. Macrophages (histiocytes), derivatives of monocytes of the blood, are also common. They are phagocytic cells that can be recognized, sometimes, by the presence of debris in their cytoplasm, which gives them a dirty appearance. Other migrants from the blood that are found in connective tissue are neutrophils, eosinophils, and lymphocytes. Plasma cells, adipocytes, mast cells, and globular leukocytes also occur in varying numbers in connective tissue.

TYPES OF EMBRYONIC CONNECTIVE TISSUE

Mesenchyme tissue is found in the embryo. It consists of a loose arrangement of pale, star-shaped (stellate) cells with interconnecting cytoplasmic processes. The mesenchyme cells are embedded in a jelly-like, amorphous, ground substance that accumulates fine fibers as development progresses.

Mucous connective tissue, another type of embryonic connective tissue, surrounds the vessels of the umbilical cord. It also occurs in limited regions in adult animals, e.g., the dermis of the comb and wattle of the chicken. It is comprised of fibroblasts and loosely arranged, fine, collagenous fibers in an abundant, amorphous ground substance.

TYPES OF CONNECTIVE TISSUE PROPER

The categories of connective tissue proper are classified according to the arrangement and proportions of their cellular and extracellular components: loose, dense, reticular, elastic, and adipose tissue.

In loose (areolar) connective tissue, the ground substance predominates. It contains many scattered cells of various types, vessels, and a loose network of fine collagenous, reticular, and elastic fibers. Loose connective tissue is widespread throughout the body. It surrounds vessels and nerves, and it is found in serous membranes such as mesenteries, the lamina propria of mucous membranes, subcutaneous tissue, the papillary (superficial) layer of the dermis, and various other places.

In contrast to loose connective tissue, dense connective tissue (often called fibrous tissue) is comprised principally of thick, collagenous fibers. It contains fewer cells than loose connective tissue, most of which are fibroblasts. In dense irregular connective tissue, the collagenous fibers course in all directions, forming a compact three-dimensional meshwork. Dense irregular connective tissue occurs in such places as the reticular (deep) layer of the dermis, submucosa of the digestive tract of some species, and capsules of organs. Dense regular connective tissue is characterized by closely packed, parallel bundles of collagenous fibers. Tendons, ligaments, and aponeuroses are formed by dense regular connective tissue.

It is helpful to know that there are no sharp lines of distinction between loose and dense irregular connective tissue, or between dense irregular and regular connective tissue. Therefore, it is not always possible to classify these types of connective tissues with great precision.

Reticular tissue is comprised of numerous reticular fibers. It forms a supportive network for the parenchyma of structures such as the spleen, lymph node, liver, kidney, and bone marrow.

Types of Connective Tissue

The numerous types of connective tissues are categorized in various ways; we will follow this format:

Embryonic Connective Tissue

Mesenchyme

Mucous

Connective Tissue Proper

Loose (Areolar)

Dense

Reticular

Elastic

Adipose

Cartilage

Bone

Blood


Bone Marrow


The term "connective tissue" used alone typically refers to the general types, namely loose and dense.


Elastic tissue is characterized by numerous regularly or irregularly arranged elastic fibers. It is exemplified by the ligamentum nuchae of grazing animals and by the vocal ligaments.

Adipose tissue consists of groups of adipocytes (also called adipose cells or fat cells) within the loose connective tissue of such places as mesenteries, subcutis, and sheaths of vessels and nerves.

| | Word Roots | | |
|-------|---------------------------|--|--|
| ROOT | MEANING | EXAMPLE SENTENCE | |
| Adipo | Fat | Adipose tissue is also called fat tissue. | |
| | A little open space | There are numerous spaces filled with ground substance between the cells and fibers of loose, or <i>areolar</i> , connective tissue. | |
| Fibro | A fiber | Fibroblasts produce fibers and ground substance of connective tissue. | |
| Phage | Eat | Macrophages, whose name means "big eaters," are important phagocytic cells. | |
| | A net or network | The <i>reticular</i> fibers of <i>reticular</i> tissue are arranged in a crisscrossing, netlike arrangement. | |










Figure 3.4 × 625




Figure 3.5 × 625

1. Amorphous ground substance 7. Fibroblast nucleus 2. Collagenous fiber 8. Lymphocyte 3. Elastic fiber 9. Mast cell 4. Eosinophil 10. Mesenchyme cell 5. Epithelium, lip 11. Neutrophil

12. Plasma cell

6. Erythrocytes in capillary

Figure 3.1. Mesenchyme,72-Hour Embryo, Chicken. Mesenchyme consists of stellate cells. Their processes touch, forming a three-dimensional latticework. The cells are surrounded by an amorphous ground substance.

Figure 3.2. Mucous Connective Tissue, Umbilical Cord, Cow. Mucous connective tissue consists of a loose framework of fibroblasts and collagenous fibers in an amorphous ground substance. Mucous connective tissue of the umbilical cord is often called Wharton's jelly.

Figure 3.3. Loose Connective Tissue, Mesentery, Cat (LeukoStat and orcein). The loose arrangement of the connective tissue cells and fibers in this whole mount preparation is evident. Fine, branching elastic fibers appear blue-gray. The thicker, collagenous fibers stain pale pink. Note the mast cell filled with purple granules.

Figure 3.4. Plasma Cells, Loose Connective Tissue, Lamina Propria, Jejunum, Dog. Plasma cells are common constituents of the lamina propria of the gastrointestinal tract. They are characterized by a basophilic cytoplasm and large blocks of nuclear heterochromatin. A lightly stained area adjacent to the usually eccentric nucleus marks the location of the Golgi apparatus.

Figure 3.5. Mast Cells, Loose Connective Tissue, Lip, Cat (Toluidine Blue). The granules of mast cells are metachromatic and are colored purple by toluidine blue.




Figure 3.6

For other labeled examples of cells of connective tissue, see these figures:


Eosinophils: 13.102, 13.105 Fibroblasts: 8.5, 8.6 Lymphocytes: 11.4, 11.31, 11.32 Macrophages: 10.4 Mast cells: 10.2 Plasma cells: 10.2

| | KEY |
|---|---|
| Collagenous fiber Eosinophil Fibroblast nucleus Globular leukocyte | 5. Lymphocyte 6. Mast cell 7. Plasma cell |

Figure 3.6. Loose Connective Tissue, Lamina Propria, Duodenum, Cow. A loose meshwork of fibers of the connective tissue and various cells are contained in an amorphous, ground substance.



 \times 250



 \times 180 Figure 3.8




Figure 3.9 \times 62.5




Figure 3.10 \times 62.5

KEY

- 1. Adipocyte
- 2. Arteriole
- 3. Collagenous fiber
- 4. Dense irregular connective tissue 10. Tendon, l.s.
- 5. Eosinophil
- 6. Epithelium, planum
- 7. Fibroblast nucleus
- 8. Loose connective tissue
- 9. Macrophage
- 11. Tendon, x.s.

Figure 3.7. Macrophages, Loose Connective Tissue, Colon, Pig.

Wandering macrophages are characterized by their oval shape. The cytoplasm of these cells often contains ingested particles and appears dirty. Eosinophils of the pig contain oval or bilobed nuclei.

Figure 3.8. Loose and Dense Irregular Connective Tissue, Dermis, Planum Nasolabiale, Cow. Note that the loose connective tissue of the papillary layer of the dermis contains finer fibers and more cells than the dense irregular connective tissue of the reticular layer.


Figure 3.9. Dense Irregular Connective Tissue, Dermis, Horse. Note the coarse, interwoven, collagenous fibers.

Figure 3.10. Dense Regular Connective Tissue, Tendon, x.s. and I.s., Nose, Pig. In tendons and ligaments, collagenous fibers are arranged in parallel order. Fibroblasts are located between the fibers.

Helpful Hints

To Distinguish Between Loose and Connective Tissue (CT):

- The main characteristics of loose CT, when contrasted with dense CT, are that loose CT has:
 - 1) more cells, and a great variety of cell types
 - 2) fewer and thinner collagenous fibers
- The main characteristics of dense CT, when contrasted with loose CT, are that dense CT has:
 - 1) fewer cells, most of which are fibroblasts
 - 2) more numerous and thicker collagenous fibers
- What about the amount of space? In loose CT the components have more space between them, and in dense CT the cells and fibers are more closely packed together. However, the amount of space may be deceiving in histologic preparations, because separation artifact (artificial space) may result due to shrinkage of the specimen that occurs during processing.
- Remember that, although elastic and reticular fibers are present in many preparations of connective tissue, typically special strains are required to distinguish them.








Figure 3.12 × 62.5

Figure 3.13 × 62.5




Figure 3.14 × 25




Figure 3.15 × 62.5

| | KEY |
|--|--|
| Adipocyte Elastic fiber Fibroblast nucleus Lymphocyte | 5. Reticular fiber 6. Tendon, x.s. 7. Tendon sheath, inner 8. Tendon sheath, outer |

Figure 3.11. Tendon and Tendon Sheath, x.s., Dog. The tendon sheath is actually made up of two sheaths. The inner sheath attaches to the surface of the tendon. The outer sheath forms a tube around the tendon and attaches to peripheral structures. The space between the two sheaths is filled with synovial fluid in living tissue. The space is not lined by an epithelium, but rather by collagenous fibers and cells of the connective tissue of the sheaths.

Figure 3.12. Elastic Tissue, Ligamentum Nuchae, I.s., Sheep. This section shows the parallel arrangement of the elastic fibers within the ligament.

Figure 3.13. Elastic Tissue, Ligamentum Nuchae, x.s., Sheep (orcein). Orcein selectively stains elastic fibers red.

Figure 3.14. Reticular Tissue, Lymph Node, Cow (Silver). Networks of reticular fibers have been blackened by silver.

Figure 3.15. Adipose Tissue, Soft Palate, Cow. Lipid content of each adipocyte (unilocular) was removed during processing, leaving an empty cavity surrounded by a thin rim of cytoplasm. Nuclei occur at the periphery of adipocytes. It is sometimes difficult to distinguish their nuclei from those of other cells of the connective tissue. See Figure 12.103 for an example of multilocular adipocytes.

For other labeled examples, see these figures:

Adipose tissue: 10.28, 10.40, 10.41, 13.72, 13.111, 13.113

Dense irregular connective tissue: 8.6 Dense regular connective tissue: 8.12

Collagenous fibers: 2.11, 2.12, 2.13, 2.17, 2.18, 8.5, 8.6, 8.13, 12.25. There are also numerous other unlabelled

examples of collagenous fibers throughout the atlas. **Elastic fibers**: 4.5, 4.6, 10.13, 10.14, 10.26, 10.27,

12.25, 15.12, 15.14, 15.18, 15.48

Mucous connective tissue: 12.90, 12.91, 12.93,

12.94

Reticular fibers: 11.24, 13.173, 14.19

CARTILAGE

GENERAL FEATURES

artilage is a special type of connective tissue. There are three basic types of cartilage: hyaline cartilage, elastic cartilage, and fibrocartilage (fibrous cartilage). Each consists of cells called chondrocytes embedded in extracellular matrix. The cartilage matrix, formed by amorphous ground substance and fibers, is firm but flexible.

Chondrocytes are often distorted in histologic specimens because they shrink during slide preparation, revealing spaces that the cells occupy, called lacunae.

The matrix of hyaline and elastic cartilage is avascular and usually invested by a **perichondrium** whose inner layer is chondrogenic, containing cells with the capacity to become chondroblasts. Its outer portion, the outer fibrous layer, is dense irregular connective tissue.

TYPES OF CARTILAGE

Hyaline cartilage, the most common type of cartilage, forms large parts of the developing vertebrate skeleton. It is also found in epiphyseal discs, articular cartilages, the trachea, bronchi, and elsewhere. The matrix of hyaline cartilage contains collagenous fibers that are not discernible with routine stains. It is separable into pale and darkly stained areas called interterritorial matrix and territorial matrix, respectively. The higher concentration of sulfated glycosaminoglycans in the latter is responsible for the darker staining. Isogenous groups, small clusters of two or four chondrocytes occupying the same lacuna, may be observed. They are the result of cell division of chondrocytes.

Elastic cartilage is similar in structure to hyaline cartilage. Its name derives from the presence of numerous elastic fibers embedded in the matrix. Among other places, it is found in the epiglottis, parts of the larynx, and the pinna.

Fibrocartilage (fibrous cartilage) is a combination of hyaline cartilage and dense connective tissue (fibrous tissue). Linear groupings of chondrocytes, embedded in a small amount of cartilage matrix, are situated between bundles of collagenous fibers of dense connective tissue. Fibrocartilage is found in such places as intervertebral discs, the pubic symphysis, and cardiac skeleton, as well as within some tendons close to their attachment to bone.

| Word Roots | | |
|------------|-------------------|---|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Chondro | Cartilage | Chondrocytes are cells in cartilage. |
| Hyal | Glassy | Fresh <i>hyaline</i> cartilage has a shiny, glassy appearance. |
| Lacuna | A space or cavity | Chondrocytes occupy spaces in the matrix called <i>lacunae</i> . |
| Peri | Around | The <i>perichondrium</i> is connective tissue that surrounds hyaline and elastic cartilage. |




Figure 4.1 × 62.5




Figure 4.2 × 125




Figure 4.3 × 125




Figure 4.4 × 62.5




Figure 4.5 × 250

| | KEY |
|--|---|
| Chondrocyte Dermis Elastic cartilage Elastic fiber Glands Interterritorial matrix Isogenous group Lacuna | Perichondrium Perichondrium, chondrogenic layer Perichondrium, fibrous layer Pseudostratified columnar epithelium Separation artifact Territorial matrix Trachea, lumen Area shown in Figure 4.5 |
| | |

Figure 4.1. Hyaline Cartilage, Trachea, Cat. Hyaline cartilage supports the wall of the trachea. Numerous chondrocytes in lacunae are evident in the cartilage. The perichondrium that surrounds the cartilage consists of an outer fibrous layer and an inner chondrogenic layer. The fibrous layer is made of mainly collagenous fibers and fibroblasts. The chondrogenic layer contains precursors of chondrocytes that secrete cartilage matrix.

Figure 4.2. Hyaline Cartilage, Trachea, Cow. The chondrocytes have dark basophilic nuclei surrounded by hazy cytoplasm. Those at the periphery of the cartilage are smaller and flattened; the more central chondrocytes are larger. Numerous isogenous groups of chondrocytes are apparent. Lacunae, evident as white spaces bordering the chondrocytes, are surrounded by dark territorial matrix. The paler interterritorial matrix is more distant from the lacunae. The collagenous fibers in the matrix of hyaline cartilage are not apparent with routine stains.


Figure 4.3. Hyaline Cartilage, Bronchus, Cat. Note the same features as in Figure 4.2.

Figure 4.4. Elastic Cartilage, Pinna, Dog (Weigert's). The pinna (auricle) of the ear consists of elastic cartilage covered by skin. The rectangle outlines the approximate area magnified in Figure 4.5.

Figure 4.5. Elastic Cartilage, Pinna, Dog (Weigert's). Numerous darkly stained elastic fibers are distinct in the matrix of the elastic cartilage. Because the chondrocytes are quite distorted in this preparation, more of each lacuna is visible. When cartilage is sectioned, the slice may not pass through each chondrocyte, only the space around it, so that lacunae sometimes look empty.

Helpful Hints

- To find hyaline or elastic cartilage at low power, look for masses that have a somewhat bubbly appearance due to the numerous scattered white spaces, the lacunae, around the chondrocytes.
- Sometimes the cartilage matrix does not stain intensely, and may appear almost white.
- To distinguish between hyaline cartilage and elastic cartilage, look for fibers in the matrix. The elastic fibers can be seen in elastic cartilage. Collagenous fibers are present but not discernible in hyaline cartilage.








Figure 4.7 × 125




Figure 4.8 × 250




Figure 4.9 × 125

| | KEY | |
|--|-------------------------------|--|
| Cartilage matrix Chondrocyte Collagenous fiber | 4. Elastic fiber 5. Lacuna | |

Figure 4.6. Elastic Cartilage, Epiglottis, Dog. In this preparation of elastic cartilage stained with H&E, the elastic fibers visible throughout the matrix are shiny and pink.

Figure 4.7. Fibrocartilage, Intervertebral Disc, Horse. Fibrocartilage is a combination of dense (fibrous) connective tissue and hyaline cartilage. Small chondrocytes in lacunae are arranged in rows and framed by a hazy rim of pale blue cartilage matrix. Lacunae appear as small white halos around the chondrocytes. Collagenous fibers of the dense connective tissue are visible between rows of chondrocytes.

Figure 4.8. Fibrocartilage, Intervertebral Disc, Horse. Note the same features as described above in this magnified view of fibrocartilage.

Figure 4.9. Fibrocartilage, Claw, Chicken. The numbers and arrangements of collagenous fibers and chondrocytes vary in fibrocartilage. Here, rows of chondrocytes in lacunae are randomly scattered among large bundles of collagenous fibers oriented in various directions.

For additional examples of cartilage, see these figures:

Hyaline cartilage: 15.16, 15.20 to 15.22, 15.43, 19.26,

19.27

Elastic cartilage: 15.7, 15.8, 15.10 **Fibrocartilage**: 10.35, 17.48

BONE TISSUE

bone, for example the temporal bone or humerus, is an organ, as it is formed by several types of tissues, including bone tissue, bone marrow, dense connective tissue, and others. **Periosteum** is dense connective tissue found on the outer surfaces of bones (except where articular cartilage is present), while epithelial cells of the **endosteum** line the bone surfaces of the **marrow** (**medullary**) **cavity** on the inside.

BONE MATRIX

Bone tissue is a living, dynamic connective tissue. It is admirably suited for its function as a skeletal substance because of its high tensile strength and relatively light weight. Its hardness and strength are provided by a matrix consisting of an amorphous ground substance, collagenous fibers that provide some flexibility, and crystals of hydroxyapatites (salts of calcium and phosphate) that make it hard.

CELLS OF BONE TISSUE

Cells of bone tissue are derivatives of **osteoprogenitor** (**osteogenic**) **stem cells** that arise from cells in the embryonic mesenchyme and have the capacity to divide and give rise to osteoblasts.

Osteoblasts are basophilic cells found on the surface of bone where they deposit the organic components (ground substance and fibers) of bone matrix. The unmineralized matrix, osteoid, subsequently becomes mineralized. Bone matrix is typically deposited as multiple layers, or lamellae.

As osteoblasts become surrounded by bone matrix, they mature into flat cells called **osteocytes** that possess numerous processes. Each osteocyte is confined to a space within

the matrix called a **lacuna** that contains tissue fluid, as do **canaliculi**, which are tiny channels in the matrix that contain the processes of the osteocytes. The networks of interconnected lacunae and canaliculi are associated with blood vessels, and this relationship allows the diffusion of nutrients and wastes between the blood and osteocytes, which are otherwise enclosed in hard bone matrix. Osteocytes function to maintain the integrity of the matrix.

Osteoclasts are large, multinucleated cells that border bone surfaces. They resorb bone matrix as necessary during the processes of bone development, remodeling, and repair.

TYPES OF BONE

There are two types of bone, spongy (cancellous) bone and compact (dense or lamellar) bone, distinguished by the amount of space (containing loose connective tissue and blood vessels) that is present. In general, compact bone has more bone matrix than space; spongy bone possesses more space than bone matrix. Flat bones of the skull are formed of spongy bone situated between two layers of compact bone. In long bones, compact bone forms the outer shells of the diaphysis and epiphysis, while spongy bone occurs in the interior of the epiphysis and the endosteal surface of portions of the diaphysis.

Spongy bone is formed by an interconnected meshwork of small pieces called spicules or trabeculae. It is called spongy bone because of the numerous medullary spaces between the delicate three-dimensional latticework of bony spicules.

Compact bone is formed by densely packed layers of bone matrix with osteocytes sandwiched between the layers. In the diaphysis of a long bone, lamellae encircle the inner and outer circumference of the bone as circumferential lamellae; other lamellae, called concentric lamellae, form ring-like layers arranged as cylinders called Haversian systems (osteons). Osteons are oriented parallel to the long axis of a bone and in the center of each one is a longitudinal Haversian canal (central canal), which contains blood vessels and nerves. The central canal is encircled by the concentric lamellae with osteocytes in lacunae. Volkmann's canals (perforating canals) are channels that contain blood vessels and nerves, which connect with vessels and nerves of the periosteum, central canals, and the medullary cavity.

BONE FORMATION

Bone tissue develops either by intramembranous or endochondral ossification. In intramembranous ossification, osteoblasts directly deposit bone matrix in or beneath a membrane. The membrane is either mesenchymal, as in the development of a flat bone of the skull, or periosteal, as in growth in diameter of a long bone.

During endochondral (intracartilaginous) ossification, bone tissue replaces hyaline cartilage. An important aspect of the growth in length of long bones is the persistence of functional epiphyseal discs. These plates of hyaline cartilage located at the junction of the epiphysis and diaphysis permit the process of endochondral ossification to continue until full growth of the bone is achieved, at which time the discs become completely replaced by bone and no further lengthening is possible.

The structure of bone is unrelated to its mode of development; that is, the lamellae formed by intramembranous ossification have the same basic makeup as those that result from endochondral bone formation. Mature bone, however, contains fewer osteocytes than the immature bone it replaces. The woven form of the latter contains numerous osteocytes and an organic matrix of interlacing collagenous fibers. Its matrix has a bluish cast in preparations stained with hematoxylin and eosin. In contrast, the matrix of mature bone is uniformly acidophilic.

Bone matrix undergoes remarkable transformations in size and shape during development. This process of bone remodeling is especially well exemplified during the formation of the skull and long bones. In both instances, transformations in shape and increases in size are accomplished through the processes of bone deposition and bone resorption.

| Word Roots | | |
|------------|-------------------|---|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Cancell | Latticework | A latticework of bits of bone characterize spongy, or <i>cancellous</i> , bone. |
| Clast | Break | Osteoclasts break down, or resorb, bone matrix. |
| Endo | Within, inner | In <i>endochondral</i> bone formation, bone forms within a hyaline cartilage model. |
| Intra | Within, inside | Intramembranous bone formation occurs within a membrane, such as mesenchyme. |
| Os, osteo | A bone | Osteoblasts produce unmineralized bone |




Figure 5.1





Figure 5.2



KEY

- 1. Bone matrix
- 2. Howship's lacuna
- 3. Immature bone
- 4. Lacuna
- 5. Mature bone
- 6. Osteoblast
- 7. Osteoclast
- 8. Osteocyte
- 9. Osteoid
- 10. Separation artifact

Figure 5.1. Intramembranous Ossification, Spongy Bone, Decalcified, Nose, Dog. Intramembranous ossification begins in centers of ossification in the embryo, where osteoblasts deposit bone matrix directly within mesenchyme, producing tiny spicules of bone matrix that interconnect with each other to form a latticework of spongy bone. Osteocytes, derived from osteoblasts that become surrounded by bone matrix, can be seen in the forming spicules. Because bone matrix is deposited without a preformed cartilage model during intramembranous ossification, the connected spicules of bone lack cores of calcified cartilage.

Figure 5.2. Intramembranous Ossification, Spongy Bone, Decalcified, Nose, Dog (Masson's). Osteoclasts are multinucleated giant cells present on the surfaces of bone where resorption is occurring. Depressions that form as osteoclasts erode bone matrix are called Howship's lacunae. The spaces between the surface of the bone matrix and the cells that border it are separation artifacts.

5.3. Immature Phalanx, **Figure** Bone, Decalcified, Fetus, Horse. There are more osteocytes per unit area in immature bone than in mature bone. Typically, immature bone shows basophilia. Both characteristics are evident in the micrograph. Note the rather even acidophilia of the matrix of the more mature bone. Newly deposited unmineralized bone matrix, osteoid, abuts the surface of the mineralized bone matrix.




Figure 5.4




Figure 5.5

KEY

- 1. Area shown in Figure 5.6
- 2. Bone matrix
- 3. Calcified cartilage matrix
- 4. Hyaline cartilage
- 5. Hypertrophied chondrocyte
- 6. Marrow cavity
- Periosteal bud
- 8. Periosteal collar
- 9. Periosteum

Figure 5.4. Hyaline Cartilage Model, Developing Long Bone. Endochondral bone formation in the embryo begins with a model of hyaline cartilage.

Figure 5.5. Primary Center of Ossification, Developing Long Bone. In this sketch, a primary center of ossification has appeared in the center of the cartilage model. The area of the primary center that is outlined (1) is shown in Figure 5.6. A bony periosteal collar has been deposited (by way of intramembranous ossification) around the mid part of the hyaline cartilage model by osteoblasts that formed from cells of the perichondrium, and the perichondrium is now the periosteum. Blood vessels of a periosteal bud have invaded the primary center of ossification, introducing cells that form the bone marrow and give rise to osteoblasts.

Figure 5.6. Primary Center of Ossification, Phalanx, I.s., Decalcified, Fetus, Horse. This section was taken from the central region of a developing phalanx and shows early endochondral ossification. The primary center of ossification in the middle of the hyaline cartilage model is characterized by the presence of chondrocytes that hypertrophy (enlarge) as the cartilage matrix around them becomes reduced. The reduced cartilage matrix consists of tiny pieces with various shapes that become calcified. The enlarged chondrocytes die, leaving spaces that merge and fill with bone marrow. Osteoblasts introduced by blood vessels of the periosteal bud deposit bone matrix (pink) on the meshwork of calcified cartilage matrix (lavender).




Figure 5.6 \times 45




Figure 5.7




Figure 5.8 \times 12.5



- 1. Area shown in Figure 5.8
- 2. Area shown in Figure 5.9
- 3. Area shown in Figure 5.10
- 4. Blood vessel
- 5. Bone marrow
- 6. Bony spicule in diaphysis 7. Chondrocyte in lacuna
- 8. Hyaline cartilage in epiphysis

- 9. Hypertrophied chondrocyte
- 10. Marrow cavity
- 11. Periosteal collar
- 12. Periosteum
- 13. Zone of hypertrophy
- 14. Zone of multiplication
- 15. Zone of reserve cartilage

Figure 5.7. Developing Long Bone, I.s. This sketch depicts the stage in development of a long bone in which the hyaline cartilage of the diaphysis has been replaced by bony spicules and bone marrow. The epiphyses are still hyaline cartilage. The area outlined with a rectangle (1) approximates that shown in Figure

Figure 5.8. Developing Long Bone, I.s., Decalcified, Dog. In this developing long bone, the epiphyses are still formed of hyaline cartilage, although the early appearance of secondary centers of ossification is indicated by the presence of blood vessels within the cartilage. Chondrocytes close to the epiphyses have become organized, so that the zones of endochondral ossification are present. Areas similar to those outlined by the rectangles in this image are shown in detail in Figures 5.9 and 5.10.

Figure 5.9. Developing Long Bone, I.s., Decalcified, Dog. This magnified view of an area similar to that indicated (2) in Figure 5.8 shows some of the zones of endochondral ossification. The zone of reserve cartilage consists of small chondrocytes. Small, somewhat flattened chondrocytes arranged in stacks mark the zone of multiplication (zone of cell division or proliferation), in which chondrocytes multiply, enabling continued growth of the cartilage. In the zone of hypertrophy, chondrocytes gradually enlarge and the matrix between them becomes reduced.




Figure 5.9 \times 180




Figure 5.10 × 180

KEY

- 1. Bone matrix
- 2. Bony spicule in diaphysis3. Calcified cartilage matrix
- 4. Hypertrophied chondrocyte5. Marrow cavity
- 6. Megakaryocyte
- 7. Osteoblast

- 8. Osteoclast
- 9. Osteocyte
- 10. Sinusoid
- 11. Zone of calcification 12. Zone of hypertrophy
- 13. Zone of ossification

Figure 5.10. Developing Long Bone, l.s., Decalcified, Dog. This is a magnified view of an area (3) similar to that outlined in Figure 5.8. In the zone of hypertrophy, chondrocytes enlarge. They eventually die, leaving spaces that merge and become occupied by bone marrow. In the zone of calcification, strands of cartilage matrix persist between hypertrophied chondrocytes and become calcified. Osteoblasts from the marrow cavity deposit bone matrix (pink) on the remnants of calcified cartilage matrix (lavender to purple) in the zone of ossification. Megakaryocytes, which produce platelets in the bone marrow, are large cells possessing a large, lobed nucleus. Osteoclasts are also present.




Figure 5.11




Figure 5.12

KEY

- 1. Area shown in Figure 5.12
- 2. Articular cartilage
- 3. Bone matrix
- 4. Calcified cartilage matrix
- 5. Epiphyseal disc
- 6. Marrow cavity
- 7. Osteoblast
- 8. Osteocyte

- 9. Periosteum
- 10. Spongy bone, diaphysis
- 11. Spongy bone, epiphysis
- 12. Zone of calcification 13. Zone of hypertrophy
- 14. Zone of multiplication
- 15. Zone of ossification
- 16. Zone of reserve cartilage

Figure 5.11. Epiphyseal Disc, Developing Long Bone, I.s. In the stage of a developing long bone represented in this sketch, most of the hyaline cartilage of the epiphyses has been replaced by spongy bone tissue, and hyaline cartilage remains only on the articular surfaces and in the epiphyseal disc (epiphyseal plate). The epiphyseal disc lies between the spongy bone of the epiphysis and the diaphysis. The area outlined by the rectangle approximates that shown in Figure 5.12.

Figure 5.12. Epiphyseal Disc, Humerus, I.s., Decalcified, Cat. This image of the area outlined in Figure 5.11 shows the cartilaginous epiphyseal disc (epiphyseal plate) between the spongy bone of the epiphysis and the diaphysis. Detail of this epiphyseal disc is portrayed in Figure 5.13.

Figure 5.13. Epiphyseal Disc, Humerus, I.s., Decalcified, Cat. The various zones of endochondral bone formation can be identified. Small, scattered cartilage cells comprise the zone of reserve (resting) cartilage. They proliferate, forming rows that constitute the zone of multiplication (zone of proliferation). The chondrocytes then enlarge (zone of hypertrophy) and the remaining cartilage matrix between the hypertrophied chondrocytes becomes impregnated with calcium salts (zone of calcification). Osteoblasts deposit bone matrix (pink) onto the calcified cartilage matrix (lavender) in the zone of ossification.




Figure 5.13 $\times 130$




Figure 5.14 × 62.5




Figure 5.15 × 62.5



Figure 5.16 × 125




Figure 5.17 × 280




Figure 5.18 × 125

| rigure 5.16 | | × 12: |
|--|---|---|
| | K | EY |
| Bone marrow Bone matrix Calcified carti Canaliculi Haversian car Haversian sys Inner circumfe Interstitial syst Lacuna | ilage matrix nal (Central canal) stem (Osteon) erential lamellae | Osteoblast Osteoclast Osteocyte Sinusoid Volkmann's canal (Perforating canal) Zone of calcification Zone of hypertrophy Zone of multiplication Zone of ossification |
| | | |

Figure 5.14. Endochondral Ossification, Phalanx, l.s., Decalcified, Dog. The zones of calcification and ossification are distinct in this preparation. Calcified cartilage matrix stains an intense purple, while bone matrix is strongly eosinophilic.

Figure 5.15. Endochondral Ossification, Epiphysis of Radius, Decalcified, Dog. Numerous spicules of bone (pink) with cores of calcified cartilage matrix (lavender) are depicted.


Figure 5.16. Compact Ground Bone, Femur, x.s., Cat (Unstained). A Haversian canal surrounded by concentric bony lamellae constitutes a Haversian system (osteon). Lacunae with canaliculi (weblike, fine, dark lines) are occupied by osteocytes and their processes, respectively, in living tissue. Volkmann's canals (perforating canals), inner circumferential lamellae, and parts of old Haversian systems, called interstitial systems, are present.


Figure 5.17. Compact Ground Long Bone, x.s., Cat (Unstained). Spider-web-like channels, canaliculi, can be seen radiating from the lacunae.

Figure 5.18. Compact Bone, Humerus, x.s., Decalcified, Chicken. In decalcified bone, hydroxyapatites have been removed, leaving the collagenous portion of the matrix. Blood vessels, osteocytes, and other tissue elements are also left intact. Compare the appearance of the compact bone in this decalcified preparation to the dried ground compact bone in Figure 5.16.


Helpful Hints

- Ground sections of bone are prepared by removing as much soft tissue and organic matter as possible, and allowing the bone to dry. Thin slices of dried bone are cut with a saw, and then ground until thin enough to study with a light microscope. Typically, they are unstained. Only inorganic matrix remains; structures such as lacunae and canals that were occupied by organic matter (cells and other soft tissues) will appear as black or translucent spaces.
 To make out details of ground compact bone, such as the
- To make out details of ground compact bone, such as the canaliculi, try increasing the contrast by closing the iris diaphragm of the microscope.









× 70

Figure 5.22




Figure 5.23 × 280

| 119010 3.20 | /\ 20\ |
|--|---|
| | KEY |
| Artery Endosteum Fibrous layer, periosteum Haversian (central) canal Marrow cavity Megakaryocyte Nerve | 8. Osteocyte 9. Osteogenic layer, periosteum 10. Periosteum 11. Sharpey's fibers 12. Tendon 13. Vein 14. Volkmann's (perforating) canal |
| | |


Figure 5.19. Compact Bone, x.s., Jaw, Decalcified, Dog. Blood vessels and nerves are evident in this cross section of a Haversian canal (central canal), as is the communication of the Haversian canal with a Volkmann's canal (perforating canal).

Figure 5.20. Compact Bone, Rib, I.s., Decalcified, Cat. Portions of Haversian canals are oriented parallel to the long axis of the bone. The collagenous fibers of a tendon extend into the bone as Sharpey's fibers. The large cells in the bone marrow are megakaryocytes.

Figure 5.21. Compact Bone, Rib, I.s., Decalcified, Cat. Flat cells of the endosteum line the marrow cavity.

Figure 5.22. Periosteum of Femur, I.s., Decalcified, Cat. The periosteum parallels the bone of the diaphysis.

Figure 5.23. Periosteum of Femur, l.s., Decalcified, Cat. A magnified view of the periosteum seen in Figure 5.22. The osteogenic layer of the periosteum abuts the bone and is more cellular than the outer fibrous layer of the periosteum.



× 62.5 **Figure 5.24**





Figure 5.25



× 25 Figure 5.26

KEY

- 1. Articulating surface, P2
- 2. Articulating surface, P3
- 3. Bone matrix 4. Cartilage
- 5. Extensor tendon
- 6. Fibrous capsule
- 8. Sharpey's fibers
 9. Synovial fold of synovial membrane
- 10. Tendon

Figure 5.24. Sharpey's Fibers, Phalanx, I.s., Decalcified, Fetus, Sheep (Mallory's). Collagenous fibers of a tendon become embedded within the bone matrix, where they are called Sharpey's fibers.

Figure 5.25. Sharpey's Fibers, Phalanx, I.s., Decalcified, Fetus, Sheep (Mallory's). Same as Figure 5.24, but magnification has been increased to show the relationship between the bone matrix and the collagenous fibers of the tendon.

Figure 5.26. Distal Interphalangeal Joint, I.s., Fetus, Horse. The mid-dorsal aspect of this developing synovial joint is shown. The joint capsule is comprised of a fibrous portion and a wellvascularized, synovial membrane. The latter lines the joint cavity, except for the articulating surfaces. Synovial folds project into the cavity. The outer, more fibrous portion of the capsule is continuous with the future periosteum of the phalanges and blends dorsally with the extensor tendon. The tendon is attached to the developing extensor process of P3. Note that the phalanges have not yet begun to ossify. After ossification, a layer of hyaline cartilage will persist on the articular surfaces.

BLOOD

lood is a special type of connective tissue comprised of formed elements in a fluid matrix. Plasma is the fluid portion, called serum when depleted of fibrinogen. The formed elements include erythrocytes (red blood cells), leukocytes (white blood cells), and platelets (thrombocytes in birds).

Blood cells and platelets are usually examined in stained blood smears (blood films). To make such a preparation, a drop of blood is spread thinly on a glass slide, dried, and stained with a Romanovsky-type stain such as Giemsa or Wright's. One end of the film is usually much thinner than the other end. Details of cell morphology are more visible in the thin film, where the cells are more flattened and less crowded. Blood smears should be scanned with the high-dry objective. With practice, most cells can be differentiated at this magnification. Oil immersion should be reserved for studying specific cells in more detail.

MAMMALS

Erythrocytes

Mature **erythrocytes** are small, anucleate cells uniquely adapted to transport oxygen and carbon dioxide. They are acidophilic and acquire an orange to red color with Romanovsky stains.

The average diameter of erythrocytes in a dried smear varies with the species. The erythrocytes of the dog are largest $(7.0\,\mu)$, while those of the goat are the smallest $(4.1\,\mu)$. Red blood cells from the same animal are all approximately the same size except in the cow, where variation in the size of the erythrocytes (anisocytosis) is not unusual. In most species the cells are disc-shaped, although in young goats they may also be angulated.

Central pallor, resulting from the erythrocyte's biconcave shape, is best defined in the erythrocyte of the dog, but may be seen in other domestic mammals. Red blood cells

sometimes adhere to each other, forming an arrangement resembling a stack of coins, called a **rouleau**. This commonly occurs in the horse and cat. It is rare in ruminants. **Crenated erythrocytes**, characterized by pointed cell margins, are observed most often in pigs.

Various factors influence the appearance of red (and white) blood cells. These include the freshness of the blood sample, the use of an anticoagulant, how quickly the smear was dried, and the thickness of the smear. The occurrence of central pallor, rouleaux, and crenation varies not only with the species, but also with each smear and within different regions of the same smear.

Leukocytes

Leukocytes are basic cellular components of the immune system. They are nucleated cells that are larger and less numerous than erythrocytes. They are classified as either granulocytes (neutrophils, eosinophils, and basophils) or agranulocytes (lymphocytes and monocytes), depending on the presence or absence of specific cytoplasmic granules. Leukocytes tend to accumulate along the edges of a blood smear, so that examples of them, although often distorted, can be found more readily in these regions.

Agranulocytes

Lymphocytes are the predominant leukocytes in ruminants and pigs. The cells range in size from 6 to $15\,\mu$ and are sometimes classified as small, medium, and large. Most of the lymphocytes in carnivores, horses, and pigs are small. Larger cells occur more often in ruminants.

Small lymphocytes have a relatively large, dense, often eccentric nucleus that is round and may be slightly indented. In the cat the nucleus is sometimes deeply indented like a kidney. The nucleus tends to be oval in the pig. Some of the lymphocytes of ruminants are binucleate.

Only a thin rim of cytoplasm may be visible in the small lymphocyte. The cytoplasm is basophilic and may show a lighter region (perinuclear halo) adjacent to the nucleus. At times, nonspecific azurophilic granules may be seen in the cytoplasm of both small and large lymphocytes.

A large lymphocyte has a less dense nucleus and paler, more abundant cytoplasm than a small lymphocyte. The nucleus may be round, oval, or kidney-shaped.

Monocytes are the largest of the leukocytes (15 to $20\,\mu$ in diameter). The nuclear chromatin tends to be diffuse, appearing lacy or sometimes patchy. The shape of the nucleus is highly variable and may be oval, irregular, kidney-shaped, or horseshoe-shaped. In the horse the nucleus is frequently kidney-shaped. In ruminants the nucleus may appear amoeboid and sometimes has a three-pronged or cloverleaf configuration.

The cytoplasm is generally pale gray-blue and may contain dustlike, azurophilic granules. It often contains vacuoles, which give it a foamy appearance. In ruminants the cytoplasm can be more basophilic and either granular or mottled in appearance.

Granulocytes

Neutrophils are the predominant leukocytes in the dog, cat, and horse. The dark-staining nucleus of the mature cell contains very densely packed chromatin. It is long and narrow and may be monolobed or segmented. The nucleus is sometimes coiled, as in the cat, and more often in the pig. When segmented, the lobes may be separated by slight indentations or thin strands of nucleoplasm. The nuclear membrane may appear irregular or tattered from bulging clumps of chromatin. In the horse, the chromatin is so heavily clumped that the nucleus appears very jagged.

The pale gray cytoplasm of the neutrophil contains pink, dustlike, specific granules that may be difficult to resolve with the light microscope. The granules are smallest in the dog, so that the cytoplasm appears nongranular and very faint. Granulation is most pronounced in the sheep and goat. In these animals, larger, more darkly stained granules occur among the finer pink granules.

Band forms of the neutrophil (and other granulocytes) may be encountered in a smear of peripheral blood. The nucleus of these cells looks like a curved or U-shaped band. A band cell can be distinguished from a mature granulocyte by the relatively smooth, rather than ragged, contour of the nucleus. Also, the chromatin of a band cell is less condensed, and the nucleus appears paler and plumper than in the mature granulocyte.

The nucleus of the **eosinophil**, although similar to that of the neutrophil, tends to be less dense and have fewer lobes. In the pig the nucleus is commonly oval or kidney-shaped rather than segmented. C-shaped, monolobed nuclei are common in ruminants.

The cytoplasm of the eosinophil stains pale blue or gray. Specific granules stain various shades of orange, pink, or red with eosin. The granules of the eosinophil of the dog, unlike those of other domestic mammals, are highly variable in size and do not usually fill the cell. Occasionally, small, clear vacuoles also occur in the cytoplasm. In the cat the granules are rod-shaped. Large, round to oblong granules are a striking feature of the eosinophil of the horse. They usually fill the cytoplasm and cause the cell membrane to bulge, so that the eosinophil resembles a raspberry. In the pig, sheep, and goat the granules are small, round to oval, and numerous, often distorting the cell membrane. In cows, the granules are round and intensely stained.

Only a small percentage (0.5% to 3%) of the leukocytes of domestic mammals are basophils. Hence, basophils are not often found in blood smears. The basophil nucleus may be irregular, bilobed, or highly segmented. The granules of basophils vary in size, number, and staining intensity. They are often fairly large, round to oval, and stain reddish purple to dark purple. The granules are a dumbbell or coccoid shape in the pig. The basophil of the cat is much different from that of the other domestic mammals. Its granules are small and lightly stained (dull gray to lavender) in a lavender cytoplasm.

Because the nuclei of the granulocytes exhibit many forms, the cells also are called **polymorphonuclear leukocytes** (polymorphs, PMNs). These terms, however, are sometimes used specifically as synonyms for the neutrophil.

Platelets

Platelets play an important role in hemostasis. Although they are also referred to as thrombocytes, they are not cells in mammals. They are membrane-bound fragments of cytoplasm from large cells called megakaryocytes found in the bone marrow and sometimes the lymph nodes and spleen. Platelets are small and possess central granules. They occur singly or in clusters in smears of peripheral blood.

CHICKEN

Erythrocytes

Mature erythrocytes of the chicken are very different from those of domestic mammals. They are large, elongated, flat cells with an oval nucleus. In dried blood smears from White Leghorn chickens, they range from approximately 9 to $12\,\mu$ long and 6 to $8\,\mu$ wide. Their size varies with the breed and the sex of the bird. The nucleus contains small, uniformly distributed clumps of chromatin. The cytoplasm stains a pale orange to pink color.

Leukocytes

Agranulocytes

Lymphocytes are the most numerous of the leukocytes in the chicken. Their size varies from small to large, as in mammals. The cytoplasm is slightly basophilic and may appear granular or homogeneous. The nucleus is round, sometimes slightly indented, and is usually centrally located. The chromatin occurs in coarse clumps, except in the larger lymphocytes where it is finer.

Monocytes are usually larger than lymphocytes and their nuclear chromatin tends to be more diffuse. Vacuoles are often seen in the cytoplasm.

Granulocytes

Heterophils are the most abundant of the granulocytes. Both heterophils and eosinophils have acidophilic, specific granules. The granules of the heterophil are rod-shaped or spindle-shaped. Their centers sometimes contain a distinctive, ruby red, spherical granule. During staining there may be partial or complete dissolution of the rods, leaving only the more stable, central granule. The granules of the eosinophil are round and pink. The eosinophil's cytoplasm is pale blue, in contrast to the clear cytoplasm of the heterophil. In both of these granulocytes the nucleus is polymorphic. In the eosinophil, the nucleus generally has fewer lobes and also exhibits dense blocks of chromatin clearly separated by lighter areas. This contrasts with the less distinctly clumped chromatin of the heterophil.

The **basophils** of the chicken are much more numerous than in mammals. Their specific granules are deeply basophilic, and the nucleus is usually unlobed and pale.

Thrombocytes

Thrombocytes are nucleated cells, related in function to the platelets of mammals. They are smaller and less elongated than erythrocytes and have a larger, more round nucleus. The pale, dull blue cytoplasm is characterized by one or more small magenta granules and vacuoles.

| Word Roots | | |
|------------|---------|---|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Cren | Notch | Crenated erythrocytes have a margin that is scalloped or notched. |
| Erythro | Red | Red blood cells are called erythrocytes. |
| Leuko | White | Leukocytes are white blood cells. |
| Phil | Love | The cytoplasmic granules of eosinophils are colored by eosin. |
| Rolel | A roll | Erythrocytes that form <i>rouleux</i> become arranged like coins stacked up in a roll of paper. |

Formed Elements of Blood

```
Mammals (covered in this atlas)
    Erythrocytes
    Leukocytes
         Agranulocytes |
              Lymphocytes
              Monocytes
         Granulocytes
              Neutrophils
              Eosinophils
              Basophils
    Platelets
Chicken
    Erythrocytes
    Leukocytes
         Agranulocytes
              Lymphocytes
              Monocytes
         Granulocytés
              Heterophils
              Eosinophils
              Basophils
    Thrombocytes
```




Figure 6.1 ×312




Figure 6.2 ×780




Figure 6.3 ×780




Figure 6.4 ×780




Figure 6.5 ×780

| | KEY |
|---|--|
| 1. Basophil 2. Endothelial cell, nucleus 3. Eosinophil 4. Erythrocyte 5. Leukocyte 6. Lymphocyte | 7. Monocyte 8. Neutrophil 9. Plasma 10. Platelet 11. Smudged eosinophil 12. Stain precipitate |

Figure 6.1. Blood Cells in Histologic Section, Artery, Cat. Biconcave, disc-shaped erythrocytes, leukocytes, and plasma (pale pink) are within the lumen of an artery.

Figure 6.2. Blood, Dog (Giemsa). Monocyte. The cytoplasm of the monocyte is typically vacuolated. The nucleus is frequently oval or U-shaped. Central pallor of the erythrocytes in this preparation is evident.

Figure 6.3. Blood, Dog (Giemsa). Lymphocyte and neutrophil. The cytoplasm of the small lymphocyte is very sparse and the nuclear chromatin is condensed. The mature neutrophil has a polymorphic nucleus. The pale cytoplasm, barely discernible, is characteristic of the neutrophil of the dog.

Figure 6.4. Blood, Dog (Giemsa). Basophil and neutrophil. The basophil has a polymorphic nucleus and coarse dark, basophilic granules of various sizes.

Figure 6.5. Blood, Dog (Giemsa). Eosinophils. In the eosinophil of the dog, the granules vary in size and number. Vacuoles occur in the cytoplasm. The eosinophil on the right is partially smudged.

Helpful Hints for Observing Blood Smears

- First, focus the specimen at low power, which may be a challenge because blood-formed elements are so small. You will mostly notice numerous tiny eosinophilic "dots," which will be the erythrocytes. Change to medium power and locate the basophilic nuclei of the leukocytes. Then place the leukocyte you want to study in the center of the field and magnify it further.
- Examine the parts of the smear where the cells are evenly spread apart from one another; avoid parts of the smear where the cells are crowded together or distorted.
- Leukocytes tend to be dragged along the edges of the slide when the smear is made, so you may find that they are more concentrated along the periphery of the smear.
- Don't try to identify every single leukocyte. Sometimes the cells become distorted or broken open when the smear is made, and the indistinct cell that results is referred to as a "smudged cell."




Figure 6.6 ×312




Figure 6.7 ×780




Figure 6.8 ×780




Figure 6.9 ×780 Figure 6.10 ×780




Figure 6.11 ×780

| KEY | | |
|---|---|--|
| Basophil Eosinophil Erythrocyte Lymphocyte Monocyte | 6. Neutrophil 7. Platelet 8. Rouleau 9. Smudged cell | |

Figure 6.6. Blood, Cat (Giemsa). Eosinophil, neutrophil, lymphocyte, monocyte and platelets. In the cat, platelets may be elongated, or cigar-shaped, as in this preparation.


Figure 6.7. Blood, Cat (Giemsa). Two monocytes. Erythrocytes are stacked in rouleaux.


Figure 6.8. Buffy Coat, Cat (Giemsa). One eosinophil, neutrophil, basophil, and platelets. A buffy coat is the layer above the packed erythrocytes of a centrifuged blood sample that contains mainly leukocytes and platelets. Eosinophils of the cat have pink, rod-shaped granules. The cytoplasm of the basophil contains numerous small, round, lavender granules that are tightly packed and may be difficult to resolve. Distinct red granules are scattered among the lavender granules. The nucleus of the neutrophil in the cat is often coiled, and the cytoplasm is slightly blue-gray with very fine granules that are difficult to see. Platelets vary greatly in size in the cat.


Figure 6.9. Blood, Cat (Giemsa). A basophil and platelets. The nuclei of basophils are remarkably long in the cat.

Figure 6.10. Blood, Cat (Giemsa). An eosinophil, a neutrophil, and platelets.

Figure 6.11. Buffy Coat, Cat (Giemsa). Three eosinophils, one neutrophil, one basophil, three lymphocytes, and platelets.










Figure 6.17 ×780




Figure 6.18 ×780

| | KEY | |
|---|---|--|
| Basophil Eosinophil Erythrocyte Lymphocyte | 5. Monocyte 6. Neutrophil 7. Platelet | |

Figure 6.12. Blood, Horse (Giemsa). Eosinophil, monocyte, two neutrophils, and lymphocyte. In the neutrophil of the horse, the nucleus often appears very jagged.

Figure 6.13. Blood, Horse (Giemsa). Eosinophil. The eosinophil of the horse has characteristic, large, round granules. The surface of the cell looks bumpy where granules are pressed against the plasma membrane, giving the cell a raspberry-like appearance.

Figure 6.14. Blood, Horse (Giemsa). Eosinophil. Note the rouleaux (stacks of erythrocytes), which are common in the horse.

Figure 6.15. Blood, Horse (Giemsa). Neutrophil and lymphocyte.

Figure 6.16. Blood, Horse (Giemsa). Monocyte. A typical monocyte with pale cytoplasm and linearly arranged chromatin.

Figure 6.17. Blood, Horse (Giemsa). Three neutrophils and one monocyte.

Figure 6.18. Buffy Coat, Horse (Giemsa). Basophil and neutrophil. Granules of the basophil are purple and vary in size and shape. The nucleus of the neutrophil has a characteristic, jagged contour.




Figure 6.19 ×780




Figure 6.20 ×780




Figure 6.21 ×780




Figure 6.22 ×780




Figure 6.23 ×780

| | KEY | |
|---|--|--|
| Basophil Basophil granules Eosinophil Erythrocyte Erythrocyte, crenated Lymphocyte | 7. Monocyte 8. Neutrophil 9. Platelet 10. Rouleau 11. Smudged cell | |

Figure 6.19. Blood, Pig (Giemsa). Two lymphocytes and a monocyte.

Figure 6.20. Blood, Pig (Giemsa). Two lymphocytes, one neutrophil, and two eosinophils. The eosinophil of the pig contains numerous pink, round granules that fill the cytoplasm. The nucleus of the eosinophil is not highly segmented. It varies from oval to deeply indented. Note the coiled appearance of the nucleus of the neutrophil, a common feature in pigs and cats. Crenated erythrocytes are commonly seen in blood smears from the pig. Rouleaux also are evident in this field.

Figure 6.21. Blood, Pig (Giemsa). Two lymphocytes and an eosinophil. In pigs, the nucleus of eosinophils varies from oval, as shown in this image, to deeply indented.

Figure 6.22. Blood, Pig (Giemsa). Eosinophil, basophil, lymphocyte, and a smudged cell. The granules of the basophil of the pig are dumbbell or coccoid in shape. Some of the granules have been squeezed from the basophil in this preparation.

Figure 6.23. Blood, Pig (Giemsa). Basophil and two lymphocytes.




Figure 6.24 ×780




Figure 6.25 ×780 Figure 6.26 ×780




Figure 6.27 ×780 Figure 6.28 ×780




Figure 6.29 ×780

| | KEY | |
|---|------------------------------|--|
| Eosinophil Erythrocyte Lymphocyte | 4. Monocyte 5. Neutrophil | |

Figure 6.24. Blood, Cow (Giemsa). Lymphocyte, monocyte, and neutrophil. The ends of the elongated nucleus of the monocyte are partly overlapped, making it appear ringlike at first glance.


Figure 6.25. Blood, Cow (Giemsa). Eosinophil. The red granules of the eosinophil are small, round, and intensely stained in the cow. The nucleus may be lobed, but is usually C-shaped.

Figure 6.26. Blood, Cow (Giemsa). Eosinophil and a lymphocyte.

Figure 6.27. Blood, Cow (Giemsa). Neutrophil.

Figure 6.28. Blood, Cow (Giemsa). Lymphocyte. The large lymphocyte of the cow often shows a deeply indented nucleus. The cytoplasm is granular and vacuolated.

Figure 6.29. Blood, Cow (Giemsa). An eosinophil and two lymphocytes.






Figure 6.35 ×780




Figure 6.36 ×780

| | KEY |
|---|---|
| Azurophilic granules Eosinophil Erythrocyte Lymphocyte | 5. Lymphocyte, binucleate6. Monocyte7. Neutrophil8. Platelet |

Figure 6.30. Blood, Sheep (Giemsa). Lymphocyte with azuro-philic granules.

Figure 6.31. Blood, Sheep (Giemsa). Binucleate lymphocyte. Some of the lymphocytes of ruminants have two nuclei.


Figure 6.32. Blood, Sheep (Giemsa). Monocyte. The nucleus of the monocytes of ruminants may be oval or indented, or have a three-pronged configuration. The cytoplasm is gray-blue and vacuolated and may contain granules.

Figure 6.33. Blood, Sheep (Giemsa). Monocyte. This monocyte has a three-pronged nucleus. Our observations have revealed that some monocytes with similar nuclei also occur in cows and goats.

Figure 6.34. Blood, Sheep (Giemsa). Lymphocyte and neutrophil. The cytoplasm of the neutrophil of sheep and goats contains numerous small and a few large, pink granules. A perinuclear halo is commonly seen around the periphery of the nucleus of lymphocytes.

Figure 6.35. Blood, Sheep (Giemsa). Neutrophil.

Figure 6.36. Blood, Sheep (Giemsa). Eosinophil. The eosinophil of the sheep contains pink, densely packed, oval granules that are uniform in size.








Figure 6.38 ×780 Figure 6.39 ×780




Figure 6.42 ×780

| | KEY |
|--|---|
| Eosinophil Erythrocyte Lymphocyte Lymphocyte, binucleate | 5. Monocyte 6. Neutrophil 7. Platelet |

Figure 6.37. Blood, Goat (Giemsa). Two lymphocytes and a neutrophil. The chromatin of the lymphocytes is in the form of closely apposed clumps. Fine granules are present in the cytoplasm of the neutrophil but are difficult to see.

Figure 6.38. Buffy Coat, Goat (Giemsa). Lymphocyte and neutrophil.

Figure 6.39. Buffy Coat, Goat (Giemsa). Eosinophil. The small, round, acidophilic granules of the eosinophil of the goat almost fill the cytoplasm.

Figure 6.40. Buffy Coat, Goat (Giemsa). Two lymphocytes. Some lymphocytes of the cow, sheep, and goat are binucleate, as is one of the lymphocytes in this image.

Figure 6.41. Buffy Coat, Goat (Giemsa). Monocyte. The cytoplasm of the monocyte is blue and contains vacuoles that are often seen in clusters.

Figure 6.42. Buffy Coat, Goat (Giemsa). Monocyte and lymphocytes. The nucleus of this monocyte appears amoeboid or three-pronged, as occurs in ruminants.




Figure 6.43 ×312




Figure 6.44 ×312




Figure 6.45 ×780




Figure 6.46 ×780

| KEY | | |
|---|--|--|
| Eosinophil Erythrocyte Granulocyte Heterophil | 5. Lymphocyte 6. Monocyte 7. Thrombocyte 8. Vacuole | |

Figure 6.43. Blood Cells in Histologic Section, Chicken. Elongated, nucleated red blood cells and a few granulocytes are shown in the lumen of a blood vessel.

Figure 6.44. Blood, Chicken (Wright-Giemsa). Erythrocytes, leukocytes, and thrombocytes.

Figure 6.45. Blood, Chicken (Wright-Giemsa). Erythrocytes and thrombocytes. An oval, coarsely granular nucleus and a vacuolated cytoplasm with one or more magenta granules characterize the thrombocyte.

Figure 6.46. Blood, Chicken (Wright-Giemsa). Eosinophil, two lymphocytes, and a heterophil. The heterophil has numerous rodshaped granules. Some of these may show a ruby-red, spherical granule at the center. The nuclear chromatin is coarse and densely packed. In contrast, the eosinophil has fewer round, pink granules in a pale blue cytoplasm. The nuclear chromatin is blocklike; the blocks are distinctly separated from each other.





Figure 6.47 ×780



×780 Figure 6.48




Figure 6.49 ×780

1. Basophil

2. Eosinophil 3. Heterophil

- 4. Immature erythrocyte
- 5. Monocyte6. Smudged heterophil

Figure 6.47. Blood, Chicken (Wright-Giemsa). Basophil, heterophil, monocyte, and immature erythrocyte. The basophil is characterized by large numbers of medium-size basophilic granules. Unlike other granulocytes, the nucleus of this cell is usually not lobed. The cytoplasm of the immature erythrocyte is more basophilic and the nucleus is less condensed than in a mature erythrocyte.

Figure 6.48. Blood, Chicken (Wright-Giemsa). Eosinophil and smudged heterophil. The rod-shaped granules of the heterophil and the round granules of the eosinophil are clearly visible.

Figure 6.49. Blood, Chicken (Wright-Giemsa). Monocyte and heterophil. Many of the rod-shaped granules of the heterophil have a ruby red, spherical granule at their center. This is a common characteristic of the heterophil. The monocyte has a pale, basophilic, and vacuolated cytoplasm.

BONE MARROW

MAMMALS

Red bone marrow is highly cellular and is specialized to produce blood cells and platelets. Along with the spleen and liver, it plays a role in hematopoiesis during prenatal development. At the time of birth it is the principal source of blood cells and is found throughout the entire skeleton of the animal. In the adult, red marrow is mostly limited to the sternum, ribs, vertebrae, skull, ilia, and the ends of long bones. Yellow marrow, rich in adipose tissue, occupies the remainder of the skeleton of an adult.

Red marrow consists of extravascular hematopoietic tissue and vascular sinusoids. The hematopoietic tissue is rich in blood cells in various stages of formation. It also contains cells of the connective tissue and is supported by a reticular meshwork. Pluripotent stem cells provide a source of unipotent stem cells committed to the formation of either erythrocytes, granulocytes, agranulocytes, or megakaryocytes. Generally, immature (early) cells of the bone marrow are relatively large, and they have a euchromatic nucleus with nucleoli. As they divide and mature, the cells become smaller, the nucleus becomes more heterochromatic, and the nucleoli disappear. Older cells predominate over the immature forms.

The progression of cell stages, from the morphologically indistinct stem cell to a specific mature blood cell, comprises a cell series (cell line). Most of the cells seen in preparations of the bone marrow belong to either the **erythroid** (red blood cell) **series** or **granulocytic** (myeloid; white blood cell) **series**. The cells of these series are presented in this chapter.

Erythroid Series

The proerythrocyte (rubriblast, proerythroblast) is a large, round cell with a basophilic cytoplasm. The nucleus is large, with finely granular chromatin and a few nucleoli. This

cell undergoes several divisions, giving rise to basophilic erythroblasts (prorubricytes). These cells are round with round nuclei. They are the earliest cells of the erythroid series that can be readily identified in smears. The basophilic erythroblast is somewhat smaller than its precursor and has a deeply basophilic cytoplasm. The nuclear chromatin is more coarsely clumped, and no nucleoli are visible. Basophilic erythroblasts give rise to polychromatophilic erythroblasts (rubricytes), which are smaller cells. The chromatin is more condensed, appearing as blocks separated by light streaks, similar to the chromatin of a plasma cell. The cytoplasm is mottled with pink and blue areas. As hemoglobin synthesis continues and ribosomes diminish, the cytoplasm becomes more pink and less blue. Mitotic division usually ceases in the late polychromatophilic erythroblast stage. Orthochromatophilic erythroblasts (normoblasts, metarubricytes) are characterized by a round, highly condensed, and deeply stained nucleus. Their cytoplasm is distinctly eosinophilic, but may show slight tinges of blue. Eventually, the nucleus is extruded, leaving an anucleate reticulocyte that matures into an erythrocyte.

Granulocytic Series

Myeloblasts are large granulocytic cells with a grainy, basophilic cytoplasm. The round to oval nucleus contains finely dispersed chromatin. Nucleoli may be present. These cells give rise to the promyelocyte, the earliest stage in the development of a granulocyte that can be readily distinguished in smears. This cell contains a relatively large nucleus with nucleoli and chromatin that is beginning to clump. The cytoplasm contains nonspecific azurophilic (magenta) granules. Promyelocytes divide and give rise to myelocytes. The myelocyte is smaller and has an oval, often eccentric, nucleus with more condensed chromatin. Specific granules, characteristic of neutrophils, eosinophils, or basophils, are apparent in the cytoplasm. Late myelocytes lose the capacity to divide. They are known as metamyelocytes when the nucleus becomes indented and more condensed. With further modification the nucleus becomes more elongated in the band-cell stage before eventually assuming the shape found in the mature granulocyte.


Other Cells in Bone Marrow


Megakaryocytes are situated in the extravascular compartment, close to sinusoids, into which they release platelets. They are very large cells with a polymorphic nucleus and a grainy cytoplasm, and they are often seen in preparations of the bone marrow together with a variety of other cells such as plasma cells, adipocytes, and cells in mitosis. Osteoblasts and osteoclasts, which are closely associated with the surface of the bone lining the marrow cavity, may also be encountered in smear preparations of marrow.


CHICKEN

The organization of bone marrow of the chicken is different from that of mammals. Erythropoiesis takes place within the vascular sinusoids, rather than in the extravascular tissue. The immature red blood cells (large cells with a basophilic cytoplasm) are found adjacent to the endothelium of a sinusoid. As division and maturation of these cells progress, the older ones move inward. Thus, mature erythrocytes (with an eosinophilic cytoplasm) accumulate in the center of the vessel. As in mammals, cells of the granulocytic series (heterophils, eosinophils, and basophils) develop in the extravascular spaces of the marrow.

| Word Roo | ots | |
|-------------------|-------------------------------|---|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Hemato Poieses | Blood Make or produce | Hematopoiesis is the formation of blood. |
| Mega Kary | Large or great The nucleus | Megakaryocytes are very large cells with a large nucleus. |
| Meta | After | Metamyelocytes form from myelocytes. |
| Myelo | Marrow | The <i>myeloid</i> series of cells consists of various stages of cells that form granulocytes in bone marrow. |
| Poly Chromato | Many Color | The cytoplasm of <i>polychromatophilic</i> erythroblasts is mottled with pink and blue colors. |
| Pro | Before | Promyelocytes give rise to myelocytes. |
| Rete | A network | Depending on the stain used, some reticulocytes show a network of blue fibers of precipitated RNA. |










Figure 7.4 × 312




Figure 7.5 × 780

| | KEY |
|---|--|
| Azurophilic debris Basophilic erythroblast Basophilic myelocyte Eosinophilic band cell Eosinophilic myelocyte Erythrocyte Erythroid cell, late Granulocytic cell, early | 9. Megakaryocyte 10. Mitotic figure 11. Neutrophilic band cell 12. Orthochromatophilic erythroblast 13. Osteoblast 14. Osteoclast 15. Plasma cell 16. Smudged cell, nucleus |

Figure 7.1. Megakaryocyte, Bone Marrow, Cat (Giemsa). The megakaryocyte is a large cell with a polymorphic nucleus and grainy cytoplasm. Blood platelets are derived from fragments of the cytoplasm. Forces generated during the formation of the smear appear to have separated a segment of the nucleus.

Figure 7.2. Osteoclast, Bone Marrow, Cat (Giemsa). The osteoclast is a large, multinucleate cell formed from fused macrophages. Azurophilic bone debris can be seen in the cytoplasm of this specimen.

Figure 7.3. Osteoblasts, Bone Marrow, Cat (Giemsa). Osteoblasts are characterized by the presence of an eccentric nucleus and basophilic cytoplasm. A perinuclear clear zone, representing the site of the Golgi apparatus, may be visible. In smears these cells often occur in clusters.

Figure 7.4. Bone Marrow, Cat (Giemsa). A variety of different cells of the bone marrow can be identified at this magnification.


Figure 7.5. Bone Marrow, Cat (Giemsa). Basophilic and eosinophilic myelocytes are represented in this field.


Erythroid (Red Blood Cell) Series


Proerythrocyte = rubriblast, proerythroblast
Basophilic erythroblast = prorubricyte
Polychromatophilic erythroblast = rubricyte
Orthochromatophillic erythroblast = normoblast or
metarubricyte
Reticulocyte = polychromatophil


Granulocytic (Myeloid; White Blood Cell) Series

Myeloblast Promyelocyte = progranulocyte Myelocyte Metamyelocyte Band cell









5 4— 6 7— 5— —8

Figure 7.10 × 780

| | KEY |
|--|---|
| Basophilic band cell Basophilic erythroblast Basophilic myelocyte Endothelial cell, nucleus Eosinophil Erythrocyte, early Erythrocyte, mature Heterophil | 9. Neutrophil 10. Neutrophilic band cell 11. Neutrophilic metamyelocyte 12. Neutrophilic metamyelocyte, late 13. Orthochromatophilic erythroblast 14. Polychromatophilic erythroblast 15. Promyelocyte 16. Reticulocyte |

Figure 7.6. Bone Marrow, Cat (Wright-Giemsa). A basophilic myelocyte, segmented eosinophil, and polychromatophilic erythroblast are evident.

Figure 7.7. Bone Marrow, Cat (Wright-Giemsa). Various myeloid developmental stages are shown in this field.

Figure 7.8. Bone Marrow, Cat (Wright-Giemsa). Various erythroid developmental stages are shown in this field.

Figure 7.9. Bone Marrow, Cat (Wright-Giemsa). The largest cell seen in this field is a promyelocyte. Note the presence of numerous magenta, azurophilic granules in its cytoplasm.

Figure 7.10. Bone Marrow, Plastic Section, Chicken (Giemsa). Intravascular developmental stages of erythrocytes are shown. In the extravascular compartment, heterophils and eosinophils can be distinguished.

MUSCLE TISSUE

GENERAL FEATURES

he cytoplasm of muscle cells, specifically called the **sarcoplasm**, contains a unique substructure of threadlike **myofilaments** (formed mainly of the proteins actin and myosin) that provides the cells with the ability to contract. Shortening occurs when the myofilaments slide past one another, resulting in movements that may be voluntary (such as moving bones of the skeleton, the tongue, and the eyes) or involuntary (such as contraction of the chambers of the heart, constriction of blood vessels and bronchioles, or peristalsis of the digestive tract).

Muscle cells of each of the three types of muscle tissue (smooth, skeletal, and cardiac) are elongated and, therefore, are also referred to as muscle fibers. This term can be misleading because the cells are true cellular units, not connective tissue fibers. Muscle cells are typically arranged parallel to one another in sheets or bundles. The endomysium is a layer of loose connective tissue, with numerous fine fibers and capillaries, which encloses and surrounds individual muscle cells.

TYPES OF MUSCLE TISSUE

Smooth muscle tissue is comprised of cells that are involuntary and lack striations. It occurs, for example, in the digestive tract, blood vessels, urinary bladder, bronchi, bronchioles, and the capsules of some organs.

The name "smooth" is derived from the fact that the cells lack cross-striations, so that the cytoplasm looks uniform rather than striped. The cells are relatively small compared to skeletal and cardiac muscle cells. They are spindle-shaped (fusiform) and have an elongated nucleus located about midway between the tapered ends of the cell.

Skeletal muscle tissue consists of cells with cross-striations that are under voluntary control. It mainly forms muscles that attach to and move the skeleton (hence the name

"skeletal"), but it is also found in organs such as the tongue, pharynx, anal canal, and muscles that move the eyes. Cross-striations result from the precise registration of the A, I, H, and M bands of the myofibrils within the cells.

Skeletal muscle cells can be exceptionally long (up to many centimeters), and they are cylindrical and multinucleated. Their nuclei are located peripherally, immediately below the **sarcolemma** (the specific term for the cell membrane of a muscle cell).

Skeletal muscle cells, surrounded by endomysium, are grouped into bundles called fascicles. Each fascicle, in turn, is surrounded by a perimysium of loose connective tissue. To form a muscle, a sheath of dense connective tissue, the epimysium, encloses and binds numerous fascicles.

Cardiac muscle tissue consists of cells that are striated and involuntary. It forms the myocardium of the heart and occurs in the walls of the major vessels that carry blood to and from the heart, including the aorta, pulmonary artery, pulmonary vein, and vena cava. Cross-striations result from the precise registration of sarcomeric bands, as in skeletal muscle tissue. A single, centrally located nucleus occurs in most cardiac muscle cells. Occasionally, however, a cell with two nuclei may be seen. Unlike skeletal or smooth muscle cells, cardiac muscle cells branch and anastomose with one another. Intercalated discs, composed of desmosomes and gap junctions, join cardiac muscle cells end to end and increase the speed of conduction of impulses, enabling them to contract as a unit within the wall of a heart chamber.

Some cells of cardiac muscle tissue are modified and function as a conducting system that helps to coordinate the heartbeat. These modified cells are the functional elements of the sinoatrial node, atrioventricular node, and Purkinje fibers.

| Word Roots | | |
|------------|------------------|--|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Cala | Insert | Intercalated disks join the ends of adjacent cardiac muscle cells. |
| Cardio | The heart | The wall of the heart contains <i>cardiac</i> muscle tissue. |
| Endo | Within, inner | The <i>endomysium</i> is connective tissue situated between individual muscle cells within a fascicle. |
| Fasci | A bundle | Skeletal muscle cells are arranged in bundles called <i>fascicles</i> . |
| Муо | Muscle | The <i>myocardium</i> is the middle, muscular layer of the wall of the heart. |
| Mys | Muscle | The <i>endomysium</i> is loose connective tissue between muscle cells within a muscle fascicle. |
| Peri | Around | Perimysium is connective tissue that surrounds a group of skeletal muscle cells to form a fascicle. |
| Striat | Streaked | muscle cells that are <i>striated</i> appear streaked or striped. |

| Main Histologic Features | | | |
|---|--|---|--|
| Smooth Muscle Cells | Skeletal Muscle Cells | Cardiac Muscle Cells | |
| NonstriatedSpindle-shapedOne central nucleus per cell | Striated Extremely long, cylinder-shaped Multinucleated, peripheral nuclei | Striated Elongated, branching One central nucleus (occasionally two) per cell | |




Figure 8.1 ×2




Figure 8.2 ×180




Figure 8.3 ×250

- 1. Area shown in Figure 8.2
- 2. Ileum, lumen
- Muscularis externa, inner circular layer
- 4. Muscularis externa, outer longitudinal layer
- 5. Separation artifact
- 6. Serosa
- 7. Simple squamous cell, nucleus
- 8. Smooth muscle cell, nucleus, l.s.
- 9. Smooth muscle cell, nucleus, x.s.
- 10. Smooth muscle cell, sarcoplasm
- 11. Villus

Figure 8.1. Smooth Muscle Tissue, Ileum, x.s., Cat. The muscularis externa of the small intestine consists of two layers of smooth muscle tissue arranged at right angles to one another, namely the inner circular layer and outer longitudinal layer. An area similar to the one outlined within a rectangle in this image is shown in detail in Figure 8.2.

Figure 8.2. Smooth Muscle Tissue, Ileum, x.s., Cat. In this cross section, the smooth muscle cells of the inner circular layer of the muscularis externa, which are oriented circularly relative to the lumen of the organ, have been cut longitudinally. The muscle cells of the outer longitudinal layer, which parallel the long axis of the organ, have been cut transversely. This image includes the simple squamous epithelium, the mesothelium, of the serosa that covers the ileum.

Figure 8.3. Smooth Muscle Tissue, Muscularis Externa, Jejunum, x.s., Sheep. This is a magnified view of the layers of the muscularis externa showing smooth muscle tissue in x.s. and I.s. Smooth muscle cells each have a single, elongated nucleus sometimes referred to as cigar-shaped. This shape is apparent in cells that have been sectioned longitudinally. In a group of smooth muscle cells, the cells are oriented in the same direction and fitted closely together with tapered ends of some cells positioned between midportions of other cells. Thus, their nuclei appear staggered in smooth muscle tissue sectioned longitudinally and, because of the snug arrangement of the cells, their boundaries are often not discernable.

The appearance of smooth muscle cells in cross section varies, depending on where they were transected along their length. If a cell was sliced through its mid region, its basophilic nucleus is visible and appears round. If the cell was cut closer to a tapered end, only eosinophilic cytoplasm, not the nucleus, is present and the cell appears smaller in diameter.

There are numerous images that show **smooth muscle tissue** throughout the atlas. See these figures for some additional examples: 10.6 to 10.19; 13.95, 13.96, 13.98, 13.99, 14.24, 14.32.






Figure 8.4



×250 Figure 8.5



 $\times 180$ Figure 8.6

KEY

- 1. Arteriole, lumen
- 2. Artery, lumen
- 3. Collagenous fiber
- 4. Connective tissue
- 5. Crypt of Lieberkühn (intestinal gland)
- 6. Dense irregular connective tissue 7. Fibroblast, nucleus
- 8. Muscularis mucosae
- 9. Nerve fascicle, unmyelinated
- 10. Simple columnar epithelium
- 11. Smooth muscle cell, nucleus, l.s. 12. Smooth muscle cell, nucleus,
- x.s. 13. Smooth muscle cell, sarcoplasm
- 14. Venule

Figure 8.4. Smooth Muscle Tissue, Coronary Artery, x.s., Pig. The smooth muscle tissue in the tunica media of this blood vessel is contracted, as evidenced by the twisted appearance of the nuclei of the smooth muscle cells. Due to the arrangement of myofilaments and the presence of additional elements in the sarcoplasm of smooth muscle cells, the cells coil and bunch up as

they shorten.

Figure 8.5. Smooth Muscle Tissue, Urinary Bladder, Pig. The spindle-shape of individual smooth muscle cells in longitudinal section and their variable appearance in cross section are particularly evident here. An arteriole, with smooth muscle cells in its wall, is present.

Figure 8.6. Smooth Muscle Tissue and Dense Irregular Connective Tissue, Duodenum, Cat. Layers of smooth muscle tissue, cut longitudinally and transversely, of the muscularis mucosa can be distinguished from the dense irregular connective tissue of the submucosa. The collagenous fibers of the connective tissue are shinier and a brighter pink color than the cytoplasm of the muscle cells. The elongated, pale nuclei of the smooth muscle cells in longitudinal section are aligned in the same direction. The dark nuclei of fibroblasts of the connective tissue are more irregularly arranged.

Helpful Hints

Students often encounter difficulty in distinguishing smooth muscle tissue from loose or dense connective tissue in H&E preparations. It is helpful to know that:

- Although collagenous fibers of connective tissue and the cytoplasm of muscle cells and are both eosinophilic, collagenous fibers are a brighter pink and distinctly shiny, whereas smooth muscle tissue tends to present an overall duller, slightly bluish pink color. Focusing up and down with the fine adjustment will help enhance these differences, especially the reflectivity of the collagenous fibers.
- The nuclei of the smooth muscle cells appear to be more orderly in their arrangement (in the same direction) than those of the cells in the loose or dense irregular connective




Figure 8.7 ×90




Figure 8.7. Skeletal Muscle Tissue, x.s. and I.s., Tongue, Cat. The tongue is highly muscular, with bundles of skeletal muscle cells oriented perpendicular to one another. For this reason, the tongue serves as a useful specimen to observe skeletal muscle tissue in both longitudinal and cross sections. Figure 8.8 is a magnified view of skeletal muscle tissue of a tongue.

Figure 8.8. Skeletal Muscle Tissue, x.s. and I.s., Tongue, Horse. Skeletal muscle cells are large and long and possess numerous, peripheral nuclei. Some of the nuclei that are near the skeletal muscle cells belong to cells of the endomysium and perimysium. In skeletal muscle cells cut longitudinally, crossstriations are evident as thin light and dark stripes perpendicular to the long axis of the cell. When cut transversely, the cells appear round to polygonal, and myofibrils are visible as dots packed in the cytoplasm. If a skeletal muscle cell is sectioned near its surface, its nuclei appear to be more centrally located.

KEY

- Capillary, x.s.
 Connective tissue
- 3. Endomysium
- 4. Myofibrils, x.s., in sarcoplasm
- 5. Perimysium
- 6. Separation artifact
- 7. Skeletal muscle cell, l.s.
- Skeletal muscle cell, nucleus Skeletal muscle cell, nucleus,
- superficial cut
- Skeletal muscle cell, oblique cut
- Skeletal muscle cell, x.s.
- 12. Stratified squamous epithelium

Helpful Hints

- Cross-striations are perpendicular to the long axis of a cell, and they are only visible when skeletal and cardiac muscle cells are sectioned longitudinally. Even then they are not apparent in some histologic preparations.
- Parallel, rod-shaped myofibrils that run the length of the cells sometimes separate enough to give individual skeletal and cardiac muscle cells in longitudinal section a filamentous or stringy look, not to be confused with the striations.
- Skeletal muscle cells are too long to be viewed in their entirety from end to end in a microscope field, so that only a portion of each cell is seen in longitudinal section.





Figure 8.9 × 1300

| KEY | |
|--|--|
| 1. A band 2. Endomysium 3. H band 4. I band | 5. Muscle cell, I.s. 6. Myofibrils 7. Z line |

Figure 8.9. Skeletal Muscle Tissue, I.s., Lip, Dog (Masson's). Individual myofibrils that have frayed apart at the cut ends of these two skeletal muscle cells in longitudinal section are distinct. The light and dark bands of the myofibrils line up with those of other myofibrils within each cell, so that each entire cell appears cross-striated.








Figure 8.11 × 625




Figure 8.12 × 62.




Figure 8.13 × 250




Figure 8.14 × 125

| | KEY |
|--|---|
| Adipose tissue Bifurcation Capillary, x.s. Cardiac muscle cell, nucleus, l.s. Collagenous fiber Dense regular connective tissue Endomysium Intercalated disk | 9. Myofibrils, x.s., in sarcoplasm 10. Separation artifact 11. Skeletal muscle cell, l.s. 12. Skeletal muscle cell, nucleus 13. Skeletal muscle cell, oblique section 14. Skeletal muscle cell, x.s. 15. Smooth muscle cell, nucleus 16. Smooth muscle cell, sarcoplasm |

Figure 8.10. Skeletal Muscle Tissue, x.s., Tongue, Horse. In cross section, the skeletal muscle cells are round to polygonal. Myofibrils are packed in the cytoplasm and peripheral nuclei are evident. Separation artifact between skeletal muscle cells makes the individual cells more distinct and reveals the capillaries and loose connective tissue of the endomysium.

Figure 8.11. Skeletal Muscle Tissue, x.s., Tongue, Horse. Myofibrils are eosinophilic rod-like structures that extend the length of the muscle cell and are formed by myofilaments. In skeletal muscle cells that are cut in cross section, their myofibrils are also cut in cross section and appear as numerous dot-like structures packed in the sarcoplasm. They are especially evident in this preparation.

Figure 8.12. Skeletal Muscle and Tendon, Oblique Cut, Tongue, Horse. The collagenous fibers of the dense regular connective tissue of a tendon can be seen associated with skeletal muscle cells

Figure 8.13. Skeletal and Smooth Muscle Tissue, Sheep (Masson's). The spindle shape of the individual smooth muscle cells cut longitudinally is evident. Cross-striations of portions of skeletal muscle cells in longitudinal section are also visible. With the connective tissue stain used to color this preparation, collagenous fibers are blue/green while the cytoplasm of the muscle cells is magenta (purplish red).

Figure 8.14. Cardiac Muscle Tissue, I.s., Heart, Cat. Branching (bifurcation) of the cardiac muscle cells is visible and endomysium is discernible where separation artifact has occurred.

For more examples, see these Figures:

Skeletal muscle tissue: 10.1, 12.13, 13.22, 15.42

Both smooth and skeletal muscle tissue: 10.8,

13.36, 13.42, 19.32

Cardiac muscle tissue: 10.36




Figure 8.15 × 360




Figure 8.16 × 180




Figure 8.17 × 625

- Capillary, x.s.
 Cardiac muscle
- 2. Cardiac muscle cell, nucleus, l.s.
- 3. Cardiac muscle cell, nucleus, x.s.
- 4. Cardiac muscle cell, x.s.
- 5. Endomysium
- 6. Fibroblast, nucleus
- 7. Intercalated disk
- 8. Myofibrils, x.s.,
- in sarcoplasm 9. Separation artifact

Figure 8.15. Cardiac Muscle Tissue, I.s., Heart, Cat. Branching and cross-striations are apparent in this preparation. Intercalated discs appear as dark, somewhat irregular lines perpendicular to the long axis of the cells and are thicker than cross-striations. (Intercalated discs are not always apparent in H&E preparations.) Nuclei of cardiac muscle cells are centrally located. Nuclei of cells of the endomysium between cardiac muscle cells are also visible.

Figure 8.16. Cardiac Muscle Tissue, x.s., Heart, Pig. Separation artifact exposes the boundaries of the cardiac muscle cells cut in cross section and the location of the endomysium between them. In many cardiac muscle cells, their centrally located nucleus is visible; in others it was not included in the plane of the section. Cross sections vary in shape and diameter, depending on the plane of section through the branching cells.

Figure 8.17. Cardiac Muscle Tissue, x.s., Heart, Pig. In cardiac muscle cells, the myofibrils separate around the centrally located nucleus, forming a perinuclear halo of "clear" cytoplasm free of myofibrils.

NERVOUS SYSTEM

ne can hardly examine a histologic preparation of any sort without finding some evidence of nervous tissue, for example, sections through myelinated or unmyelinated nerves, isolated neurons, encapsulated nerve endings, or perhaps a nerve plexus.

Nervous tissue consists of various sizes and kinds of neurons (nerve cells) and their supporting elements, and it is basically similar in all domestic animals. Neurons each have a cell body (soma) that contains a nucleus with a prominent central nucleolus. The nucleus is surrounded by cytoplasm speckled with Nissl granules, which are clumps of rough endoplasmic reticula. Radiating from the cell bodies are cytoplasmic processes, called dendrites, which receive stimuli, and axons, which generate and conduct impulses. Multipolar neurons, such as motor neurons and interneurons, have numerous dendrites and a single axon. Bipolar neurons, such as in the retina and other special sense organs, have a single axon and a single dendrite that extend from the cell body. Unipolar neurons (pseudounipolar neurons) are sensory neurons that have one short process that arises from the cell body and bifurcates.

In the central nervous system, which includes the brain and spinal cord, there is a structural framework provided by **neuroglial cells**. The four types of neuroglia are astrocytes, oligodendrocytes, microglia, and ependymal cells; each has a unique structure and performs various functions. Additionally, the **meninges** (the pia mater, arachnoid mater, and dura mater) are formed of connective tissue and surround the brain and spinal cord, providing support and protection.

In the peripheral nervous system, which includes cranial nerves, spinal nerves, and ganglia, neurons are supported by connective tissue and special cells. For example, bundles of axons are fastened together by both loose and dense irregular connective tissue, forming peripheral nerves, and the axons of neurons are supported by Schwann cells (neurolemmocytes). Clusters of cell bodies of neurons of the peripheral nervous system, along with supportive satellite cells and connective tissue, form ganglia.

The myelin sheath of myelinated axons in the peripheral nervous system is formed by Schwann cells that encircle an axon along its length, separated by small gaps (nodes of Ranvier). The cell membrane of each Schwann cell layers around the axon many times, forming white, lipidrich myelin.

Myelin is formed by oligodendrocytes in the central nervous system. White matter in the CNS consists mainly of myelinated axons. Gray matter lacks myelinated structures, and is formed mostly of the cell bodies of neurons, dendrites, and neuroglia.

Selected examples of nervous system elements as they typically appear in histologic preparations of various kinds are presented in this chapter. Additionally, sections through portions of the brain, brain stem, and spinal cord have been included. The organs of special sense, the eye and ear, are treated in separate chapters.

| Word Roots | | |
|------------|----------------------|---|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Dendr | Tree | Dendrites branch like a tree. |
| Gangli | A knot or a swelling | Cell bodies of neurons in the peripheral nervous system form clusters called ganglia. |
| Glia | Glue | Neuroglia are cells that provide a supportive role. |
| Lemma | Sheath | Schwann cells, also called neurolemmocytes, envelop axons like a sheath. |
| Menin | A membrane | The meninges are membranes associated with the brain and spinal cord. |
| Neuro | Nerve | Nerve cells, also called neurons, are cells of nervous tissue. |






Figure 9.1 × 25








Figure 9.4 × 250




Figure 9.5 × 62.5

| | KEY | |
|--|---|--|
| 1. Blood vessel 2. Bone, skull 3. Dendrite 4. Dura mater 5. Granular layer 6. Mesothelium 7. Molecular layer 8. Outer granular layer 8. Outer granular layer | 9. Periosteum 10. Pia mater 11. Purkinje cell 12. Pyramidal cell 13. Pyramidal cell layer 14. Subarachnoid space 15. White matter | |

Figure 9.1. Cerebellum, Sheep. The molecular and granular layers of the gray matter and Purkinje cells are shown. The white matter lies deep to the gray matter.

Figure 9.2. Cerebellum, Sheep. Portions of the dendritic tree of the multipolar Purkinje cells are visible.

Figure 9.3. Cerebral Cortex, Dog. Outer portions of cerebral cortex with numerous blood vessels.

Figure 9.4. Cerebral Cortex, Pyramidal Cells, Dog. Magnified view of cells of the pyramidal layer.

Figure 9.5. Dura Mater, Goat. The dura remains attached to the skull when the latter is separated from the brain. It is a dense fibroelastic layer lined by a mesothelium. The dura merges with the periosteum of the skull.





Figure 9.6 × 260

| KEY | | |
|---|---|--|
| Arachnoid mater Cerebrospinal fluid Collagenous fiber | 4. Molecular layer 5. Pia mater 6. Small artery | |

Figure 9.6. Meninges, Cerebellum, Sheep (Mallory's). The arachnoid mater, subarachnoid space, and pia mater are shown. The subarachnoid space is filled with cerebrospinal fluid, which is stained purple. Wisps of collagenous fibers (blue) can be seen within the subarachnoid space. These connect the arachnoid layer with the pia.








Figure 9.8 × 12.5




Figure 9.9 × 125




Figure 9.10 × 62.5




Figure 9.11 × 125

| | KEY |
|---|---|
| Astrocyte, cell body Astrocyte, process Brain sand Capillary Cerebellum, vermis Choroid plexus Epithelium Fourth ventricle Granular layer | 10. Medulla 11. Molecular layer 12. Neuron 13. Perivascular space 14. Pia mater 15. Purkinje cell 16. Venule 17. Villus, x.s. |
| | |

Figure 9.7. Cerebellum, Dog. Portion of a sulcus containing a venule. The perivascular space surrounding the vessel is continuous with the subarachnoid space and separates the vessel from the pia mater on either side.

Figure 9.8. Choroid Plexus, Cat (Cresyl Violet). Portion of the fourth ventrical with choroid plexus in the roof of the medulla.

Figure 9.9. Choroid Plexus, Dog. The simple cuboidal epithelium and large, thin-walled capillaries are major constituents of the villi of the choroid plexus.

Figure 9.10. Brain Sand, Hypothalamus, Dog. Calcified, granular material called brain sand can be found dispersed through various parts of the brain, including the hypothalamus, cerebellum, and pineal gland.

Figure 9.11. Fibrous Astrocytes, Medulla, Cat (Golgi). These neuroglia cells have long processes that show little or no branching.




Figure 9.12




Figure 9.13 × 26

- Area shown in Figure 9.13
 Central canal
 Gray matter, ventral horn

- 4. Multipolar neuron, cell body 5. White matter

Figure 9.12. Spinal Cord, Cervical, x.s., Sheep. The area outlined by the rectangle is shown in Figure 9.13.

Figure 9.13. Spinal Cord, Cervical, x.s., Sheep (Masson's). The central canal, gray matter, and white matter are shown. The cell bodies of multipolar neurons reside in the gray matter.




Figure 9.14 × 125




Figure 9.15 × 125




Figure 9.16 × 180




Figure 9.17 × 62.5


| KEY | | |
|---|--|--|
| Axon hillock Central canal Cilia Dendrite Gray matter | 6. Nerve fibers 7. Neuron cell body 8. Nissl granules 9. Nucleolus 10. Nucleus | |

Figure 9.14. Central Canal, Spinal Cord, x.s., Sheep. Tall ependymal cells, some of which are ciliated, line the central canal.

Figure 9.15. Multipolar Neurons, Spinal Cord, Sheep. The axon hillock of two neurons can be seen. Nissl granules are absent from the hillock region, but extend into the dendrites.

Figure 9.16. Multipolar Neurons, Spinal Cord, Cow. Two multipolar neurons are shown in this smear preparation. Note prominent nucleoli and Nissl granules.

Figure 9.17. Dorsal Root Ganglion, Dog (Luxol Fast Blue/Cresylecht Violet). Portion of a dorsal root ganglion showing neurons and nerve fibers. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)



× 250 Figure 9.18




Figure 9.19 × 125




Figure 9.20 × 125




Figure 9.21 \times 62.5




Figure 9.22 \times 12.5

- 1. Adipose tissue
- 2. Epineurium
- 3. Muscularis externa, inner circular
- Muscularis externa, outer longitudinal
 Muscularis mucosae
- 6. Nerve
- 7. Nerve fiber

- 8. Neuron cell body
- 9. Nissl granules
- 10. Perineurium
- 11. Plexus
- 12. Satellite cell
- 13. Submucosa

Figure 9.18. Dorsal Root Ganglion, Dog (Luxol Fast Blue/ Cresylecht Violet). Flattened satellite cells envelop the round neuron cell bodies of the unipolar neurons. (Photograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 9.19. Parasympathetic Ganglion, Pulmonary Artery, Dog. A ganglion and associated nerve located in the adventitia of the artery.

Figure 9.20. Meissner's Plexus, Esophagus, x.s., Pig. These parasympathetic plexuses are located in the submucosa of the digestive tract. Note the characteristic large "owl's eye" nucleus of the neurons.

Figure 9.21. Auerbach's Plexus, Jejunum, I.s., Dog. These parasympathetic plexuses are located between the inner circular and outer longitudinal layers of the muscularis externa of the digestive tract.

Figure 9.22. Nerve, Myelinated, x.s., Pig (Masson's). The nerve shown is comprised of many fascicles bounded by a connective tissue sheath, the epineurium. Each fascicle is surrounded by a perineurium and contains numerous axons. A magnified view of a nerve fascicle in this preparation is shown in Figure 9.23.




Figure 9.23 × 260

Axon
 Endoneurium
 Myelin sheath

4. Perineurium5. Space artifact

Figure 9.23. Nerve Fascicle (portion), Myelinated, x.s., Pig (Masson's). Magnified view of one of the nerve fascicles shown in Figure 9.22. Delicate connective tissue fibers of the endoneurium are visible around individual myelinated axons.




Figure 9.24 \times 250




Figure 9.25 × 250




Figure 9.26 \times 180




Figure 9.27




Figure 9.28 \times 625

- 1. Axon
- 2. Blood vessel
- 3. Groove in plasma membrane of Schwann cell
- 4. Myelin sheath
- 5. Nerve fibers, l.s. 6. Nerve fibers, x.s.
- Perineurium
- 8. Schwann cell, nucleus

Figure 9.24. Nerve, Myelinated, Thoracic Wall, Cat. Myelin sheaths (pink) often present a scalloped or vacuolated appearance, an artifact of processing. Axons in this preparation are round, oval, or shriveled and either blue-gray or purple.


Figure 9.25. Axons, Myelinated, Medulla, Horse (Hagguist). Axons (blue) with myelin sheaths (pink) are seen in longitudinal sections and cross sections.


Figure 9.26. Nerve Fascicles, Unmyelinated, Ureter, Pig. Note the wavy appearance of the fibers that have been cut longitudinally. See Figures 9.27 and 9.28 for comments on the relationship of unmyelinated axons to Schwann cells.


Figure 9.27. Schwann Cell With Unmyelinated Axons. Unmyelinated axons of the peripheral nervous system are enveloped by Schwann cells. A Schwann cell may have 20 or more grooves in its surface, each groove containing one or more axons.

Figure 9.28. Nerve Fascicle (portion), Unmyelinated, Left Ventricle, Pig. Each Schwann cell enwraps several unmyelinated axons within grooves (which appear as vesicles in cross section) of its plasma membrane. Axons can be seen filling some of the grooves, while in other grooves the axons have shrunken from the plasma membrane of the Schwann cell. Unmyelinated axons are typically smaller in diameter than myelinated axons.

For other examples of labeled unmyelinated nerve fas**cicles**, see Figures 8.4, 10.13, 10.23, 10.38, 10.39, 13.95, 15.39, 15.40.










Figure 9.32 × 250




Figure 9.33 × 12.5

| Arteriovenous shunt Capsule Central canal Fluid-filled space Glycogen body Gray matter Inner core Mesenteric blood vessel | 9. Multipolar neurons 10. Nerve 11. Nuclear bag fiber 12. Nuclear chain fiber 13. Outer core 14. Pancreas 15. Skeletal muscle cell, x.s. 16. White matter |
|--|--|


Figure 9.29. Pacinian Corpuscle, Wholemount, Mesentery, Cat (Carmine). Two Pacinian corpuscles (one on the right somewhat distorted) are shown. Each is surrounded by a capsule of connective tissue within which are located concentric laminae of flattened cells that form the core. A nerve ending courses through the center of the corpuscle. The closely packed inner core cells surround the nerve. The peripheral laminae form a looser, outer core.


Figure 9.30. Pacinian Corpuscle, x.s., Pancreas, Cat. The Pacinian corpuscle is frequently seen in the pancreas of carnivores. See Figure 9.29 for description.


Figure 9.31. Small Encapsulated Nerve Endings, Dermis, Planum, Cow. Numerous encapsulated sensory nerve endings occur in the dermis of the planum near the epithelium.

Figure 9.32. Neuromuscular Spindle, x.s., Thoracic Muscle, Cat. A neuromuscular spindle is a proprioceptor located within a muscle. It consists of sensory and motor nerve endings and intrafusal fibers, which are narrow, modified skeletal muscle cells. Nuclear chain fibers are intrafusal fibers with a single row of nuclei, whereas nuclear bag fibers are intrafusal fibers that contain many closely packed nuclei. A capsule encloses the fluid-filled space that surrounds the intrafusal fibers.

Figure 9.33. Glycogen Body, Lumbosacral Enlargement, Spinal Cord, x.s., Chicken. The glycogen body is found only in birds. It consists of polyhedral, vesicular cells, each containing a central mass of glycogen and a peripherally displaced nucleus.










Figure 9.37 × 125




Figure 9.38 × 62.5

| | KEY |
|--|--|
| Axon Bone Capsule of Herbst corpuscle Cell of glycogen body Central canal Core Dermis Dorsal root ganglion Epidermis | 10. Muscle (moves feather) 11. Gray matter 12. Multipolar neuron 13. Nerve 14. Sharpey's fibers 15. Space artifact 16. Unipolar neuron 17. Ventral root 18. White matter |

Figure 9.34. Glycogen Body, Lumbosacral Enlargement, Spinal Cord, x.s., Chicken. Cells of the glycogen body in detail. See Figure 9.33 for description.

Figure 9.35. Dorsal Root Ganglion, Lumbosacral Enlargement, Spinal Cord, x.s., Chicken. Portions of the spinal cord, ventral root of a spinal nerve, dorsal root ganglion, and vertebra.

Figure 9.36. Dorsal Root Ganglion, Lumbosacral Enlargement, Spinal Cord, x.s., Chicken. Neuron cell bodies of unipolar neurons and myelinated axons are shown.

Figure 9.37. Herbst Corpuscle, Upper Beak, x.s., Chicken. These encapsulated nerve endings occur frequently in the skin of the bird. They are similar to Pacinian corpuscles of mammals and consist of an outer capsule of connective tissue, a laminated core, and an axial sensory nerve ending.

Figure 9.38. Herbst Corpuscle, Skin, Neck, Chicken. The Herbst corpuscles associated with follicles of feathered skin are sausage-shaped.

10

CARDIOVASCULAR SYSTEM

he heart pumps blood and conveys it to the tissues and organs through blood vessels. Fluid that escapes from the blood is returned to the venous system by lymphatic vessels.

BLOOD VESSELS

Vessels of the cardiovascular system are lined by an **endothelium**, which is typically a single layer of squamous cells. The smallest of the blood vessels, **capillaries**, are tiny endothelial tubes. They are easily overlooked in histologic sections, especially if they are compressed or collapsed.

The walls of arteries and veins are arranged into concentric layers: the inner tunica intima, middle tunica media, and outer tunica adventitia. The composition and thickness of these layers vary with the size and type of vessel. The tunica media is not always present.

Small arteries can be defined, arbitrarily, as possessing up to eight or nine layers of smooth muscle cells in the tunica media. The smallest of these vessels is usually termed an arteriole. Its wall is comprised of an endothelium (tunica intima), one or two layers of circularly arranged smooth muscle cells (tunica media), and a bit of surrounding loose connective tissue (tunica adventitia). Some of the larger small arteries have an internal elastic membrane (a sheet-like membrane forming the outer surface of the tunica intima). Small arteries are accompanied by small veins. The smallest veins are called venules. These are similar to arterioles, but have relatively thin walls and lack a tunica media of smooth muscle. An internal elastic membrane is not found in small veins.

As the diameter of a vessel increases, the tunics become larger and more elaborate. For example, the tunica intima of a medium artery contains connective tissue interspersed between the endothelium and internal elastic membrane. The thick tunica media, with

varying proportions of smooth muscle and elastic fibers, comprises the bulk of the wall. The connective tissue of the tunica adventitia contains collagenous and elastic fibers, small blood vessels (vasa vasorum), and nerves. A medium vein, in contrast, has less smooth muscle and fewer elastic fibers in the tunica media and possesses a thicker tunica adventitia.

Arteries ordinarily appear round in cross section and have an obvious, rippled, internal elastic membrane. Conversely, accompanying veins are larger in diameter with an irregular or collapsed lumen and thinner walls, and, except for some of the largest, they have no internal elastic membrane. The lumens of blood vessels in tissue sections often contain blood cells, plasma, or both. Although it can be difficult to distinguish between veins and lymphatic vessels, the latter have thinner walls than veins of similar size and normally do not contain erythrocytes. Valves may occur in both veins and lymphatic vessels.

There are several variations from the "typical" blood vessels: The tunica adventitia of large veins adjacent to the heart contains cardiac, rather than smooth, muscle. Some arteries have smooth muscle in the tunica intima, as well as the tunica media. Smooth muscle may be oriented either longitudinally or circularly. The tunica adventitia of arteries may be either abundant or scant.

The arteries of arteriovenous anastomoses lack an internal elastic membrane, but possess epithelioid (epithelial-like) longitudinally arranged smooth muscle cells. Special structures, the aortic and carotid bodies, are closely associated with the tunica adventitia of their respective arteries.

Many special vessels unique to certain organs such as the sinusoids of the liver, postcapillary venules of lymph nodes, and helicine arteries of the penis are presented elsewhere with their appropriate organ systems.

HEART

The heart is a muscular organ whose wall is composed of an endocardium, myocardium, and epicardium. The thickness and composition of the wall vary, being thickest in the ventricles and thinnest in the atria. The middle layer of cardiac muscle, the myocardium, predominates. Valves of connective tissue covered by an endothelium are extensions of the endocardium. Regions of the heart, including the base of the aorta and pulmonary trunk, as well as the atrioventricular orifices and septum, are supported by the cardiac skeleton. This cardiac skeleton may be in the form of dense irregular connective tissue, fibrocartilage, hyaline cartilage, or bone, and varies with age and among individuals.

A small amount of fluid occurs in the pericardial cavity between the epicardium (visceral pericardium) and the parietal pericardium.

| Word Roots | | |
|-------------------|---|---|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Adventicius | Coming from abroad, foreign; the outermost covering, derived from connective tissue, of a body part | The tunica adventitia of a blood vessel is the outermost layer that blends with surrounding structures. |
| Cardio Endo | The heart Within | The endocardium is the innermost layer of the wall of the heart. |
| Intima | Innermost | The innermost layer of the wall of a blood vessel is the tunica intima. |
| Tunica Media | A covering, cloak Middle | The tunica media forms the middle coat, or layer, of a typical blood vessel. |








Figure 10.2 × 625




Figure 10.3 × 125




Figure 10.4 × 250




Figure 10.5 × 250

1. Arteriole, x.s. 2. Capillary, I.s. 3. Capillary, x.s. 4. Endothelial cell, nucleus 5. Endothelial cell, surface cut 6. Erythrocytes 7. Macrophage 8. Mast cell 12. Small artery, x.s. 12. Small vein 13. Smooth muscle cell, nucleus 14. Uterine gland 15. Venule

Figure 10.1. Capillaries, x.s. and l.s., Diaphragm, Dog. Extensive capillary networks occur around muscle cells.


Figure 10.2. Capillary, I.s., Lamina Propria, Duodenum, Sheep. Erythrocytes are lined up in the lumen of this capillary.


Figure 10.3. Arterioles and Venules, Eyelid, Pig. Small blood vessels of various sizes are present in the dermis.


Figure 10.4. Arterioles, x.s., Endometrium, Uterus, Dog. The smallest of the arterioles shown have only one layer of smooth muscle in their walls.


Figure 10.5. Venule, l.s., Connective Tissue, Epiglottis, Goat. The wall of the venule consists of an endothelium surrounded by a small amount of connective tissue.

For other labeled examples of **capillaries**, see Figures 2.18, 8.8, 8.10, 8.11, 8.16, 8.17, 9.7, 12.31, and 14.26












Figure 10.10 × 250

KEY 1. Arteriole 2. Endothelial cell, nucleus 3. Erythrocytes 4. Internal elastic membrane 5. Plasma 6. Skeletal muscle 7. Small artery, x.s. 8. Small vein, collapsed, x.s. 9. Sweat gland 10. Tunica adventitia 11. Tunica media 12. Valve 13. Venule


Figure 10.6. Arterioles and Venule, I.s., Submucosa, Esophagus, Cat. The circular arrangement of the smooth muscle of the tunica media can be seen in the region where the arterioles have been cut tangentially.

Figure 10.7. Small Artery and Vein, x.s., Eyelid, Pig. These vessels are surrounded by portions of sweat glands in the dermis. Veins such as the one shown often have an irregular or collapsed lumen.


Figure 10.8. Small Artery, I.s., Esophagus, Pig.

Figure 10.9. Small Vein with Valve, I.s., Nose, Sheep. Valves are thin flaps of connective tissue covered on both sides by an endothelium.

Figure 10.10. Small Artery, x.s., with Branch, Subcutis, Dog.












Figure 10.13



Figure 10.14 \times 62.5




Figure 10.15 \times 62.5

- 1. Adipose tissue
- Artery
 Elastic fiber
- 4. Erythrocytes
- 5. External elastic membrane
- 6. Internal elastic membrane
- 7. Nerve fascicle, unmyelinated
- 8. Pancreas
- Tunica adventitia
- 10. Tunica intima, endothelium
- 11. Tunica media
- 12. Vein

Figure 10.11. Small Artery, Vein, and Nerve, x.s., Pancreas, Cat. Note that both of the vessels have a sparse adventitia.

Figure 10.12. Artery and Vein, x.s., Wattle, Rooster. Note the especially thick tunica media of the artery.

Figure 10.13. Medium Artery, x.s., Lymph Node, Pig. The rich pink color of the elastic fibers contrasts with the paler pink of the collagenous fibers and smooth muscle.

Figure 10.14. Medium Artery, x.s., Lymph Node, Pig (Orcein). Elastic fibers are stained reddish brown with orcein.

Figure 10.15. Medium Vein, I.s., Lymph Node, Pig. This vein accompanied the artery in Figures 10.13 and 10.14.

Summary of the Layers of the Wall of a Typical Medium Artery:

Tunica Intima (Tunica Interna)

Endothelium: A simple squamous epithelium Small amount of connective tissue

Internal elastic membrane

Tunica Media: Circularly arranged smooth muscle cells; some elastic and collagenous fibers

Tunica Adventitia (Tunica Externa): A layer of connective tissue that blends with surrounding structures

Helpful Hints


How to Distinguish Between a Typical Medium Artery and Medium Vein:

Artery


- Tends to be more round in cross section
- Has an internal elastic membrane
- Has a thicker tunica media


Vein


- Tends to be larger in diameter and have a collapsed, irregular lumen
- Does not have an internal elastic membrane
- Has a thicker tunica adventitia













× 25 **Figure 10.20**

| KEY | | |
|---|--|--|
| Adipose tissue External elastic membrane Internal elastic membrane Nerve fascicle, unmyelinated | 5. Tunica adventitia 6. Tunica intima, endothelium 7. Tunica media 8. Valve | |
| | | |


Figure 10.16. Medium Vein, I.s., Lymph Node, Pig (Orcein). Elastic fibers are stained reddish brown with orcein.


Figure 10.17. Medium Artery, x.s., Lymph Node, Cat. Longitudinally oriented smooth muscle is present in the tunica intima between the endothelium and internal elastic membrane.


Figure 10.18. Renal Artery, Near Aorta, x.s., Pig. Note both an inner and outer layer of smooth muscle in the tunica media. The inner layer is arranged longitudinally.

Figure 10.19. Portal Vein, x.s., Dog. Note the bundles of longitudinally arranged smooth muscle in the tunica adventitia, a characteristic of large veins.

Figure 10.20. Vein with Valves, x.s., Lip, Pig. Portions of two leaflets of a valve extend into the lumen of this vein.










Figure 10.24 × 250




Figure 10.25 × 12.5

| KEY | | |
|--|--|--|
| Anastomotic artery Aorta, lumen Epithelioid cells Nerve fascicle, unmyelinated Pulmonary artery, lumen | 6. Tunica adventitia 7. Tunica intima 8. Tunica intima, endothelium 9. Tunica media | |


Figure 10.21. Umbilical Artery (Right) and Vein (Left), x.s., Horse (Masson's). The tunica media of the umbilical artery is thicker than that of the umbilical vein.


Figure 10.22. Umbilical Artery, x.s., Horse (Masson's). The umbilical artery lacks an internal elastic membrane. The innermost smooth muscle cells of the tunica media are epithelioid (epithelial-like) and oriented longitudinally.


Figure 10.23. Glomus, Nose, Pig. The highly convoluted anastomotic artery, surrounding connective tissue, and nerves forming this organized arteriovenous anastomosis can be seen.


Figure 10.24. Arteriovenous Anastomosis, x.s., Lip, Pig. Longitudinally directed cells of smooth muscle of the tunica media are characteristically epithelioid (epithelial-like) in an anastomotic artery. These arteries lack an internal elastic membrane and have a small lumen.

Figure 10.25. Aorta (Left) and Pulmonary Artery (Right), x.s., Pig (Orcein). This preparation was stained with orcein to highlight elastic tissue (red-brown).












Figure 10.30 × 25

| KEY | | | |
|---|---|--|--|
| Adipose tissue Cardiac muscle Collagenous fibers Elastic fibers Endocardium | 6. Myocardium 7. Smooth muscle 8. Tunica intima, endothelium 9. Tunica media 10. Vasa vasorum | | |

Figure 10.26. Aorta, x.s., Dog (Orcein). This specimen was stained with orcein to emphasize elastic tissue (red-brown).


Figure 10.27. Pulmonary Artery, x.s., Sheep. Portion of the tunica intima and tunica media. Smooth muscle of the tunica media is oriented in various directions. Wavy, pink elastic fibers occur among the smooth muscle.


Figure 10.28. Vena Cava, x.s., Dog. This section was taken from a region near the heart. The tunica adventitia consists largely of cardiac muscle and adipose tissue.

Figure 10.29. Right Auricle, Pig (Orcein). The section was stained with orcein to show the distribution of elastic fibers (red-brown).

Figure 10.30. Right Auricle, Pig (Mallory's). This preparation shows the distribution of smooth muscle in the endocardium.










Figure 10.34 \times 12.5

- 1. Adipose tissue

- Adorda, lumen
 Atrium, myocardium
 Cardiac skeleton, cartilaginous
 Cardiac skeleton, fibrous
- 6. Mesenchyme-like tissue
- 7. Pulmonary artery, lumen 8. Tunica adventitia
- 9. Tunica intima, endothelium
- 10. Tunica media
- 11. Valve

Figure 10.31. Pulmonic (Semilunar) Valve, x.s., Dog. Pulmonic valves are located in the pulmonary artery near the heart. The section shows a portion of the fibrous cardiac skeleton.

Figure 10.32. Pulmonic (Semilunar) Valve, x.s., Dog (Masson's). Portions of two adjacent pulmonic valves are visible. The connective tissue of the valves and the tunica media of the pulmonary artery blend with the fibrous cardiac skeleton. A cushion of mesenchyme-like connective tissue lies adjacent to the cardiac skeleton.

Figure 10.33. Pulmonic (Semilunar) Valve, x.s., Dog. The valve consists of a core of dense irregular connective tissue sandwiched between two layers of endothelium.

Figure 10.34. Pulmonic (Semilunar) Valve, x.s., Dog. A portion of the atrial wall, pulmonary artery, aorta, and cardiac skeleton (cartilaginous and fibrous) are visible.




Figure 10.35 × 125




Figure 10.36 × 180




Figure 10.37 × 62.5




Figure 10.38 × 125




Figure 10.35. Cardiac Skeleton, Dog. The cartilaginous portion of the cardiac skeleton of the dog is formed from fibrocartilage containing numerous scattered chondrocytes.

Figure 10.36. Purkinje Cells, x.s. and l.s., Left Ventricle, Goat. Myofibrils are limited to the periphery of these large, modified cardiac muscle cells.

Figure 10.37. Visceral and Parietal Pericardium, Cat. The pericardium consists of a mesothelium (simple squamous epithelium) and underlying connective tissue. The mesothelium of the visceral pericardium (epicardium) covers the surface of the heart. The remainder of the pericardial cavity is lined by the mesothelium of the parietal pericardium.

Figure 10.38. Aortic Body, Pig. The aortic body is located between the pulmonary artery and aorta. It is a small, encapsulated structure containing blood vessels, nerves, and two types of parenchyma cells (see Figure 10.39).










Figure 10.41 \times 62.5




Figure 10.42 \times 62.5

- 1. Adipose tissue 9. Nerve fascicle, unmyelinated 2. Arteriole
 - 10. Plasma
 - 11. Type I cell
 - 12. Type II cell 13. Valve

 - 14. Vein 15. Venule

6. Erythrocytes
7. Lymph
8. Lymphatic vessel

3. Artery
4. Cecum, wall
5. Endothelium

Figure 10.39. Aortic Body, Pig. Two types of parenchyma cells can be distinguished in the aortic body. The type I (glomus) cell has a round nucleus and a granular cytoplasm. The type II (sustentacular) cell has few or no cytoplasmic granules and an oval nucleus. Type I cells usually occur in clusters surrounded by Type Il cells and connective tissue.

Figure 10.40. Lymphatic Vessel with Valve, Artery, and Vein, Cecal Tonsil, Chicken. Lymphatic vessels have a large lumen and a relatively thin wall. Valves may be present.

Figure 10.41. Lymphatic Vessel with Valve, Lymph Node, Pig. The valves of lymphatic vessels consist of a connective tissue core surrounded on each side by an endothelium.

Figure 10.42. Large Lymphatic Vessel, Submucosa, Cecum, Horse. The wall of the lymphatic vessel consists of an endothelium surrounded by a small amount of connective tissue. The latter blends with the connective tissue of the submucosa.

LYMPHATIC SYSTEM

ymphatic tissue consists predominantly of lymphocytes. These and a variable number of plasma cells, macrophages, and other cells occur among a framework of reticular cells and fibers. In H&E preparations lymphatic tissue appears purple because of the presence of numerous small lymphocytes, each with a basophilic nucleus and little cytoplasm.

MAMMALS

Types of Lymphatic Tissue

Diffuse lymphatic tissue is characterized by a moderate concentration of scattered lymphocytes. A round, oval, or irregularly circumscribed aggregation of mostly small, densely packed lymphocytes is called a lymphatic nodule. A nodule may contain a central pale area, the germinal center. Because the majority of cells of the germinal center are larger lymphocytes with more cytoplasm and lightly staining nuclei, this region appears pale in contrast to the dense corona (marginal zone, peripheral zone) of small lymphocytes. Diffuse lymphatic tissue and lymphatic nodules are components of most lymphatic organs. They also appear in the connective tissue of the digestive, respiratory, urinary, and reproductive organs, among other locations.

Aggregations of lymphatic nodules form Peyer's patches in the lamina propria and submucosa of the small intestine, particularly the ileum.

Tonsils

Tonsils are collections of lymphatic nodules and diffuse lymphatic tissue. They occur in the connective tissue below the epithelium in specific regions of the tongue, pharynx, and larynx.

Follicular tonsils are characterized by deep invaginations of the surface epithelium called crypts. A crypt together with its associated lymphatic tissue is a tonsillar follicle. Collectively, several follicles form the tonsil. Examples of tonsils with crypts include the following: lingual tonsils of the horse, pig, and cow; tubal tonsils of the pig; paraepiglottic tonsils of the pig, sheep, and goat; palatine tonsils of the horse, pig, and ruminant. In the palatine tonsils of ruminants the crypts lead into a common sinus, which then opens onto the surface.

Tonsils without crypts have a smooth, somewhat folded, or bulging surface, but lack deep invaginations of the epithelium. Examples of these are the tubal tonsils of ruminants, the paraepiglottic tonsil of the cat, and the palatine tonsils of carnivores.

Salivary glands associated with tonsils are typically mucous glands except those in carnivores, where they are mixed (mucous and serous combined).

Lymph Nodes and Hemal Nodes

A lymph node is organized into a cortex and medulla. The cortex consists of lymphatic nodules surrounded by diffuse lymphatic tissue. Extensions of the latter tissue into the medulla are called medullary cords. Lymphocytes, other leukocytes, macrophages, and plasma cells can be found in the medullary cords.

A capsule of connective tissue, with some smooth muscle and elastic fibers, covers the lymph node. Parts of the capsule extend inward as **trabeculae**. Afferent lymphatic vessels penetrate the capsule to join the subcapsular sinus. Cortical sinuses connect the **subcapsular sinus** to **medullary sinuses**. The latter lead to efferent lymphatic vessels at the hilus. The various sinuses are less cellular than the parenchyma and appear pale by comparison. They are lined by a discontinuous endothelium and are spanned by a webwork of cytoplasmic processes of reticular cells. They contain some free cells such as lymphocytes and macrophages.

Blood vessels enter and leave the node mostly from the region of the hilus. Unique blood vessels called **postcapillary venules** are found in the deep cortex. They are lined by elongated cells that appear cuboidal when cut in cross section. Lymphocytes migrate between these cells.

The amount or arrangement of cortical and medullary tissue can vary from that of the "typical" lymph node. The lymph node of the pig, for example, is characteristically atypical with the location of the cortical and medullary tissue, as well as the flow of lymph, being reversed.

Hemal nodes occur along blood vessels of ruminants. They are characterized by blood-filled sinuses between cellular cords. Connective tissue and some smooth muscle form the capsule and trabeculae (which are sparse). Hemal nodes lack lymphatic vessels. Hemolymph nodes, in contrast to hemal nodes, possess lymphatic vessels. Their sinuses receive a mixture of blood and lymph.

Spleen

The spleen has a capsule that is rich in smooth muscle and elastic fibers. In horses and cows two or three layers of

muscle are oriented perpendicular to each other, while in carnivores, pigs, sheep, and goats, the muscle fibers are interwoven. The capsule is thickest in the horse and cow and thinnest in carnivores. **Trabeculae** project into the interior of the spleen from the capsule. They tend to be especially large in cows and sheep.

The parenchyma of the spleen is divisible into the white and red pulp. Dense accumulations of lymphocytes, arranged around central arteries, form the periarterial lymphatic sheaths (PALS). These, along with lymphatic nodules, comprise the white pulp. White pulp appears purple in H&E preparations because of the high concentration of numerous small lymphocytes. Red pulp, because of the large numbers of erythrocytes it contains in its reticular meshwork and blood vessels, is stained red in H&E preparations.

The splenic artery enters the hilus of the spleen and branches into trabecular arteries. When these enter the parenchyma of the spleen and become surrounded by white pulp, they are called central arteries (not necessarily located in the center of the PALS). On leaving the white pulp, the central artery branches into a group of pulp arteries. These, in turn, branch into two or three arterioles, which terminate in two or more capillaries. Commonly, the pulp arteries and their branches are called a penicillus because, collectively, they resemble the bristles of an artist's brush. A portion of the capillaries of the penicillus becomes surrounded by concentric layers of macrophages contained in a reticular framework. These cellular and fibrous thickenings are called ellipsoids (pericapillary macrophage sheaths). The term sheathed capillary is used by some authors for the combined unit consisting of the capillary and the ellipsoid. Ellipsoids are especially abundant in the marginal zone, the region between the red and white pulp. They are very large and numerous in pigs. The capillaries of the ellipsoids continue as terminal arterial capillaries. Arterial capillaries may join venous sinuses or pulp veins (closed circulation), or they may empty directly into the spaces of the reticular meshwork of the red pulp (open circulation).

The spleen of the dog is a **sinusal** spleen. The red pulp contains typical **venous** (splenic, vascular) **sinuses**. These are wide channels lined by elongated, longitudinally oriented endothelial cells. The spleens of the cat, horse, pig, and ruminant are classified as **nonsinusal**, having poorly developed or no sinuses. Wisps of smooth muscle in the red pulp are most numerous in pigs and ruminants.

Thymus

The **thymus** gland is covered by a thin capsule of connective tissue that projects inward as **septa**, partially dividing the organ into **lobules**. The parenchyma of each lobule is organized into a **cortex** of mostly small, densely packed lymphocytes and a **medulla** with fewer and larger lymphocytes. The medulla is continuous between lobules. The thymus lacks lymphatic nodules and is supported by a unique cytoreticulum of stellate, epithelial reticular cells and only a few reticular fibers.

Hassall's (thymic) corpuscles occur in the medulla of each lobule. They are concentric whorls of acidophilic and

flattened reticular cells that may become swollen, keratinized, and calcified centrally. They are found exclusively in the thymus gland.

As an animal ages, much of the thymus becomes replaced by adipose tissue.

CHICKEN

Lymph nodes do not occur in the chicken. However, diffuse lymphatic tissue and lymphatic nodules are widespread.

Spleen

The spleen of the chicken is covered by a muscular capsule, but trabeculae are absent. Areas of red and white pulp are less distinct than in the mammalian spleen. White pulp is diffusely scattered throughout the spleen and is composed primarily of small lymphocytes. It contains sheathed arteries and, occasionally, lymphatic nodules. Red pulp is formed from venous sinuses and anastomosing cords of reticular cells, macrophages, lymphocytes, and red blood cells.

Thymus

As in mammals, the **thymus** is arranged into incompletely separated lobules of cortical and medullary tissue. Typical Hassall's corpuscles, similar to those found in mammals, are seen infrequently. Instead, diffuse forms of Hassall's corpuscles, called **reticular structures**, are abundant in the medulla. These are pale, irregular masses of reticular cells with vesicles that contain acidophilic material and degenerating cells. **Myoid cells**, characterized by a fibrous cytoplasm, also occur in the medulla.

Bursa of Fabricius

The bursa of Fabricius is a saclike dorsal diverticulum of the proctodeum that is unique to birds. It is characterized by tall, thick mucosal folds (plicae) filled with numerous polyhedral follicles. Each follicle, composed of lymphatic tissue, is divided into a cortex and medulla. A layer of undifferentiated epithelial cells occupies the periphery of the medulla, which is separated from the cortex by a capillary layer. The bursa is lined by a pseudostratified columnar epithelium, except at the apex of each follicle, which is covered by an epithelial tuft of simple columnar cells.




Figure 11.1 ×25




Figure 11.2 ×12.5




Figure 11.3 ×25




Figure 11.4 ×250




Figure 11.5

| KEY | | |
|-----------------------------|-----------------------|--|
| 1. Corona | 7. Mucosa | |
| 2. Diffuse lymphatic tissue | 8. Muscularis externa | |
| 3. Germinal center | 9. Reticular cell | |
| 4. Lymphatic nodule | 10. Submucosa | |
| 5. Lymphocyte, large | 11. Villus | |
| 6. Lymphocyte, small | | |

Figure 11.1. Lymphatic Nodules and Diffuse Lymphatic Tissue, Pyloric Stomach, Cat. Dense aggregations of lymphocytes form lymphatic nodules in the lamina propria.

Figure 11.2. Lymphatic Nodules and Diffuse Lymphatic Tissue, Colon, x.s., Pig. The mucosa and submucosa contain diffuse lymphatic tissue and large lymphatic nodules with germinal centers. Lymphatic nodules are especially numerous in the digestive tract of the pig.

Figure 11.3. Lymphatic Nodule, Colon, Pig. Cells of the germinal center and corona. Many small lymphocytes occur in the peripheral corona; fewer and larger cells are seen in the germinal center.

Figure 11.4. Lymphatic Nodule, Colon, Pig. Detail of cells of the germinal center and corona. Small lymphocytes are characterized by a heterochromatic nucleus and scant cytoplasm.

Figure 11.5. Peyer's Patch, Ileum, x.s., Cat. A Peyer's patch is an aggregation of lymphatic nodules in the lamina propria and submucosa of the small intestine.




Figure 11.6 ×12.5




Figure 11.7 ×12.5




Figure 11.8




Figure 11.9 ×12.5




Figure 11.10 ×125

| | KEY |
|-----------------------------|---|
| 1. Diffuse lymphatic tissue | 9. Serous demilune |
| 2. Epiglottis | 10. Skeletal muscle |
| 3. Fossa semilunar fold | Stratified squamous epithelium, |
| 4. Lymphatic nodule | semilunar fold |
| 5. Mucous acinus | 12. Stratified squamous epithelium, tonsil |
| 6. Muscularis externa | 13. Submucosa |
| 7. Salivary glands | 14. Vestibular fold |
| 8. Serous acinus | 15. Villus |

Figure 11.6. Peyer's Patch, Ileum, x.s., Dog. Eight lymphatic nodules of a portion of a Peyer's patch are visible in this section.

Figure 11.7. Paraepiglottic Tonsil, Larynx, I.s., Cat. In the cat an accumulation of lymphatic tissue in the lateral wall of the larynx, between the epiglottis and the vestibular fold, forms a tonsil without crypts.

Figure 11.8. Palatine Tonsil, Dog. In the dog the entire tonsil lies within a fossa (a small hollow) and is covered in part by a semilunar fold. The palatine tonsils of carnivores lack crypts.

Figure 11.9. Palatine Tonsil, Dog. A portion of a tonsil and semilunar fold.

Figure 11.10. Palatine Tonsil, Dog. Mixed salivary glands are associated with the wall of the tonsils in carnivores. In other species only mucous glands are present.







Figure 11.12 ×12.5



1 -9 -8 4 -5 -7 12

Figure 11.14




Figure 11.15 ×12.5

| rigure 11.15 | ×12.5 |
|---|---|
| | KEY |
| Capsule Cortical sinus Crypt Deep cortex Diffuse lymphatic tissue Leukocyte infiltration Lymphatic nodule | 8. Medullary cord 9. Medullary sinus 10. Salivary glands, mucous 11. Stratified squamous epithelium 12. Subcapsular sinus 13. Trabecula |
| | |

Figure 11.11. Palatine Tonsil, Horse. The palatine tonsils of noncarnivores have crypts (surface invaginations lined by stratified squamous epithelium).

Figure 11.12. Palatine Tonsil, Horse. Note that the continuity of each crypt with the surface is not always evident.

Figure 11.13. Palatine Tonsil, Horse. Lymphocytes and other leukocytes have infiltrated and partly obliterated the epithelial lining of this crypt.

Figure 11.14. Lymph Node, Cow. The lymph node is surrounded by a capsule. Trabeculae project inward from the capsule. The cortex contains sinuses, diffuse lymphatic tissue, and lymphatic nodules. The medulla is composed of medullary cords and sinuses.

Figure 11.15. Lymph Node, Dog. This histologic section of a lymph node at low magnification shows the same features as the drawing in Figure 11.14.




Figure 11.16 ×12.5




Figure 11.17 ×12.5




Figure 11.18 ×12.5




Figure 11.19 ×12.5




Figure 11.20 ×12.5

| KEY | | |
|------------------------------|-----------------------------|--|
| 1. Capsule | 6. Lymphatic nodule | |
| 2. Cartilage plate, bronchus | 7. Meďulla, lymph node | |
| 3. Cortex, lymph node | 8. Medullary cord | |
| 4. Cortical tissue | 9. Medullary sinus | |
| 5. Diffuse lymphatic tissue | 10. Primary bronchus, lumen | |

Figure 11.16. Tracheobronchial Lymph Node, Dog. This lymph node is adjacent to the wall of a primary bronchus near the tracheal bifurcation.

Figure 11.17. Lymph Node, Horse. The arrangement of cortical and medullary tissues may be atypical in some of the lymph nodes of mammals. In the example shown the distribution of cortical and medullary components is the reverse of that commonly expected.

Figure 11.18. Lymph Node, Horse. Not only is the arrangement of cortical and medullary components reversed in this section, but the proportion of the medullary tissue is much greater than usual.

Figure 11.19. Lymph Node, Pig. The lymph nodes of pigs consistently show an atypical pattern. In this section cortical tissue is predominantly central, while medullary tissue occurs both superficially and internally.

Figure 11.20. Lymph Node, Cow. The lymph node of the cow is often characterized by the presence of large lymphatic nodules.




Figure 11.21 ×62.5




Figure 11.22 ×125




Figure 11.23 ×125



Figure 11.24 ×125




Figure 11.25 ×125

| | KEY |
|---|---|
| Capsule Cortical sinus Diffuse lymphatic tissue Lymphatic nodule | 5. Reticular fiber6. Smooth muscle7. Subcapsular sinus8. Trabecula |
| | |

Figure 11.21. Cortex, Lymph Node, Horse. The subcapsular sinus below the capsule continues as cortical sinuses that parallel the trabeculae through the cortex.

Figure 11.22. Cortex, Lymph Node, Cow (Masson's). The inner portion of the capsule contains smooth muscle (pink).

Figure 11.23. Cortex, Lymph Node, Cow. The trabecula contains smooth muscle.

Figure 11.24. Cortex, Lymph Node, Cow (Silver). A network of fine, branching reticular fibers provides a supportive framework for the diffuse and nodular lymphatic tissue.

Figure 11.25. Cortex, Lymph Node, Sheep (Mallory's). Continuity of the subcapsular sinus with the cortical sinus is evident.




Figure 11.26 ×250




Figure 11.27 ×250




Figure 11.28 ×12.




Figure 11.29 ×250




Figure 11.30 ×125

| KEY | | |
|------------------------------|--------------------------------|--|
| 1. Diffuse lymphatic tissue | 8. Multinucleate giant cell | |
| 2. Endothelial cell, nucleus | 9. Postcapillary venule, l.s. | |
| 3. Lymphocyte | 10. Postcapillary venule, x.s. | |
| 4. Macrophage | 11. Reticular cell | |
| 5. Mast cell | 12. Smooth muscle | |
| 6. Medullary cord | 13. Trabecula | |
| 7. Medullary sinus | | |


Figure 11.26. Cortex, Lymph Node, Sheep (Mallory's). Smooth muscle cells (pink) among collagenous fibers (blue) of the capsule.

Figure 11.27. Deep Cortex, Lymph Node, Dog. Postcapillary venules, l.s. and x.s. These vessels are lined by elongated cells that appear cuboidal in cross section. Lymphocytes migrate between the endothelial cells of the postcapillary venules.

Figure 11.28. Cortex, Lymph Node, Horse. Multinucleate giant cells, derived from the coalescence of macrophages, are sometimes found in lymph nodes.

Figure 11.29. Medulla, Lymph Node, Cow. In this preparation reticular cells, endothelial cells, and macrophages contain numerous pigment granules.

Figure 11.30. Medulla, Lymph Node, Sheep. Smooth muscle is distributed throughout the medullary sinuses.








Figure 11.32 ×360 2 1




Figure 11.33 ×25




Figure 11.34 ×250

| KEY | | |
|---|--|--|
| 1. Adipose tissue | 8. Lymphocyte | |
| 2. Capsule | Macrophage with erythrocytes | |
| Diffuse lymphatic tissue Endothelial cell, nucleus | 10. Medullary cord | |
| 4. Endothelial cell, nucleus | 11. Medullarý sinus | |
| 5. Erythrocytes | 12. Megakaryocyte | |
| 6. Granulocyte | 13. Reticular cell | |
| 7. Lymphatić nodule | 14. Subcapsular sinus | |

Figure 11.31. Medulla, Lymph Node, Dog. Cellular medullary cords surround medullary sinuses that are lined incompletely by endothelial cells. A megakaryocyte is present in a medullary cord.

Figure 11.32. Medulla, Lymph Node, Dog. Macrophages containing phagocytized erythrocytes are evident in the medullary sinuses.

Figure 11.33. Hemal Node, Sheep. The general organization is much like that of a lymph node, but the sinuses are filled with blood. Lymphatic nodules are scarce, and trabeculae of connective tissue are not apparent.

Figure 11.34. Hemal Node, Sheep. The subcapsular (marginal) sinus is filled with blood. Reticular cells of the sinus contain phagocytized material.




Figure 11.35




Figure 11.36 ×12.5




Figure 11.37 ×62.5




Figure 11.38 ×250




Figure 11.39

| KEY | | |
|---------------------|------------------|--|
| 1. Capsule | 7. Mesothelium | |
| 2. Central artery | 8. Red pulp | |
| 3. Ellipsoid | 9. Trabecula | |
| 4. Endothelial cell | 10. Venous sinus | |
| 5. Lymphatic nodule | 11. White pulp | |
| 6. Marginal zone | · · | |

Figure 11.35. Spleen, Dog. This drawing is of a small portion of the spleen.

Figure 11.36. Spleen, Dog. The parenchyma of the spleen is organized into red pulp and white pulp (periarterial lymphatic sheaths and lymphatic nodules). Trabeculae extend inward from the capsule and are seen throughout the red pulp.

Figure 11.37. Spleen, Dog. Note the smooth muscle in the capsule and trabeculae. The spleen of the dog is a sinusal spleen, containing venous sinuses. See Figure 11.38.

Figure 11.38. Spleen, Dog. Venous sinuses are lined by longitudinally oriented, elongated endothelial cells. The nuclei may or may not be apparent in cross sections of such lining cells. Erythrocytes fill the sinuses and the spaces of the red pulp.

Figure 11.39. Spleen, Dog. Ellipsoids can be seen in the marginal zone between the periarterial lymphatic sheath (white pulp) and the red pulp. They are also present in the red pulp.




Figure 11.40 ×125




Figure 11.41 ×25




Figure 11.42 ×62.5





Figure 11.43 ×62.5



igure 11.44 ×62.5

| rigure 11.44 | ×02. |
|---|--|
| | KEY |
| Capillary lumen Capsule Central artery Elastic fiber Ellipsoid Marginal zone | 7. Periarterial lymphatic sheath 8. Red pulp 9. Serosa 10. Smooth muscle 11. Trabecula |

Figure 11.40. Spleen, Pig (Orcein). The capsule and trabeculae are rich in elastic fibers (red-brown).

Figure 11.41. Spleen, Pig (Mallory's). Ellipsoids are abundant in the pig. They are especially numerous in the vicinity of the marginal zone of a periarterial lymphatic sheath. See Figure 11.44 for details of ellipsoids.

Figure 11.42. Capsule, Spleen, Horse. The capsule of the spleen of the horse and cow contains layers of smooth muscle oriented at right angles to each other, rather than being interwoven as in carnivores, pigs, sheep, and goats. In this preparation there are three distinct layers of muscle. Compare with Figures 11.37, 11.43, and 11.45.

Figure 11.43. Capsule, Spleen, Cow. The capsule contains two thick layers of smooth muscle oriented at right angles to each other.

Figure 11.44. Spleen, Pig. Ellipsoids are especially abundant in the spleen of the pig. Each consists of macrophages and reticular fibers that surround a capillary.




Figure 11.45 ×62.5




Figure 11.46 ×25




Figure 11.47 ×12




Figure 11.48 ×12.5


| KEY | | |
|-------|----------------|------------------|
| 1. C | apsule | 7. Septum |
| 2. C | ortex | 8. Serosa |
| 3. El | astic fiber | 9. Smooth muscle |
| 4. Ly | mphatic vessel | 10. Trabecula |
| 5. M | Nedulla | 11. White pulp |
| 6. Re | ed pulp | |

Figure 11.45. Capsule, Spleen, Sheep. In sheep the bulk of the capsule contains many interwoven smooth muscle cells. The smooth muscle in the capsule of the spleen of carnivores (Figure 11.37), pigs, and goats has a similar arrangement. In the horse (Figure 11.42) and cow (Figure 11.43) the muscle cells are arranged in layers instead. Elastic fibers can be observed as faint pink spirals.

Figure 11.46. Spleen, Sheep. Note the thick trabecula. Characteristically, the spleens of cows and sheep have thick trabeculae. Compare with Figure 11.41.

Figure 11.47. Red Pulp, Spleen, Sheep. Wisps of smooth muscle are scattered throughout the red pulp.

Figure 11.48. Thymus, Puppy. A thin capsule of connective tissue covers the thymus. Lobules, incompletely divided by connective-tissue septa, consist of an outer, dark cortex and an inner, pale medulla. The medulla is continuous between adjacent lobules.



| | THE REAL PROPERTY AND ADDRESS OF THE PARTY AND | THE RESERVE AND ADDRESS OF THE PARTY OF THE |
|--------------|--|---|
| Figure 11.49 | | ×260 |

| | KEY | |
|---|------------|--|
| Cortex Hassall's corpuscle | 3. Medulla | |

Figure 11.49. Thymus, Puppy. Portion of the medulla and cortex. The cortex consists predominantly of small lymphocytes. The lymphocytes of the medulla are larger and less abundant. The medulla contains concentrically arranged, swollen, and keratinized reticular cells that form Hassall's corpuscles, which are characteristic of the thymus.




Figure 11.50 ×12.5




Figure 11.51 ×12.5




Figure 11.52 ×25




Figure 11.53 ×250




Figure 11.54 ×360

- 1. Adipose tissue
- 2. Blood vessel
- 3. Connective tissue
- Crypt of Lieberkühn (intestinal gland)
- 5. Erythrocyte
- 6. Granulocyte
- 7. Lymphatic nodule
- 8. Mesothelium

- 9. Muscularis externa
- 10. Red pulp
- 11. Reticular cell
- 12. Sheathed artery, lumen
- 13. Smooth muscle of capsule
- 14. Thymic tissue
- 15. Viĺlus
- 16. White pulp

Figure 11.50. Thymus, Cat (Old). In older animals, functional thymic tissue is largely supplanted by adipose tissue.

Figure 11.51. Cecal Tonsil, x.s., Chicken. The accumulation of diffuse and nodular lymphatic tissue in the lamina propria and submucosa near the opening of each cecum is called the cecal tonsil.

Figure 11.52. Spleen, Chicken. Red pulp (pink) intermingles with white pulp (purple). The white pulp contains a few lymphatic nodules. Trabeculae of connective tissue are absent.

Figure 11.53. Capsule, Spleen, Chicken. Layers of smooth muscle make up a substantial part of the capsule.

Figure 11.54. Spleen, Chicken. Sheathed arteries, x.s., in white pulp. These vessels are lined by plump endothelial cells surrounded by reticular cells.




Figure 11.55 ×12.5




Figure 11.56 ×250



Figure 11.57 ×12.5




Figure 11.58 ×125




Figure 11.59 ×250

| KEY | | |
|------------------------------|---|--|
| 1. Bursa of Fabricius, lumen | 8. Muscularis | |
| 2. Capillary layer | 9. Myoid cell | |
| 3. Cortex | Pseudostratified epithelium | |
| 4. Epithelial tuft | 11. Reticular structure | |
| 5. Follicle | 12. Septum | |
| 6. Lamina propria | 13. Undifferentiated epithelial cell | |
| 7. Medulla | 14. Vesicle | |

Figure 11.55. Thymus, Chicken. The thymus of the chicken is similar to that of mammals. The pale areas throughout the medulary regions of the lobules are called reticular structures (Figure 11.56)

Figure 11.56. Medulla, Thymus, Chicken. Myoid cells, cut obliquely, are characterized by a fibrous cytoplasm and peripheral nuclei. The pale-staining reticular structure in this section is considered a diffuse form of a Hassall's corpuscle. It consists of diffuse groups of reticular cells and scattered vesicles. The vesicles may contain eosinophilic material or degenerating cells.

Figure 11.57. Bursa of Fabricius, Chicken. Portions of the long mucosal folds (plicae) project into the lumen of the bursa. Numerous follicles, each composed of a cortex and medulla, fill the lamina propria of each fold.

Figure 11.58. Bursa of Fabricius, Chicken. Where the apex of a follicle contacts the epithelium, tall, pale columnar cells with apical nuclei form an epithelial tuft. Elsewhere, mucosal folds are covered by a pseudostratified columnar epithelium.

Figure 11.59. Bursa of Fabricius, Chicken. A portion of a follicle. The darkly stained cortex is composed mostly of many small lymphocytes. The paler medulla contains fewer cells of various sizes. A layer of undifferentiated epithelial cells, which are cuboidal with an acidophilic cytoplasm, occupies the periphery of the medulla. A capillary network separates the cortex and the medulla.

INTEGUMENT

The integument includes the skin and its derivatives. Skin consists of two layers, an epidermis and dermis, joined to underlying structures such as muscle and bone by the subcutis (subcutaneous tissue). Sweat, sebaceous, and mammary glands, as well as hair and feather follicles, are epidermal structures that are located in the dermis and subcutis. The highly keratinized claws and hooves of mammals, and the beak, claws, and scales of fowl are also skin derivatives.

MAMMALS

Skin

Epidermis

The epidermis of thick skin is a keratinized, stratified, squamous epithelium. The stratum basale is a single layer of cuboidal to columnar cells that lies on a basement membrane adjacent to the dermis. These cells give rise to the stratum spinosum, a layer of variable thickness whose polygonal cells become squamous toward the surface. Cells of the stratum granulosum contain basophilic keratohyalin granules in their cytoplasm. The stratum lucidum is a thin, pale, eosinophilic, translucent layer. It is limited to regions where the epidermis is very thick, such as the digital pads of carnivores. In structures composed of hard keratin (rather than soft keratin), such as hooves and claws, both the stratum granulosum and stratum lucidum are absent. The most superficial layer of skin, the stratum corneum, is composed of dead, keratinized squamous cells that slough from the surface. Cell division within the stratum basale and stratum spinosum allows continued growth of the epidermis.

The epidermis of thin skin is composed of relatively few cells, but the number varies with the location. Thin skin lacks a stratum lucidum, and a stratum granulosum is not always evident.

Dermis

The dermis consists of loose and dense irregular connective tissue containing blood vessels, lymphatic vessels, and nerves. The superficial, loose connective tissue of the dermis, the papillary layer (superficial layer), forms projections called dermal papillae that interdigitate with the epidermis and serve to anchor the two layers. Dermal papillae are especially prominent in thick skin. The deep layer of dense irregular connective tissue is called the reticular layer (deep layer) of the dermis.

Epidermal Derivatives

Hairs and Hair Follicles

Hairs are associated with regions of the body covered by thin skin. They arise from germinal (matrix) cells of the hair bulb at the base of the hair follicle. Multiplication of germinal cells results in growth of the hair. Near its origin, a hair consists of a central medulla of cuboidal cells, a cortex of flattened cells oriented parallel to the long axis of the hair, and an outer cuticle consisting of scalelike cells that partially overlap so their free edges are directed upward toward the surface of the skin. The cells of the hair become keratinized as they are pushed toward the surface from the region of the hair bulb. Within the hair the medulla may become reduced distally, and it is absent entirely in wool hairs.

Hair follicles are set obliquely in the dermis or subcutis, although in sheep they tend to be vertical. A vascular dermal papilla projects into the hair bulb. Melanocytes, located close to the dermal papilla among matrix cells, have cytoplasmic processes that provide pigment to the hair cells. The germinal cells of the matrix, in addition to forming new hair cells, give rise to the inner root sheath of the follicle. The cuticle of the inner root sheath is composed of overlapping, scalelike cells similar to those of the cuticle of the hair, but their free edges are directed downward so that the hair and inner-root sheath interlock. The inner root sheath becomes keratinized and tapers distally, ending close to the opening of sebaceous glands into the follicle. The peripheral external root sheath represents a downward continuation of the epidermis. A connective tissue (dermal) sheath surrounds the follicle and abuts the basement (glassy) membrane of the external root sheath. It blends with the rest of the dermal connective tissue. An arrector pili muscle (smooth muscle) inserts on the connective tissue sheath of the follicle and originates from the superficial layer of the dermis.

Single (simple) hair follicles are evenly distributed in the skin of horses and ruminants and occur in groups of three in pigs. Most of the follicles in carnivores are compound follicles. Each compound follicle is formed from a single primary follicle and several secondary follicles. The follicles unite at the level of the openings of the sebaceous glands, forming a common follicle, which extends from the point of union to the skin surface. The hairs that are produced exit as a group to the surface through the common follicular opening. Sinus (tactile) hairs are limited to the face region. They are produced by large follicles that are well innervated and that contain blood-filled sinuses within

their connective-tissue sheaths. In horses, pigs, and ruminants, the sinus is trabeculated throughout its length. In carnivores, the upper region is nontrabeculated, forming an annular sinus.

Glands and Other Epidermal Deriviatives

The short ducts of sebaceous glands usually empty into hair follicles, although they may also empty directly onto the skin surface. Basal (stem) cells of sebaceous glands divide and give rise to vacuolated secretory cells that synthesize lipid. The innermost, mature secretory cells die and break apart, forming an oily product called sebum. This form of product release is called holocrine secretion.

Sweat glands may be winding (serpentine) or highly coiled, and may be either tubular or saclike. They empty their secretion through a duct, either into a hair follicle or onto the skin surface. The epithelium of the secretory portion of the gland varies from flattened to columnar. Contractile myoepithelial cells surround the secretory cells and the initial portion of their ducts.

Traditionally, sweat glands have been classified as either merocrine (secretory product released by exocytosis) or apocrine (secretory product released when small pieces of cytoplasm containing the product are pinched off the free surface of the cell). Recent evidence, however, has suggested that this may not be true and that all sweat glands use the merocrine (eccrine) form of release.

Special regions in the skin of various species have numerous, well-developed glands. The carpal glands of pigs consist of masses of merocrine sweat glands. Numerous apocrine sweat glands characterize the mental organ of pigs and the interdigital and inguinal pouches of sheep. The submental organ of cats, the supracaudal gland of carnivores, the infraorbital pouch of sheep, and the scent (horn) glands of goats contain many large sebaceous glands.

The skin of the nose of horses is thin with fine hairs, sebaceous and sweat glands, and occasional sinus hairs. The planum of the nose of the other domestic mammals is covered by a thick, highly keratinized epidermis. The planum nasale of carnivores is devoid of glands and hairs. In cats, the epidermis forms numerous small bumps, while that of the dog is rather flat with surface grooves. The planum rostrale of the pig contains numerous merocrine sweat glands and sparse hairs. The planum nasolabiale of the cow and the planum nasale of sheep and goats are hairless and contain compound acinar glands that produce a serous secretion.

Digital pads of cats and dogs are covered by a very thick epidermis that is smooth in the dog and roughened by conical papillae in the cat. Coiled merocrine sweat glands occur in the dermis and the digital cushion of the pads.

Lobules of mammary glands are situated in the subcutis and consist of tubuloacinar glands and intralobular ducts. When a mammary gland is active, secretory tissue is prominent, and intralobular and interlobular connective tissue is reduced. When a gland is inactive, only the duct system is evident. Cellular thickenings at the termination of intralobular ducts represent gland remnants or gland precursors in the inactive gland. Interlobular ducts, with a bistratified cuboidal to columnar lining, drain the lobules and lead to the lactiferous ducts and lactiferous sinuses at the base of the teat. The teat sinus, with a bistratified columnar to cuboidal lining, leads to the teat canal that opens onto the tip of the teat. The teat canal is lined by a stratified squamous epithelium that is continuous with the skin. Single teat sinuses and canals pass through the teats of ruminants, while the teats of carnivores, horses, and pigs contain multiple teat sinuses and canals, each opening separately onto the surface. The skin surface of the teat of cows and pigs lacks sebaceous glands, sweat glands, and hairs. Chestnuts and ergots are epidermal thickenings characteristic of the horse. The claws of carnivores, hooves of ungulates, and horns of ruminants are highly specialized derivatives of the skin composed of hard keratin.

CHICKEN

Skin

The epidermis of the chicken is generally thinner than that of mammals. It is composed of an inner stratum germinativum and an outer stratum corneum. The stratum germinativum includes a basal layer, an intermediate layer of one to several layers of polygonal cells, and a thin transitional layer of flat vacuolated cells just below the stratum corneum.

The dermis of feathered skin lacks papillae and is nonglandular. Mutlilocular as well as unilocular adipocytes occur in the subcutis.

Epidermal Derivatives

The epidermally derived feathers may be classified into three main types in the adult chicken: contour, down, and filoplume. A contour feather has a central shaft that is divisible into a hollow calamus (quill) and a rachis. A vane extends laterally from each side of the rachis and is composed of barbs and barbules with interlocking hooklets. Down feathers are soft and fluffy. Their barbules lack hooklets. Filoplumes are small, hairlike feathers.

Feathers are situated in tubelike follicles oriented obliquely in the dermis or subcutis. The follicle wall of a developing feather is lined by a stratum corneum and underlying stratum germinativum surrounded by a layer of connective tissue. The epidermal collar, a thick ring of epidermal cells at the base of the follicle, gives rise to the feather. It surrounds the dermal (feather) papilla, which gives rise to a well-vascularized, mesenchyme-like feather pulp that is present during growth of the feather. A network of feather muscles, each composed of one to several bundles of smooth muscle, attaches the follicles to each other. No muscles are associated with the follicles of filoplumes.

Wattles and combs are appendages of the skin whose dermis contains an extensive, superficial network of sinus capillaries and abundant mucous connective tissue. The sinus capillaries are responsible for the striking red color of the appendages.


Digital pads are covered by a thick stratum corneum and contain a cushion of adipose tissue in their subcutis. Scales, claws, and beaks are keratinized derivatives of the skin.

The uropygial (preen) gland is a bilobed holocrine gland located in the dorsal base of the tail. It produces an oily secretion. Simple tubular glands radiate outward from the lumen of each lobe like the bristles of a bottle brush. Each tubule is divided into a sebaceous zone and a glycogen zone, named according to their histochemical staining properties. The glycogen zone is continuous with the lumen of the lobe. Each lobe is drained by a primary duct that passes through the isthmus to the papilla (nipple) to open onto the surface.

| Word Ro | ots | |
|-------------------|---------------------------|--|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Arrect Pili | Steep, upright Hair | When the <i>arrector pili</i> contracts, it causes hairs to stand upright. |
| Corn or Kerato | Horn | The stratum <i>corneum</i> is formed by <i>keratinized</i> cells. |
| Cutis Sub | Skin Under or below | The <i>subcutis</i> is a layer of loose connective tissue below the skin. |
| Derm | Skin | The dermis is a layer of the skin. |
| Epi | Above, over | The <i>epidermis</i> is above the dermis. |
| Papilla | Nipple | The <i>papillary</i> layer of the dermis has nipple-like projections called dermal <i>papillae</i> . |
| Stratum | A layer | The <i>stratum</i> basale is the deepest layer of the epidermis. |



Figure 12.1 ×12.5



1. Dermal papilla
2. Dermis
3. Epidermal peg
4. Epithelium, hairy skin
5. Epithelium, planum
6. Hair follicle
7. Stratum basale

8. Stratum corneum
9. Stratum granulosum
10. Stratum lucidum
11. Stratum spinosum
12. Subcutis
13. Surface groove
14. Sweat gland


Figure 12.1. Planum Nasale and Hairy Skin, Nose, Dog. Junction of the hairless planum nasale (thick skin) and the hairy portion (thin skin) of the nose. No glands are associated with the planum of carnivores. There are surface grooves in the planum of the dog.

Figure 12.2. Planum Nasale and Hairy Skin, Nose, Cat. No hairs or glands are associated with the planum of carnivores. The surface of the planum bears numerous small, raised tubercles that are characteristic of the cat.

Figure 12.3. Epithelium, Planum Nasale, Cat. Portions of the small tubercles typical of the cat's planum. All layers of the epidermis are evident. Note how the papillae of the dermis interdigitate with the epidermal pegs.



Figure 12.3 × 90






Figure 12.5 ×625

- 1. Dermis
- 2. Epidermis
- 3. Hair follicle
- 4. Intercellular bridges
- 5. Melanocyte
- 6. Sebaceous gland
- 7. Sinus hair follicle
- 8. Stratum basale
- 9. Stratum spinosum
- 10. Sweat gland

Figure 12.4. Nose, Horse. Numerous small sebaceous glands, sweat glands, fine hairs, and the follicle of a sinus hair are evident. The epidermis is heavily pigmented.

Figure 12.5. Nose, Horse. A melanocyte with numerous pigment granules is located in the deep portion of the epidermis. Surrounding cells have phagocytized melanin granules produced by melanocytes. The granules are aggregated like a cap just above the nucleus of some of the cells of the stratum spinosum.








Figure 12.7 ×250




Figure 12.8 ×18




Figure 12.9 ×62.5


| rigure 12.9 | | ∧02. € |
|---|--|---------------|
| | KEY | |
| Dermal papilla Dermis Duct Epidermis Merocrine sweat gland Myoepithelial cell, nucleus | 7. Nasolabial gland 8. Stratum basale 9. Stratum corneum 10. Stratum spinosum 11. Surface groove | |
| | | |

Figure 12.6. Planum Rostrale, Pig. The very thick epidermis of the snout of the pig has low, wide elevations and an especially thick stratum corneum. Long dermal papillae project into the epidermis. Numerous merocrine sweat glands occur in the subcutis. Hairs, which are sparse on the planum of the pig, are not shown.

Figure 12.7. Planum Rostrale, Pig. Detail of the merocrine sweat glands. Secretory cells are either columnar or cuboidal and are surrounded by myoepithelial cells.

Figure 12.8. Planum Nasolabiale, Cow. The surface of the planum of the cow is hairless and marked by grooves. Long dermal papillae project into the thick epidermis. Glands are abundant in the subcutis of the planum of ruminants.

Figure 12.9. Planum Nasolabiale, Cow. The tissue was cut parallel to the surface of the planum, so that cross sections of dermal papillae appear in the stratum spinosum. Each papilla is surrounded by cells of the stratum basale and each contains several blood vessels.



×360 **Figure 12.10**




Figure 12.11 ×250





Figure 12.12



× 125 **Figure 12.13**

- 1. Dermis, papillary layer
- Dermis, reticular layer
- 3. Digital cushion
- 4. Duct of sweat gland5. Intercellular bridges
- 6. Intralobular duct
- 7. Keratinocyte nucleus
- 8. Merocriné sweat gland
- 9. Pacinian corpuscle
- 10. Secretory acinus
- 11. Skeletal muscle

Figure 12.10. Planum Nasolabiale, Cow. During tissue processing, the keratinocytes of the stratum spinosum shrink away from each other, but remain attached at multiple sites where desmosomes are located. As a result of this artifact, there appear to be cytoplasmic connections between adjacent cells. Some early workers called these "intercellular bridges" because they were thought to represent cytoplasmic connections between cells. Others thought the artifacts resembled spines, hence the name stratum spinosum.


Figure 12.11. Planum Nasolabiale, Cow. The planum of the cow, sheep, and goat contains many tubuloacinar serous glands. Branches of an intralobular duct can be seen entering secretory acini.

Figure 12.12. Digital Pad, Dog. The digital pad is hairless and covered by a very thick epidermis that is roughened by small conical projections in the dog. Compare with Figure 12.14.

Figure 12.13. Digital Pad, Dog. Coiled merocrine sweat glands and pacinian corpuscles among skeletal muscle and loose connective tissue of the digital pad.



Figure 12.14 ×52



| | KEY |
|----------------------------|-----------------------|
| 1. Dermis, papillary layer | 6. Stratum basale |
| 2. Dermis, reticular layer | 7. Stratum corneum |
| 3. Digital cushion | 8. Stratum granulosum |
| 4. Duct of sweat gland | 9. Stratum lucidum |
| 5 Enidermis | 10 Stratum spinosum |

Figure 12.14. Digital Pad, Cat. The surface of the digital pad of the cat is smooth, lacking the conical papillae that are typical of the dog (Figure 12.12). Portions of the excretory ducts of sweat glands spiral through the stratified squamous epithelium.

Figure 12.15. Digital Pad, Cat. Detail of the epidermis and dermis shown in Figure 12.14. All five layers of the epidermis are evident.




Figure 12.16

| KEY | | |
|--|--|--|
| Dermis Hair, x.s. Stratum basale | 4. Stratum corneum5. Stratum granulosum6. Stratum spinosum | |

Figure 12.16. Skin, Mid-ventral Abdomen, Dog. The epidermis is thin and consists of four layers. Note that only a few layers of cells comprise the stratum spinosum. The stratum corneum is also relatively thin, and the keratinized cells have loosened and separated from the surface.

Summary of the Composition of the Skin and Subcutis

Skin:

A. Epidermis: A keratinized, stratified squamous epithelium

- 1. Stratum basale: A single layer of cuboidal to columnar cells that lies on the basement membrane.
- 2. Stratum spinosum: A layer of variable thickness formed of polygonal cells that become flattened toward the surface.
- 3. Stratum granulosum: Flattened cells with basophilic cytoplasmic granules.
- 4. Stratum lucidum: Thin, pale, translucent layer present in regions where the epidermis is very thick.5. Stratum corneum: Outermost layer; composed of dead, keratinized squamous cells.

B. Dermis: Connective tissue below the epidermis formed of two layers

- 1. Papillary (superficial) layer: Loose connective tissue just deep to the epidermis; includes projections, dermal papillae, that interdigitate with the epidermis.
- 2. Reticular (deep) layer: Dense, irregular connective tissue.

Subcutis (Hypodermis; Superficial Fascia)

- A layer of loose connective tissue, often rich in adipose tissue, below the skin.
- Joins the dermis of the skin to underlying structures, such as muscle and bone.




Figure 12.17 ×52




Figure 12.18 ×125




Figure 12.19 ×125

- 1. Arrector pili muscle
- 2. Common follicular opening
- 3. Dermis, deep
- 4. Dermis, superficial
- 5. Epidermis
- 6. Hair
- 7. Hair bulb

- 8. Sebaceous gland
- 9. Skeletal muscle
- 10. Stratum basale
- 11. Stratum corneum
- 12. Stratum spinosum cell, nucleus
- 13. Subcutis

Figure 12.17. Skin, Back, Cat. Two compound follicles are visible in the dermis. In carnivores the hairs of compound follicles merge at the level of the sebaceous glands and share a common follicular opening to the surface. Bits of hairs are evident in the follicles as shiny, yellow-brown structures. The arrector pill muscles of the skin of the back are especially well developed in cats and dogs. A space artifact separates the subcutis from the underlying skeletal muscle.

Figure 12.18. Skin, Caudal Abdomen, Cat. The epidermis is extremely thin. Cells of the stratum spinosum are sparse, and those of the stratum granulosum are visible only as occasional dark granular areas just beneath the stratum corneum. Hairs are visible within the compound follicles.

Figure 12.19. Skin, Back, Cat (Masson's). Portions of an arrector pill muscle, sebaceous gland, and a hair within a follicle.




Figure 12.20 ×52

| KEY | | |
|--|--|--|
| Dermis Epidermis Hair Hair bulb | 5. Hair follicle 6. Sebaceous gland 7. Sweat gland | |

Figure 12.20. Skin, Neck, Horse. Simple hair follicles occur in the skin of noncarnivores.




Figure 12.21 ×360




Figure 12.22 ×125

- 1. Connective tissue sheath
- 2. Cortex, hair
- 3. Cuticle, hair
- 4. Cuticle, inner root sheath
- 5. Dermal papilla
- 6. Dermis

- 7. Inner root sheath
- 8. Medulla, hair
- 9. Outer root sheath
- 10. Pyknotic nucleus
- 11. Sweat gland

Figure 12.21. Skin, Neck, Horse. Mid-region of a longitudinal section of a hair follicle. Large, clear cells of the medulla of the hair have round nuclei that become pyknotic as they progress distally from the hair bulb. The pigment-laden cortex of the hair is formed from closely packed elongated cells that have become keratinized. Scalelike, keratinized cells of the cuticle of the hair partially overlap so that their free edges point upward. They interlock with cells of the cuticle of the inner root sheath, whose free edges are directed downward.

Figure 12.22. Skin, Neck, Horse. A dermal papilla projects into the hair bulb at the base of the follicle. Cells of the cortex of the hair are nearly obscured by pigment granules provided by melanocytes of the bulb.




Figure 12.23




Figure 12.24




Figure 12.25 ×62.5




Figure 12.26 ×12.5



Figure 12.27

| | KEY |
|------------------------------|------------------------|
| 1. Adipose tissue | 9. Hair |
| 2. Collagenous fiber | 10. Hair bulb |
| 3. Dartos muscle | 11. Sebaceous gland |
| 4. Dermis | 12. Stratum basale |
| 5. Dermis, deep layer | 13. Stratum corneum |
| 6. Dermis, superficial layer | 14. Stratum granulosum |
| 7. Elastic fiber | 15. Stratum spinosum |
| 8. Epidermis | 16. Sweat gland |

Figure 12.23. Skin, Dorsal Neck, Pig. Note the extremely thick dermis. Sweat glands and adipose tissue are seen in the subcutis.

Figure 12.24. Skin, Dorsal Neck, Pig. The epidermis and part of the dermis are shown in detail.

Figure 12.25. Skin, Dorsal Neck, Pig (Orcein). The dermis of the skin contains numerous branching elastic fibers. The fibers of the superficial layer are fine, while those of the deep layer are coarse.

Figure 12.26. Skin, Back, Sheep. Various portions of numerous hair follicles are embedded in the thick superficial layer of the dermis. The hair follicles of sheep tend to be arranged vertically, rather than diagonally, in the dermis. Compare with Figures 12.17 and 12.20.

Figure 12.27. Scrotum, Goat. The epidermis of the scrotum is remarkably thin. Portions of two simple hair follicles are located in the dermis. Bundles of smooth muscle among fibroelastic tissue in the dermis comprise the tunica dartos.




Figure 12.28 ×125




Figure 12.29 ×12.5




Figure 12.30 ×62.5




Figure 12.31 ×250




Figure 12.32 ×250

| 1. Adipose tissue 2. Blood vessel 3. Capillary 4. Connective tissue sheath 5. Cortex, hair 6. Dermis, superficial layer 7. Duct of sweat gland 8. Elastic cartilage 9. Epidermis 10. Enidermis inner surface 11. Epidermis, outer surface 12. Hair follicle 13. Inner root sheath 14. Medulla, hair 15. Outer root sheath 16. Primary hair 17. Sebaceous gland 18. Smooth muscle 19. Venule | | |
|---|--|--|
| 2. Blood vessel 3. Capillary 4. Connective tissue sheath 5. Cortex, hair 6. Dermis, superficial layer 7. Duct of sweat gland 8. Elastic cartilage 9. Epidermis 12. Hair follicle 13. Inner root sheath 14. Medulla, hair 15. Outer root sheath 16. Primary hair 17. Sebaceous gland 18. Smooth muscle 19. Venule | | KEY |
| 10. Epideimis, illier soridee | Blood vessel Capillary Connective tissue sheath Cortex, hair Dermis, superficial layer Duct of sweat gland Elastic cartilage | 12. Hair follicle 13. Inner root sheath 14. Medulla, hair 15. Outer root sheath 16. Primary hair 17. Sebaceous gland 18. Smooth muscle |

Figure 12.28. Scrotum, Goat. The thin epidermis and a portion of a hair follicle are shown in detail.

Figure 12.29. Pinna (Auricle), External Ear, Dog. A plate of elastic cartilage is covered by the skin of the outer (convex) and inner (concave) surfaces of the pinna. Hair follicles are more numerous in the skin of the outer surface.

Figure 12.30. Pinna, External Ear, Dog. Clusters of compound hair follicles cut in cross section vary in appearance at different levels of the dermis. The cells of the cortex and medulla of the hairs are evident in the deepest portions of the follicles. More superficially, the cells become keratinized and appear shiny pink (medulla) and yellow (cortex). Several hairs have merged to share a common follicle wall near the epidermis.

Figure 12.31. Pinna, External Ear, Dog. Detail of follicles, shown in cross section, from the deep region of the dermis, similar to those in Figure 12.30.

Figure 12.32. Skin, Back, Sheep. A wool hair, shown in cross section, lacks a medulla.





Figure 12.33 ×18



Figure 12.34



| 13 | |
|---|---|
| igure 12.35 | ×125 |
| K | EY |
| Annular sinus Connective tissue sheath, inner Connective tissue sheath, outer Endothelial cell, nucleus Epidermis Erythrocytes Hair | 8. Hair bulb 9. Inner root sheath 10. Merkel's cell 11. Outer root sheath 12. Sebaceous gland 13. Sinus pad 14. Trabecula |
| | |

Figure 12.33. Sinus Hair Follicle, I.s., Nose, Cow. The entire length of the blood-filled sinus is crossed by numerous trabeculae in sinus hair follicles of ruminants, horses, and pigs.

Figure 12.34. Sinus Hair Follicle, Oblique Section, Nose, Dog. The large sinus hair follicle contains a blood-filled sinus, lined by an endothelium, between the inner and outer layers of the connective tissue sheath. In carnivores only the lower region of the sinus is spanned by a network of trabeculae of connective tissue. The upper region contains an annular sinus, free of trabeculae, into which protrudes a thickening of the inner connective tissue sheath called the sinus pad.

Figure 12.35. Sinus Hair Follicle, Nose, Dog. Detail of Figure 12.34. Portion of the sinus pad, annular sinus, and trabeculated sinus. Note the Merkel's cells, associated with tactile stimulation, in the external (outer) root sheath.




Figure 12.36 ×125




Figure 12.37 ×62.5




Figure 12.38 ×250




Figure 12.39 ×250




Figure 12.40 ×125

| | KEY |
|--------------------------------|------------------------------------|
| 1. Capillary | 9. Outer root sheath |
| 2. Duct | 10. Pyknotic nucleus |
| 3. Follicular folds | 11. Sebaceous gland |
| 4. Gland cells, surface cut | 12. Secretory portion, sweat gland |
| 5. Hair | 13. Stem cell |
| 6. Hair follicle | 14. Sweat gland, l.s. |
| 7. Inner root sheath | 15. Sweat gland, x.s. |
| 8. Myoepithelial cell, nucleus | 16. Vacuolated cell |

Figure 12.36. Sebaceous Gland and Hair Follicle, I.s., Lip, Sheep. The inner root sheath forms follicular (circular) folds below the entrance of sebaceous glands into the follicle.

Figure 12.37. Nose, Dog. Longitudinal sections of two serpentine sweat glands. Their tubular structure is evident.

Figure 12.38. Sebaceous Gland and Sweat Gland, Nose, Dog. The secretory portion of a sweat gland, lined by cuboidal to columnar cells, is continuous with the bistratified, flattened cells of its duct. The cell types found in the holocrine sebaceous gland are evident: small, flat, peripheral stem cells; maturing, round cells with pale, vacuolated cytoplasm; inner degenerating cells with pyknotic nuclei.

Figure 12.39. Sweat Gland, Skin, Horse (Trichrome). Cross and oblique sections of a coiled sweat gland in the dermis are lined by cuboidal to columnar cells and surrounded by myoepithelial cells. The solid sheet of several cells represents a surface cut through the wall of the gland.

Figure 12.40. Sweat Gland and Duct, Teat, Sheep. Low columnar secretory cells with apical blebs end abruptly where the duct epithelium begins. Both the secretory cells and the initial segment of the duct are surrounded by myoepithelial cells. The cytoplasm of the myoepithelial cells appears as a pink, sometimes rippled band.




Figure 12.41 ×125




Figure 12.42 ×250




Figure 12.43 ×12.5




Figure 12.44 ×250




Figure 12.45 ×25

| | KEY |
|------------------------|---|
| 1. Basement membrane | 8. Hair follicle |
| 2. Clear cell | Myoepithelial cell, cytoplasm |
| 3. Dark cell | 10. Myoepithelial cell, nucleus, l.s. |
| 4. Dermis | 11. Myoepithelial cell, nucleus, x.s. |
| 5. Duct of sweat gland | 12. Sebaceous gland |
| 6. Epidermis | 13. Secretory cell, nucleus |
| 7. Hair bulb | 14. Sweat gland |


Figure 12.41. Sweat Gland, Teat, Sheep. The secretory cells of sweat glands vary from squamous to tall columnar. They are squamous in this preparation, but are columnar in Figure 12.40. Note that one of the secretory portions is cut tangentially, revealing the elongated shape of the myoepithelial cells.

Figure 12.42. Sweat Gland, Teat, Sheep (Silver and Eosin). The basement membrane of a sweat gland is blackened with silver. Myoepithelial cells occur between the flattened secretory cells and the basement membrane.

Figure 12.43. Carpal Gland, Pig. Lobules of merocrine sweat glands occur in the subcutaneous tissue on the medial side of the carpus of the pig.

Figure 12.44. Carpal Gland, Pig. Dark and clear cells of the secretory units of these merocrine sweat glands are surrounded by myoepithelial cells.

Figure 12.45. Infraorbital Pouch (Sinus), Sheep. Many large sebaceous glands occupy the wall of the infraorbital pouch of sheep. Some apocrine sweat glands lie deep to the sebaceous glands.






Figure 12.47 ×12.5




Figure 12.48 ×2:




Figure 12.49 ×125




Figure 12.50 ×25

| KEY | | |
|---|--|--|
| Corpora amylacea Epidermis Gland Glandular epithelium Hair follicle | 8. Intralobular connective tissue 9. Intralobular duct 10. Lobule 11. Lymphatic vessel 12. Sebaceous gland | |
| 6. Interlobular connective tissue7. Interlobular duct | 13. Secretory unit 14. Sweat gland | |

Figure 12.46. Inguinal Pouch, Sheep. The skin of the inguinal pouch of sheep contains a few hairs, sebaceous glands, and an abundance of apocrine sweat glands.

Figure 12.47. Mammary Gland, Inactive, Cat. Lobules of glandular tissue and ducts are surrounded by fibroelastic subcutaneous tissue. The overlying skin contains a few hairs. Large lymphatic vessels lie deep to the parenchyma.

Figure 12.48. Mammary Gland, Inactive, Cow. Abundant interlobular connective tissue and components of the duct system are evident in an inactive gland. Interlobular ducts branch into the lobules as intralobular ducts.

Figure 12.49. Mammary Gland, Inactive, Cow. Lobules are composed of intralobular ducts and intralobular connective tissue, which is moderately rich in cells. Thickenings at the terminations of intralobular ducts represent remnants or precursors of glandular epithelium. When these are cut in cross section, they cannot always be distinguished from ducts.

Figure 12.50. Mammary Gland, Active, Cow. In the active gland, secretory parenchyma is well developed and connective tissue is reduced. Compare with Figure 12.48. The lumens of the secretory glands and ducts are filled with secretion (deep pink).




Figure 12.51 ×125




Figure 12.52 ×250




Figure 12.53 ×12.5




Figure 12.54 ×62.5




Figure 12.55 ×12.5

- 1. Connective tissue
- 2. Corpora amylacea
- 3. Epidermis
- 4. Intralobular connective tissue
- 5. Myoepithelial cell, nucleus
- 6. Secretion

- 7. Secretory cell, nucleus
- 8. Secretory unit
- 9. Smooth muscle
- 10. Sweat gland
- 11. Teat sinus

Figure 12.51. Mammary Gland, Active, Cow. A portion of a lobule containing numerous tubuloacinar secretory units. Some of the alveoli contain round concretions of casein and cellular debris called corpora amylacea.

Figure 12.52. Mammary Gland, Active, Cow. Many secretory cells have basally displaced nuclei and indistinct lateral cell borders. These cells appear pale because their cytoplasmic lipids have been extracted. Sloughed cells, whose dark nuclei are visible in the lumens, are part of the secretory product. Some of the flat nuclei surrounding the alveoli belong to myoepithelial cells.

Figure 12.53. Teat, x.s., Dog. A portion of the teat shows numerous sinuses among intermingling bundles of smooth muscle and fibroelastic connective tissue. Nonruminants have multiple teat sinuses and teat canals. Some glands and hairs are associated with the skin of the teat of carnivores, horses, sheep, and goats.

Figure 12.54. Teat Sinus, x.s., Dog. Detail of a teat sinus of Figure 12.53 reveals a highly folded lining. Glandular areas, composed of small secretory units, are associated with the wall of the sinus.

Figure 12.55. Teat Sinus, x.s., Cat. This cross section through a teat reveals five teat sinuses. Some of these contain a secretion that is stained pink.




Figure 12.56 ×12




Figure 12.57 ×12.5





Figure 12.58 ×180



igure 12.59 ×25

| K | EY | |
|-------------------------------------|----------------------|--|
| 1. Bistratified columnar epithelium | 8. Hair follicle | |
| 2. Bistratified cuboidal epithelium | 9. Lamina propria | |
| 3. Blood vessel | 10. Lymphatic vessel | |
| 4. Collagenous band | 11. Sebaceous gland | |
| 5. Elastic fibers | 12. Smooth muscle | |
| 6. Epidermis | 13. Sweat gland | |
| 7. Epithelium | 14. Teat sinus | |

Figure 12.56. Teat Sinus, x.s., Horse (Orcein). A band of collagenous fibers lies between the epithelium and the underlying fibroelastic connective tissue.

Figure 12.57. Teat Sinus, x.s., Cow. The mucosa of the teat sinus blends with the middle layer of the teat. The latter contains well-developed, longitudinally oriented blood vessels (cut in x.s.), bundles of smooth muscle, fibroelastic tissue, and lymphatic vessels. The outer layer, the skin surface, is not shown.

Figure 12.58. Teat Sinus, x.s., Cow. The teat sinus is lined by a bistratified cuboidal to columnar epithelium.

Figure 12.59. Teat Sinus, x.s., Sheep, Male. The skin of the teat contains hairs, sebaceous glands, and sweat glands, except in the cow and pig. Compare with Figure 12.63.




Figure 12.60 ×62.5




Figure 12.61 ×12.5




Figure 12.62 ×62.5




Figure 12.63 ×25

- 1. Bistratified columnar epithelium
- 2. Bistratified cuboidal epithelium
- 3. Blood vessel
- 4. Dermis
- 5. Epidermis
- 6. Epithelium

- 7. Lamina propria
- 8. Smooth muscle
- 9. Stratified squamous epithelium
- 10. Stratum basale
- 11. Stratum corneum
- 12. Stratum spinosum

Figure 12.60. Teat Sinus and Canal, Junction, x.s., Horse. Patches of the bistratified epithelium (columnar and cuboidal) of the teat sinus intermingle with the stratified squamous epithelium of the teat canal.

Figure 12.61. Teat Canal, x.s., Cow (Trichrome). The keratinized stratified squamous lining of the teat canal is encircled by a papillated layer (green) of connective tissue and bundles of smooth muscle (pale yellow).

Figure 12.62. Teat Canal, x.s., Cow (Trichrome). Detail of the thick, keratinized, stratified squamous epithelium and the surrounding connective tissue and smooth muscle shown in Figure 12.61.

Figure 12.63. Skin Surface, Teat, x.s., Cow (Trichrome). The skin surface of the teat of the cow and pig is hairless.




Figure 12.64 ×26



Figure 12.65 ×12.5




Figure 12.66 ×12.5

KEY

- 1. Claw fold
- 2. Dermis
- 3. Digital pad
- 4. Distal phalanx
- 5. Epidermis, digital pad
- 6. Laminae

- 7. Limiting furrow
- 8. Merocrine sweat gland
- 9. Middle phalanx
- 10. Sole
- 11. Wall

Figure 12.64. Developing Claw, I.s., Fetus, Dog. The claw of carnivores consists of a dorsal and lateral wall (body, claw plate) and a ventral sole of hard keratin that cover the distal phalanx. The claw fold is the skin that covers the wall at the base of the claw. Endochondral bone formation has begun in the phalanges of this specimen.

Figure 12.65. Sole of Claw and Digital Pad, Dog. The limiting furrow separates the digital pad from the sole of the claw.

Figure 12.66. Apex of Claw, I.s., Dog. The dermis of the wall bears laminae (lamellae) at the apex of the claw.



Figure 12.67 ×12.5




Figure 12.68 ×12.5




Figure 12.69 ×62.5




Figure 12.70 ×125

| 1/ | 7 | • | ú |
|----|---|---|---|
| | | | |
| | | | |

1. Dermal papilla

2. Dermis

3. Hair follicle

4. Horn tubule

5. Intertubular horn

- 6. Stratum basale
 - 7 (. .
 - 7. Stratum corneum8. Stratum granulosum
 - 9. Stratum spinosum
- Figure 12.67. Horn, Cow. Horns of ruminants are composed of bone of the cornual process covered by a dermis and epidermis. The epidermis with a thick stratum corneum of hard keratin (horn) and a portion of the underlying papillated dermis are shown here. (Photograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 12.68. Chestnut, Horse. Junction of the hairy skin and chestnut. Chestnuts (and ergots) of horses are keratinized thickenings of the epidermis composed of horn tubules (tubular horn) and intertubular horn. Horn tubules arise from the cells of the stratum basale that cover the apex and sides of dermal papillae. Intertubular (interpapillary) horn arises from the cells of the stratum basale that are located between the bases of the dermal papillae. Only a small portion of the very thick stratum corneum of the chestnut is shown.

Figure 12.69. Chestnut, Horse. Detail of a portion of the epidermis.

Figure 12.70. Chestnut, Horse. The section was cut parallel to the surface of the chestnut at the level of the stratum corneum. Horn tubules appear in cross section between intertubular horn.

THE EQUINE HOOF

The equine hoof is the keratinized portion of the epidermis that covers the distal end of the digit. The various regions of the hoof are depicted in Figures 12.71 and 12.72. The perioplic, coronary, and laminar regions comprise the wall (the portion of the hoof that is visible when the digit is on the ground). The wall turns in ventrally at an acute angle to form the bars. The sole, which forms most of the ventral surface of the hoof, is attached to the bars and the adjacent, inner border of the wall. The frog, a caudal, edge-shaped mass, lies between the bars. The apex of the frog merges with the sole cranially. The bulbs are the convex protuberances located above and behind the frog.

The keratinized tissue comprising the hoof is in the form of tubular, intertubular, and laminar horn (Figures 12.75, 12.76, 12.79, and 12.80). The underlying, living layers of the epidermis include the stratum spinosum, whose cells are undergoing keratinization, and the stratum basale. The stratum basale borders on the dermis (corium), which is rich in blood vessels and nerves. The dermis may be papillated or laminated, depending on whether the overlying epidermis contains tubular or laminated horn, respectively. The dermis blends with underlying structures such as the subcutaneous cushions and the periosteum of the third phalanx.

The perioplic epidermis forms a band of soft, nonpigmented tubular horn. It merges with the epidermis of the skin above and extends downward as a thin, glossy, flaky layer of keratin that forms the outer coating of the wall of the hoof, called the **stratum tectorium** (stratum externum). This layer is well developed in young animals, but tends to be worn away in older horses. The perioplic epidermis

widens at the heels to form the bulbs. The **perioplic dermis** is characterized by the presence of fine, short, papillae (1 to 2 mm).

The germinal cells of the **coronary epidermis** form horn tubules (tubular horn) and intertubular horn that extend from the coronary region to the ground surface, forming the bulk of the wall of the hoof, the **stratum medium**. The horn tubules are orientated at an angle to the ground. They parallel the external surface of the hoof. The **coronary dermis** is marked by long dermal papillae (4 to 6 mm).

The laminar epidermis of the wall is in the form of laminae (lamellae) that are arranged parallel to the horn tubules of the stratum medium. They extend from the deep edge of the coronary region to the sole. Each primary lamina bears numerous secondary laminae (not present in the hooves of pigs and ruminants) that project at right angles along its length. The primary epidermal laminae are keratinized and fused with the inner portion of the stratum medium of the wall. The secondary epidermal laminae consist of a core of cells of the stratum spinosum bordered by cells of the stratum basale. The epidermal laminae form the stratum internum of the wall of the hoof. They interdigitate with primary and secondary dermal laminae of the laminar dermis. This extensive interdigitation serves to suspend the third phalanx from the hoof. At the ground surface, the junction of the epidermal laminae of the wall (unpigmented) with the sole is called the white line.

The tubular and intertubular horns of the bulbs, sole, and frog are softer than that of the wall of the hoof. The dermis of these regions, like that of the periople and coronary region, is papillated. The epidermis and dermis of the bars are laminated, being continuous with the laminar region of the wall.




Figure 12.71. Hoof, Horse. The various regions of the hoof are shown. The inner surface of the periople and coronary region and that of the sole, frog, and bulbs are stippled in the drawing. In the intact toe, dermal papillae extend into the funnel-shaped depressions whose openings are represented by the stipples.




Figure 12.72. Sole, Hoof, Horse. The sole forms most of the ventral surface of the hoof. It is bordered by the bars, the apex of the frog, and the inner border of the wall of the hoof.




Figure 12.73 ×12.5




Figure 12.74 ×25




Figure 12.75 ×125





Figure 12.76 $\times 12.5$



| | KEY |
|-----------------------------|------------------------------------|
| 1. Bone, P3 | 10. Hair follicle, developing |
| 2. Cartilage, developing P3 | 11. Horn tubule |
| 3. Coronary dermis | 12. Horn tubule, cortex |
| 4. Coronary epidermis | 13. Horn tubule, medulla |
| 5. Dermal papilla | 14. Intertubular horn |
| 6. Dermis, sole | 15. Laminar dermis |
| 7. Dermis, wall | Perioplic dermis |
| 8. Epidermis, sole | 17. Perioplic epidermis |
| 9. Epidermis, wall | · · |

Figure 12.73. Developing Hoof, I.s., Fetus, Horse. The regions that form the three layers of the hoof wall are apparent: the perioplic, coronary, and laminar regions.

Figure 12.74. Developing Hoof, Coronary Region, I.s., Fetus, Horse. Portion of coronary epidermis and coronary dermis, later in development than in Figure 12.73, showing tubular and intertubular horn. The medulla and cortex of the horn tubules are formed from the cells of the stratum basale that cover the tip and the sides of dermal papillae, respectively. Intertubular horn is formed by cells of the stratum basale that are located between the bases of the dermal papillae. (Photograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 12.75. Developing Hoof, Coronary Region, I.s., Fetus, Horse. Detail of Figure 12.74, showing two horn tubules in longitudinal section. (Photograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 12.76. Hoof, Coronary Region, x.s., Horse. Dermal papillae and horn tubules of tubular horn. (Photograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 12.77. Developing Hoof, Wall and Sole, x.s., Fetus, Horse. The dermis of the wall is laminated, while that of the sole is papillated.




Figure 12.78 ×125




Figure 12.79 ×12.5



- 1. Basal cell
- 2. Blood vessel
- 3. Bone, P3
- 4. Dermal lamina, primary
- 5. Dermal lamina, secondary
- 6. Epidermal lamina, primary
- 7. Epidermal lamina, secondary
- 8. Horn tubule
- 9. Laminar dermis
- 10. Stratum medium

Figure 12.78. Developing Hoof, Wall, x.s., Fetus, Horse. Detail of Figure 12.77. The epidermal laminae, at this time, consist mainly of a layer of basal cells (stratum basale). Primary epidermal laminae have begun to form secondary laminae. The primary and secondary dermal laminae are extensions of the laminar dermis.

Figure 12.79. Hoof, Laminar Region, x.s., Horse. Horn tubules of the stratum medium are seen in cross section. Primary and secondary epidermal laminae of the stratum internum interdigitate with the laminar dermis, which anchors the third phalanx to the wall of the hoof. Epidermal laminae, which are long ridges, appear featherlike in cross section. (Photograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 12.80. Hoof, Laminar Region, x.s., Horse. Primary epidermal laminae of the stratum internum, continuous with the stratum medium, bear secondary epidermal laminae. These interdigitate with primary and secondary dermal laminae. The secondary epidermal laminae and the dermal laminae comprise the sensitive laminae. Nuclei of the basal cells appear as small dark spots along the periphery of secondary epidermal laminae. (Photograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)




Figure 12.80 ×90




Figure 12.81 ×25




Figure 12.82 ×125




Figure 12.83 ×62.5



Figure 12.84 ×25




Figure 12.85 ×250

| KEY | | |
|------------------------|--------------------------|--|
| 1. Axial blood vessel | 10. Feather muscle | |
| 2. Barbs, pigmented | 11. Feather pulp | |
| 3. Corneous cells | 12. Feather sheath | |
| 4. Corneous connection | 13. Horn tubule | |
| 5. Dermal papilla | 14. Intertubular horn | |
| 6. Dermis | 15. Melanocyte | |
| 7. Epidermal collar | 16. Stack of nuclei | |
| 8. Epidermis | 17. Stratum corneum | |
| 9. Feather follicle | 18. Stratum germinativum | |

Figure 12.81. Hoof, Sole, Oblique Section, Horse. Dermal papillae and horn tubules of the sole are shown. (Photograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 12.82. Skin, Neck, Chicken. The epidermis of feathered skin is very thin and composed of a stratum germinativum and stratum corneum. The layers of the stratum germinativum are evident in Figure 12.92. Nuclei of the epidermal cells are often organized into stacks perpendicular to the surface. Abundant small blood vessels appear in the superficial region of the dermis.

Figure 12.83. Feather Follicle, Skin, Neck, Chicken. Oblique section through the basal region of a follicle with a developing feather. An epidermal collar surrounds the dermal papilla. The upper portion of the dermal papilla blends with the feather pulp.

Figure 12.84. Skin, Chicken. Oblique sections of developing contour feathers.

Figure 12.85. Skin, Chicken. Oblique section of a developing contour feather. Melanocytes lie among cells of the barb. Barbule cells, beginning with the outermost ones, receive pigment from the processes of the melanocytes.






Figure 12.86 ×125





12. Stratum germinativum, skin

Figure 12.86. Skin, Chicken. Longitudinal section of a contour feather showing several developing barbs, later in development than in Figure 12.85. The pale, cuboidal cells at the base of each barb form the stem of the barb.

5. Dermis

6. Feather muscle

Figure 12.87. Skin, Chicken. Portion of contour feather follicle, x.s., at level of calamus (quill). Note that feather pulp has been replaced by an air space.

Figure 12.88. Skin, Chicken. Cross section of a contour feather showing numerous barbs.






Figure 12.88 ×90










Figure 12.91 $\times 12.5$




Figure 12.92 ×250

| KEY | | |
|--|---|--|
| Basal layer Calamus (quill) Dermis, central layer Dermis, superficial layer Elastic tendon | 10. Follicle, contour feather11. Follicle, filoplume feather12. Intermediate layer13. Mucous connective tissue14. Sinus capillary | |
| Endothelial cell, nucleus Epidermis Erythrocyte Feather muscle | 15. Stratum corneum16. Stratum germinativum, skin17. Transitional layer | |

Figure 12.89. Skin, Eyelid, Chicken. Feather follicles, x.s. The follicle wall of the tiny filoplume feather is relatively thick. A feather muscle attaches to the sheath of connective tissue of the follicle of a contour feather by an elastic tendon.

Figure 12.90. Wattle, x.s., Rooster. Numerous sinus capillaries in the superficial layer of the dermis impart a red color to the wattle (and comb) when filled with blood. Mucous connective tissue of the intermediate layer of the dermis surrounds the central layer of dense connective tissue.

Figure 12.91. Comb, Rooster. The point of a comb is similar in appearance to the wattle (Figure 12.90). The collagenous fibers of the central layer of the dermis arise from the periosteum of the skull and carry vessels and nerves to the extremities of the comb. These vessels of the comb are shown in Figure 12.92.

Figure 12.92. Comb, Rooster. Portion of epidermis and superficial dermis. The stratum corneum and layers of the stratum germinativum (basal, intermediate, and transitional layers) are evident. The lower ends of the cells of the basal layer bulge into the dermis, so that the epidermal-dermal boundary is uneven. Numerous anastomosing, blood-filled sinus capillaries in the superficial layer of the dermis are lined by pigment-laden endothelial cells.








Figure 12.94 ×12.5




Figure 12.95 ×12.5




Figure 12.96 ×250




Figure 12.97 ×25

| KEY | |
|-------------------------------|------------------------------|
| 1. Adipose tissue | 12. Merkel's cell |
| 2. Blood Vessel | 13. Mucous connective tissue |
| 3. Bone | 14. Oral cavity |
| 4. Dermal papilla | 15. Palatine ridge |
| 5. Dermis, central layer | 16. Periosteum |
| 6. Dermis, intermediate layer | 17. Sinus capillary |
| 7. Dermis, superficial layer | 18. Sinus capillaries |
| 8. Epidermal peg | 19. Stack of nuclei |
| 9. Epidermis | 20. Stratum corneum |
| 10. Fibroblast | 21. Stratum germinativum |
| 11. Herbst corpuscle | 22. Tomial edge |

Figure 12.93. Comb, Hen. The comb of a laying hen, compared with that of a rooster, contains less mucous connective tissue and more dense connective tissue in the intermediate layer, as well as fewer and smaller sinus capillaries in the superficial layer of the dermis (Figure 12.94).

Figure 12.94. Comb, Hen. Detail of the epidermis and a portion of the dermis in Figure 12.93. Fewer and smaller sinus capillaries are in the superficial layer of the dermis of the comb of the laying hen as compared with that of the rooster. Note the arrangement of the nuclei of epidermal cells into stacks.

Figure 12.95. Comb, Hen. Numerous Merkel's cells are located along the inner surface of the epidermis. These cells are associated with tactile nerve endings.

Figure 12.96. Upper Beak, x.s., Chicken. One side of the upper beak is shown. The bone of the premaxilla is covered by a periosteum, dermis, and epidermis with a thick layer of hard keratin. The dermis of the lateral surface of the upper beak often contains Herbst corpuscles; one corpuscle is shown here. See Figure 9.35 for detail of this corpuscle. Dermal papillae of the lateral surface diminish medially. The lower beak slips inside the upper beak between the palatine ridge and the stratum corneum of the medial surface.

Figure 12.97. Upper Beak, x.s., Chicken. Detail of dermis and epidermis of the tomial edge of Figure 12.96. Cells of the stratum basale vary in height and width, so that intermittent groups of tall, slender cells form epidermal pegs between which project dermal papillae.




Figure 12.98 ×25




Figure 12.99 ×250




Figure 12.100 ×12.5



Figure 12.101 ×62.5




Figure 12.102 ×25

- 1. Adipose tissue, subcutis
- 2. Dermis
- 3. Keratin, hard
- 4. Keratin, soft
- 5. Stratum corneum
- 6. Stratum germinativum
- 7. Sulcus

Figure 12.98. Spur, Hen. The spur cap consists of an extremely thick stratum corneum of hard keratin.

Figure 12.99. Scutes, I.s., Anterior Metatarsus, Chicken. Scutes are large scales that are covered by hard keratin. Their region of overlap, shown here, forms a sulcus lined by soft keratin.

Figure 12.100. Scutes, I.s., Anterior Metatarsus, Chicken. The region of overlap of two scutes shows the transition from the hard keratin to the deeper-staining, soft keratin of the sulcus.

Figure 12.101. Reticulate Scale, x.s., Digit, Chicken. The hard keratin of the stratum corneum covers the outer surface and lines the sulci of these small scales from the lateral metatarsus.

Figure 12.102. Digital Pad, Chicken. A thick keratinized epithelium, a dermis, and a thick cushion of adipose tissue in the subcutis characterize the digital pad.




Figure 12.103 ×12.5




Figure 12.104 ×62.5




Figure 12.105 ×25




Figure 12.106 ×12.5



Figure 12.107 ×250

| KEY | | |
|---------------------------------|-------------------------------------|--|
| 1. Bone, distal phalanx | 8. Multilocular fat cell, nucleus | |
| 2. Capsule of connective tissue | 9. Sebaceous zone | |
| 3. Dermis | 10. Stratum germinativum | |
| 4. Dorsal plate, claw | 11. Tendon | |
| 5. Footpad | 12. Unguinal scale, dorsal surface | |
| 6. Glycogen zone | 13. Unguinal scale, ventral surface | |
| 7. Lobe, lumen | 14. Ventral plate, claw | |
| | | |

Figure 12.103. Multilocular Fat, Digital Pad, Chicken. Multilocular adipocytes, containing numerous lipid vacuoles and a central nucleus, are common in the subcutis of the chicken.

Figure 12.104. Claw, l.s., Chicken. The dotted line indicates the approximate location of the drawing (cross section) of the claw shown in Figure 12.105.

Figure 12.105. Claw, Base, x.s., Chicken. The region of the base of the claw indicated with a dotted line in Figure 12.104 is represented in cross section in this drawing.

Figure 12.106. Claw, Base, x.s., Chicken. Compare this photomicrograph with Figures 12.104 and 12.105. The free edge of the dorsal, unguinal scale (scute type) overlaps the base of the claw, so that a cross section reveals a dorsal and ventral surface of the scale. The soft keratin of the ventral surface of the scale abuts the dorsal plate of hard keratin of the base of the claw. The dorsal plate curves ventrally over the bone of the distal phalanx.

Figure 12.107. Uropygial Gland, l.s., Chicken. A portion of one lobe of this bilobed holocrine gland shows branched tubular glands surrounded by a capsule of connective tissue. Each tubular gland is composed of a peripheral, sebaceous zone and an inner, glycogen zone. The latter communicates with the lumen of the lobe.



Figure 12.108 ×12.5




Figure 12.109 ×12.5




Figure 12.110 ×250




Figure 12.111 ×250

- 1. Adipose tissue
- 2. Basal cell, nucleus
- 3. Central layer
- 4. Epidermis
- 5. Feather follicle
- 6. Gland, lumen
- 7. Glycogen zone

- 8. Intermediate cell
- 9. Isthmus
- 10. Lobe, lumen
- 11. Primary duct
- 12. Sebaceous zone
- 13. Smooth muscle

Figure 12.108. Uropygial Gland, I.s., Chicken. The lumen of a lobe communicates with a primary duct, which passes through the isthmus toward the papilla of the gland.

Figure 12.109. Uropygial Gland, I.s., Chicken. The two primary ducts pass through the papilla (nipple). Their openings (not shown) onto the surface are surrounded by feathers.

Figure 12.110. Sebaceous Zone, Uropygial Gland, Chicken. Portions of tubular glands are shown in cross section. The basal layer of the glandular epithelium is represented by the oval to flat nuclei of the small basal cells. A single layer of acidophilic and grainy intermediate cells lies on the basal layer. Cells of the thick, central (transitional) layer accumulate lipid, hypertrophy, and degenerate toward the luminal surface.

Figure 12.111. Glycogen Zone, Uropygial Gland, Chicken. Portions of tubular glands are shown in cross section. Intermediate cells, with an acidophilic and grainy cytoplasm, form a thick layer in the glycogen zone. The pale cells of the central layer are less numerous. Compare with Figure 12.110.

DIGESTIVE SYSTEM

MAMMALS

he digestive tract extends from the mouth to the anus. Generally, its wall is composed of an outer serosa (or an adventitia), muscularis externa, submucosa, and inner mucosa. The mucosa consists of an inner epithelium, a middle lamina propria, and an outer muscularis mucosae. A muscularis mucosae is absent from the mouth, pharynx, portions of the esophagus, and the rumen. The mouth lacks a submucosa and muscularis externa.

From the lips through the nonglandular stomach, the epithelium of the mucosa is stratified squamous. Among other places the epithelium is keratinized on the dental pad, surface of the tongue, hard palate, cheek, and the nonglandular stomach of ruminants, horses, and pigs. The epithelium of the mucosa in the glandular stomach, the small intestine, and most of the large intestine is simple columnar; in the anal canal it is stratified squamous.

From the mouth through the esophagus, the mucosa is moistened by the secretions (mucous or serous) of various glands, including the major salivary glands. Surface mucous cells and mucous neck cells of the stomach, and goblet cells of the small and large intestines also contribute lubricating secretions.

Tongue

The tongue has various small outgrowths, papillae, located primarily on its upper surface. These vary considerably in size and appearance. Some (filiform) have threadlike projections or bear spines. Some are cushion-shaped (circumvallate, fungiform), while others (foliate) take the form of a succession of folds. Taste buds occur in the epithelium of circumvallate, foliate, and fungiform papillae.

Oropharynx

The **oropharynx** is lined by a stratified squamous epithelium and contains mucous glands, except in carnivores, where the glands are mixed. A muscularis externa of skeletal muscle is surrounded by an adventitia.

Esophagus

Throughout most of its length, the **esophagus** is surrounded externally by an adventitia. The muscularis externa varies in composition. In the dog it is composed of skeletal muscle throughout its length, except in the vicinity of the stomach, where skeletal muscle is replaced by smooth muscle. In ruminants the entire muscularis externa is comprised of skeletal muscle. In the horse and cat a switch from skeletal to smooth muscle occurs in the caudal third of the esophagus, whereas in the pig the change occurs just cranial to the diaphragm.

The mucosa of the esophagus is lined by stratified squamous epithelium. The longitudinally arranged smooth muscle of the esophageal muscularis mucosae varies in amount from anterior to posterior. It is in the form of isolated bundles anteriorly and a continuous sheet posteriorly in the cat, horse, and ruminant. In the dog and pig it is absent anteriorly and appears as a continuous sheet posteriorly.

Mucous or mixed glands occur in the submucosa of the esophagus. In the cat, horse, and ruminant, glands occur only at the junction of the pharynx and esophagus. In the pig, they occur anteriorly, diminish in the midregion, and are sparse caudally. In the dog, they occupy the entire length of the esophagus and extend into the stomach for a short distance.

Stomach

The horse, ruminant, and pig have a nonglandular fore-stomach and a glandular stomach. In ruminants, the forestomach is divisible into a rumen, reticulum, and omasum. The glandular stomach of ruminants is the abomasum. The cat and dog have a glandular stomach, but lack a forestomach. In all of these animals the glandular stomach consists of cardiac, fundic, and pyloric gland regions. The cardiac gland region is relatively small in all but the pig.

The epithelium of the glandular stomach invaginates into the lamina propria, forming tubular structures called **gastric pits** (foveolae). Depressions of the mucosa known as **gastric furrows** are also present.

Various tubular glands empty into the bottom of the gastric pits. Mucous glands with occasional parietal cells are the principal type found in the cardiac gland region. In the fundic gland region, glands are constructed mostly of parietal and chief cells, which secrete hydrochloric acid and pepsinogen, respectively. The glands of the pyloric gland region are mainly of the mucous type with interspersed parietal cells.

In carnivores, the mucosa of the fundic gland region is separated into an adoral, narrow, and thin **light zone**, and an aboral, wide, and thick **dark zone**. These zones are

readily visible on gross examination of the mucosa and are distinguishable histologically. The stomach of the cat has a thick layer of connective tissue between the base of the glands and the muscularis mucosae, called the **stratum compactum**. This may be capped by a layer of fibroblasts, the **stratum granulosum**. The combination of these cells and the stratum compactum is called the **lamina subglandularis**. The latter may be absent in dogs. A submucosa, muscularis externa of smooth muscle, and serosa complete the wall of the stomach.

Intestines

The intestines of mammals consist of a small intestine (duodenum, jejunum, and ileum) and a large intestine (cecum, colon, rectum, and anal canal). In both the small and large intestine, the epithelium is simple columnar with a striated border. Goblet cells occur among the columnar cells. The former increase in number from anterior to posterior, with the greatest number occurring in the large intestine.

Small Intestine

Villi are confined to the small intestine in mammals. They are short and thick in ruminants but long and slender in carnivores. At the bases of the villi are invaginations of the epithelium, the crypts of Lieberkühn (intestinal glands). Replacement of the mucosal epithelium occurs by cell division, primarily within the crypts. A muscularis mucosae, consisting of two layers of smooth muscle, separates the crypts from the underlying submucosa. The latter is formed from loose connective tissue in the horse, ruminant, and pig. In contrast, it is composed of moderately dense connective tissue in carnivores. A lamina subglandularis may be present in the intestine of carnivores. The remainder of the wall of the intestine is comprised of a muscularis externa of smooth muscle and a serosa.

Compound, tubuloacinar Brünner's glands (duodenal glands, submucosal glands) are mucous glands occurring within the submucosa and often within the lamina propria of the duodenum. In carnivores, sheep, and goats, they are limited to the initial or mid-region of the duodenum; in horses, pigs, and cows, they extend into the jejunum. Brünner's glands also project into the pyloric stomach for a short distance. Aggregations of lymphatic nodules, Peyer's patches, are present in the lamina propria and submucosa of the small intestine, especially the ileum.

Large Intestine

The mucosa of the large intestine presents a flat surface. Villi are absent. Crypts are longer than in the small intestine. Flat bands, taenia coli, consisting of longitudinally arranged smooth muscle and elastic fibers, occur in the colon of horses and pigs. Similar structures, taenia ceci, occur in the cecum. The rectum terminates at the anal canal, which is lined by a stratified squamous epithelium. The epithelium is nonkeratinized in the anterior portion of the canal and keratinized in the posterior portion, which is continuous with the hairy skin. Tubuloacinar anal glands

occur in the submucosa and muscularis of the anal canal in carnivores and pigs. Circumanal glands occur in the subcutis around the anus of the dog. The upper portion of these glands is sebaceous, whereas the lower portion is nonsebaceous. The cells of the latter resemble hepatocytes. Accordingly, the nonsebaceous region is often called a hepatoid gland.

Paired anal sacs occur lateral to and below the anus of carnivores. Each is lined by a keratinized, stratified squamous epithelium and is located between the inner smooth muscle of the internal anal sphincter and the outer skeletal muscle of the external anal sphincter. The excretory duct of each gland opens into the keratinized portion of the anal canal. Glands of the anal sac are apocrine tubular in the dog. In the cat, both apocrine tubular glands and sebaceous glands surround the anal sac.

Liver

The liver is a large, lobed gland. Each lobe is covered by a mesothelium, beneath which is a thin connective tissue layer, the capsule of Glisson. Each lobe is divided into numerous classic lobules. These consist of sinusoids and of plates of parenchyma cells, hepatocytes, radially organized about a central vein. Lobules are indistinctly separated from one another in all animals except the pig, in which an abundance of connective tissue between lobules clearly identifies their boundaries. Portal tracts (areas) occur at the interstices of three or more lobules. Each tract contains one or more branches of a portal vein, hepatic artery, bile ductule, and lymphatic vessel. These various components are supported by a framework of connective tissue.

Bile, secreted by hepatocytes, enters tiny bile canaliculi from which it flows into the canals of Hering, located close to each portal tract. The canals unite with the bile ductule of a portal tract. Bile ductules lead into bile ducts. The epithelium of bile ductules is simple cuboidal, whereas that of the bile ducts is simple columnar. Goblet cells occur in the largest bile ducts.

Gallbladder

The gallbladder is a storage depot for bile. Its mucosa is thrown into numerous folds when the bladder is contracted. When it is distended, they mostly disappear. The simple columnar epithelial lining has a striated border. Goblet cells have been reported in the epithelium of the cow. They have also been observed in the goat. Mucous, serous, or mixed glands are often seen in the wall of the gallbladder of ruminants. The smooth muscle of the muscularis is arranged circularly (mostly oblique, according to some) for the most part. The gallbladder is absent in the horse.

Pancreas

The pancreas consists of numerous tubuloacinar secretory units, which form the exocrine component of the organ. Clusters of epithelial cells, the endocrine islets of Langerhans

(pancreatic islets), are scattered among the secretory units. Tubuloacinar units drain into long, narrow intercalated ducts, which are lined by elongated cells that present a cuboidal appearance when sectioned transversely. Intercalated ducts communicate directly with interlobular ducts. Striated (secretory) ducts are not present. Unlike salivary glands, myoepithelial cells are lacking around the secretory units. Pacinian corpuscles are commonly found within the connective tissue of the pancreas of dogs and cats.

CHICKEN

In the chicken, the **salivary glands** are all of the mucous variety. They are located in the roof and floor of the oral cavity, tongue, and pharynx. **Taste buds** are present but sparse. They are associated with the ducts of salivary glands at the base of the tongue and the pharynx.

Esophagus

The esophagus has the usual seven layers. It is lined by a thick, nonkeratinized, stratified squamous epithelium. The muscularis externa is composed of smooth muscle along the entire length of the esophagus. Mucous glands occur in the lamina propria, but are lacking throughout most of the crop, which is a caudal diverticulum located approximately two-thirds of the way down the esophagus. The crop has a structure identical to the rest of the esophagus, but it lacks mucous glands.

Stomach

The stomach of the chicken consists of a glandular proventriculus and a muscular ventriculus (gizzard). The mucosa of the proventriculus is thrown into folds (plicae). Depressions between the folds are called sulci. The epithelium is simple columnar except at the base of the sulci, where it is cuboidal. The wall of the proventriculus consists of large, compound, tubular glands. The secretory cells, which are cuboidal to low columnar, produce both pepsinogen and hydrochloric acid, thus combining the function of mammalian chief and parietal cells. Each gland opens to the lumen of the stomach through a conical papilla.

The ventriculus is a highly muscular grinding organ. It is lined by an epithelium that invaginates into the lamina propria, forming elongated pits, each of which bears terminal tubular gastric glands. Cells of the latter secrete a thick, horny material. Although keratin-like, this substance, usually called keratinoid, is not chemically equivalent to keratin. It forms the tough inner lining, about 1 mm thick, of the ventriculus.

Intestine

The intestine of the chicken is similar in structure throughout its length. It consists of a duodenum, jejunum, ileum,

and large intestine. A pair of blind, elongated ceca join the intestine at the junction of the ileum and large intestine. The terminal end of the large intestine joins the coprodeum of the cloaca. Villi are present throughout the small and large intestines. They are longest in the duodenum, but gradually shorten and thicken caudally. In the coprodeum, they are stumpy and rounded. Villi also are present in the ceca, becoming flattened toward the blind end. Crypts of Lieberkühn (intestinal glands) are short and open between the villi, as in mammals. Although the wall of the intestine of the chicken is similar to that of the mammal, the absence of duodenal glands and an extremely thin submucosa in the chicken are notable differences.

Liver

As in mammals, the liver is covered by a mesothelium beneath which is a layer of connective tissue, Glisson's capsule. Lobes of the liver are subdivided into numerous lobules indistinctly separated from one another. The radiating plates of hepatocytes in each lobule are two cells wide

in the chicken. In contrast, those of mammals are one cell wide.

Gallbladder

The gallbladder of the chicken is similar to that of the mammal. The mucosa is lined by a simple columnar epithelium and is strongly folded into villuslike projections when contracted.

Pancreas

The pancreas of the chicken resembles that of the mammal. The exocrine portion is tubuloacinar. Lobulation is indistinct because of the lack of interlobular connective tissue. Islets of Langerhans are abundant. Two types of islets, alpha and beta, can be easily recognized. Columnar alpha cells characterize the alpha islet. Polygonal beta cells are the principal cells of the beta islets. Alpha islets produce glucagon, whereas beta islets form insulin.

| Word Roots | | |
|------------|-------------------------------|--|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Circum | Around | Circumanal glands occur in the subcutis around the anus of the dog. |
| Fili | A thread | Filiform papillae of the tongue have threadlike, keratinized projections. |
| Gast | Stomach | The epithelium of the glandular stomach invaginates, forming <i>gastric</i> pits. |
| Hepat | Liver | Lobules of the liver contain plates of hepatocytes. |
| Lamina | A thin plate, sheet, layer | The <i>lamina</i> subglandularis is a layer below the glands in the lamina propria. |
| Villi | Shaggy hairs | Numerous tiny fingerlike projections, the villi, project into the lumen of the small intestine, giving it a shaggy appearance. |




Figure 13.1 ×25




Figure 13.2 ×25




Figure 13.3 ×125




Figure 13.4 ×12.5




Figure 13.5 ×12.5

| KEY | | |
|---|---|--|
| Alveolar bone Connective tissue, papilla of Dental lamina Dental papilla Dental sac | 11. Mucous acinus12. Outer enamel epithelium13. Sebaceous gland14. Serous acinus15. Serous demilune | |
| 6. Duct7. Hair follicle8. Inner enamel epithelium9. Intralobular duct10. Labial gland | 16. Skeletal muscle17. Stellate reticulum18. Stratified squamous epithelium, keratinized19. Stratum granulosum | |

Figure 13.1. Lip, Sheep. The section was taken through the junction of the hairy and nonhairy portions of the lip. The stratum granulosum is present in the epidermis of the hairy portion of the lip, but disappears at the junction with the nonhairy portion of the lip. Portions of hair follicles are present.

Figure 13.2. Lip, Sheep. Oral surface of the lip with mixed labial glands among the skeletal muscle.

Figure 13.3. Lip, Sheep. Mixed labial glands within the skeletal musculature.

Figure 13.4. Cheek, Sheep. The mucous membrane of the cheek of ruminants is characterized by numerous, conical papillae. The apex and lateral surfaces of the papillae are highly keratinized.

Figure 13.5. Developing Permanent Tooth, Dog. The ectodermally derived enamel organ has differentiated into the outer and inner enamel epithelium and the stellate reticulum. The dental papilla, derived from mesenchyme, is in contact with the inner enamel epithelium.




Figure 13.6 ×12.5




Figure 13.7 ×125




Figure 13.8 ×62.5




Figure 13.9 ×25

| KEY | |
|--------------------------|---|
| 1. Alveolar bone | 12. Epithelium, free gingiva |
| 2. Ameloblasts | 13. Fiber bundle |
| 3. Attachment epithelium | 14. Lamina propria |
| 4. Blood vessel | 15. Odontoblasts |
| 5. Cementum | Outer enamel epithelium |
| 6. Dental papilla | 17. Precementum |
| 7. Dental pulp | 18. Predentin |
| 8. Dental sac | 19. Space artifact |
| 9. Dentin | 20. Stellate reticulum |
| 7. DOI | 20. Giolidio Tolicololli |

Figure 13.6. Developing Permanent Tooth, Dog. Dentin and enamel formation has begun (Figure 13.7).

21. Stratum intermedium

10. Enamel

11. Enamel space

Figure 13.7. Dentinoenamel Junction, Developing Permanent Tooth, Dog. Odontoblasts cover the surface of the mesenchymal dental papilla. These cells produce predentin (uncalcified dentin). The pale layer of predentin abuts the recently calcified dentin. The enamel organ consists of tall, columnar ameloblasts that produce enamel; a stratum intermedium; and the stellate reticulum.

Figure 13.8. Root of Tooth, x.s., and Periodontal Ligament, Dog. The periodontal ligament consists of bundles of collagenous fibers, blood and lymphatic vessels, nerves, and cells (mostly fibroblasts). The fiber bundles extend between, and anchor to, the cementum of the tooth and the alveolar bone. The ends of the fibers that are embedded in either cementum or bone are called Sharpey's fibers. They are indistinct in this micrograph.

Figure 13.9. Upper Deciduous Tooth, Decalcified, and Gingiva, I.s., Dog. The enamel space identifies the location of enamel before it was lost during decalcification. The attachment (junctional) epithelium of the gingiva is nonkeratinized stratified squamous and lacks papillae of connective tissue. It abuts the enamel region and is continuous with the papillated, keratinized stratified squamous epithelium of the free gingiva.




Figure 13.10 ×12.5




Figure 13.11 ×62.5




Figure 13.12 ×2




Figure 13.13 ×25




Figure 13.14 ×12.5

| | KEY |
|----------------------------------|------------------------------------|
| 1. Adipose tissue | 8. Palatine bone |
| 2. Connective tissue, papilla of | 9. Stratified squamous epithelium, |
| 3. Horn tubule-like structure | keratinized |
| 4. Lamina propria | 10. Stratum basale |
| 5. Loose connective tissue | 11. Stratum corneum |
| 6. Mixed gland | 12. Stratum spinosum |
| 7. Mucous gland | 13. Submucosa |

Figure 13.10. Dental Pad, Cow. The dental pad of ruminants is distinguished by its thick stratum corneum and well-developed papillae of connective tissue.

Figure 13.11. Epithelium, Dental Pad, Sheep. The section shows a thick stratum corneum and underlying stratum spinosum. A horn, a tubulelike structure, extends through the stratum corneum.

Figure 13.12. Hard Palate, Caudal, Dog (Masson's). All domestic mammals, except the pig, have glands (mucous or mixed) in the submucosa of the caudal portion of the hard palate. The cranial portion lacks glands in all of the domestic mammals.

Figure 13.13. Hard Palate, Pig. Large irregular patches of pale, loose connective tissue are scattered throughout the submucosa.

Figure 13.14. Soft Palate, Cow. Mucous glands and adipose tissue occupy portions of the lamina propria and submucosa. The epithelium is stratified squamous and keratinized.








Figure 13.16 ×25



Figure 13.17 ×25



Figure 13.18 ×250

- 1. Connective tissue, papilla of
- 2. Connective tissue, papilla of (caudal)
- 3. Connective tissue, papilla of (rostral)
- 4. Filiform papilla, projection of
- 5. Fungiform papilla, core
- 6. Sensory cell, nucleus
- 7. Skeletal muscle
- 8. Small papilla
- 9. Spine
- 10. Stratum spinosum
- 11. Supporting cell, nucleus
- 12. Taste pore

Figure 13.15. Filiform Papilla, Tongue, Cat. The filiform papilla of carnivores contains several small, rostral connective tissue papillae and a large, caudal connective tissue papilla. A large, keratinized spine is associated with the caudal papilla.

Figure 13.16. Filiform Papillae, Tongue, Horse. In horses and pigs, delicate, threadlike, keratinized projections extend from the upper surfaces of the filiform papillae. The papillae of connective tissue are long, but are not subdivided.

Figure 13.17. Fungiform and Filiform Papillae, Tongue, Goat. This section is from the tip of the tongue. The fungiform papilla is moundlike in section with a broad core of connective tissue containing numerous nerves. Portions of keratinized projections of filiform papillae appear on either side. In ruminants, the connective tissue of filiform papillae is subdivided into several small papillae.

Figure 13.18. Taste Bud, Fungiform Papilla, Tongue, Horse. The taste bud is embedded within the keratinized stratified squamous epithelium of a fungiform papilla. Supportive and sensory cells are visible within the bud. The nucleus and cytoplasm of the sensory cells are slightly darker than those of the supporting cells.




Figure 13.19 ×5

| KEY | |
|---|---|
| Connective tissue, core of Connective tissue, papilla of Duct | 4. Lingual salivary gland5. Stratified squamous epithelium6. Taste buds |

Figure 13.19. Circumvallate Papilla, Tongue, Goat. This large papilla lies within a depression of the lingual epithelium. Taste buds occur within the epithelium of the papilla facing the cavity of the depression, but are usually absent from the upper surface of the papilla.




Figure 13.20 ×25




Figure 13.21 ×25





Figure 13.22 \times 36




| Figure 13.23 | ×25 |
|---|---|
| l l | KEY |
| Adipose tissue Core of connective tissue Filiform papilla, epithelium of Interlobular connective tissue Interlobular duct Intralobular duct Lobule Mucous gland | 9. Muscle, longitudinal 10. Muscle, transverse 11. Muscle, vertical 12. Serous gland 13. Skeletal muscle 14. Stratified squamous epithelium, keratinized 15. Space artifact |
| | |


Figure 13.20. Conical Papilla, Tongue, Goat. This highly keratinized papilla is located on the upper surface of the tongue.

Figure 13.21. Lingual Salivary Glands, Horse. Both mucous and serous glands occur between the skeletal muscle bundles of the tongue.

Figure 13.22. Musculature, Tongue, I.s., Cat. The vertical, horizontal, and transverse arrangement of the lingual skeletal musculature can be seen below the mucosal papillae.

Figure 13.23. Parotid Gland, Horse. Portions of several lobules are shown. Lobules are often delineated by space artifacts. See Figure 13.24 for detail of a lobule.










Figure 13.26 ×250




Figure 13.27 ×125




Figure 13.28 ×250

| KEY | |
|--|--|
| Basal striations Intercalated duct Interlobular connective tissue Mucous acinus | 5. Secretory unit6. Serous acinus7. Serous demilune8. Striated duct |

Figure 13.24. Parotid Gland, Horse. Serous acini, intercalated ducts, and striated (secretory) ducts are present. The latter show clearly defined basal striations.

Figure 13.25. Parotid Gland, Dog. Serous acini and intralobular ducts are shown.

Figure 13.26. Parotid Gland, Cow. The secretory units are lined by pale, acidophilic cells with large nuclei. The cells vary in height, giving the luminal surface a scalloped appearance. This feature is unique to the cow.

Figure 13.27. Submandibular Gland, Dog. Mucous acini (some with serous demilunes) and serous acini characterize the parenchyma.

Figure 13.28. Submandibular Gland, Sheep. Intercalated ducts branching from a striated duct.




Figure 13.29 ×250

- 1. Intercalated duct
- 2. Interlobular connective tissue
- 3. Mucous acinus
- 4. Serous acinus
- 5. Serous demilune
- 6. Striated duct
- 7. Tubular mucous unit

Figure 13.29. Submandibular Gland, Sheep (Mallory's). The junction between a mucous acinus and an intercalated duct is illustrated.

Figure 13.30. Sublingual Gland, Dog. In the cat, dog, and horse, the sublingual gland contains mucous secretory units, serous acini, and serous demilunes. Long tubular mucous units are a characteristic feature of the gland in the dog.




Figure 13.30 ×130




Figure 13.31



12

Figure 13.33

| KEY | |
|--|--|
| Adipose tissue Blood vessel Coat of connective tissue Duct Epithelium Goblet cells Intercalated duct | 8. Interlobular connective tissue 9. Mixed glands 10. Mucous acinus 11. Serous demilune 12. Skeletal muscle 13. Stratified squamous epithelium 14. Striated duct |

Figure 13.31. Sublingual Gland, Pig. In the pig and ruminant, mucous tubuloacinar secretory units predominate. Serous demilunes are sparse.

Figure 13.32. Interlobular Duct, Sublingual Gland, Pig. This large interlobular duct is lined by a columnar epithelium. The latter is bistratified in places. Goblet cells occur in the epithelium.

Figure 13.33. Oropharynx, Dog. The section shows mixed glands among the skeletal muscle and within the submucosa. Mixed glands are shown in detail in Figure 11.10.




Figure 13.34 ×62.5

- 1. Blood vessel
- 2. Duct
- 3. Elastic fibers
- 4. Lamina propria
- 5. Mixed glands6. Mucous glands
- 7. Muscularis externa, inner circular
- 8. Muscularis externa, outer longitudinal
- 9. Muscularis mucosae
- 10. Stratified squamous epithelium
- 11. Submucosa

Figure 13.34. Oropharynx, Dog (Orcein). A thick band of connective tissue, containing numerous elastic fibers, parallels the mucosa.

Figure 13.35. Esophagus, Mid-region, x.s., Dog. The glands of the dog's esophagus are predominantly mucous. They are located in the submucosa throughout the length of the esophagus of the dog. The muscularis externa is skeletal muscle, except very near the stomach (Figure 13.36).



Figure 13.35 ×52





Figure 13.36



 $\times 25$ **Figure 13.37**




Figure 13.38 ×12.5




Figure 13.39 $\times 12.5$

11. Muscularis mucosae

12. Skeletal muscle 13. Smooth muscle

- 1. Adipose tissue
- 2. Adventitia
- 3. Duct
- 4. Esophagus, lumen
- 5. Lamina propria
- 7. Mucous glands
- 8. Muscularis externa, inner circular
- 6. Mixed glands
- 14. Stratified squamous epithelium

9. Muscularis externa, outer longitudinal

10. Muscularis externa, outer oblique

- 15. Stratified squamous epithelium, keratinized
- 16. Submucosa
- Figure 13.36. Esophagus, Near Stomach, I.s., Dog. The muscularis externa of the dog's esophagus is composed of skeletal muscle, except in the region caudal to the diaphragm, where the musculature is mixed.

Figure 13.37. Esophagus, Cranial, I.s., Cat. A few mixed glands are present in the submucosa. In the cat, horse, and ruminant, esophageal glands occur only near the junction of the esophagus and pharynx. The stratified squamous lining of the esophagus of carnivores is typically nonkeratinized. In the cat and hors, the muscularis externa is skeletal muscle throughout much of the esophagus. The transition from skeletal to smooth occurs in the caudal one-fifth to one-third of the esophagus in these animals.

Figure 13.38. Esophagus, Cranial, x.s., Horse. The papillated stratified squamous epithelium shows a distinct keratinized layer. The epithelium is also keratinized in pigs and ruminants. A sparse muscularis mucosae is present in the cranial esophagus in the horse, cat, and ruminant. The muscularis externa consists of skeletal muscle in this region.

Figure 13.39. Esophagus, Cranial, I.s., Pig. The stratified squamous epithelium is also keratinized in horses and ruminants. Note the abundance of mucous glands in the submucosa. In the cranial portion of the esophagus, a muscularis mucosae is absent in the pig and dog. The muscularis externa consists of skeletal muscle in this region.




Figure 13.40 ×12.5




Figure 13.41 ×12.5




Figure 13.42 ×62.5




Figure 13.43 ×62.5

- 1. Adipose tissue
- 2. Adventitia
- 3. Connective tissue, papilla of
- 4. Esophagus, lumen
- 5. Lamina propria
- 6. Mucous gland
- 7. Muscularis externa, inner circular
- 8. Muscularis externa, outer longitudinal
- Muscularis externa, outer oblique
- 10. Muscularis mucosae
- 11. Serosa
- 12. Skeletal muscle
- 13. Smooth muscle
- 14. Stratified squamous epithelium
- 15. Submucosa

Figure 13.40. Esophagus, Mid-region, I.s., Pig. Note the decrease in glandular tissue and the presence of a muscularis mucosae. Compare with Figure 13.39. The muscularis externa consists of skeletal muscle in this region.

Figure 13.41. Esophagus, Caudal, x.s., Pig. The region of the pig's esophagus just cranial to the diaphragm shows the presence of smooth and skeletal muscle in the muscularis externa, a lack of glands, and a thick muscularis mucosae.

Figure 13.42. Esophagus, Caudal, x.s., Pig. Detail of Figure 13.41 shows the smooth and skeletal musculature of the muscularis externa.

Figure 13.43. Esophagus, Mid-Region, x.s., Sheep. The muscularis mucosae is less developed than in the pig (see Figure 13.40).

Summary of typical layers of the wall of organs of the digestive tract

A. Mucosa

- 1. Epithelium: Simple columnar or stratified squamous
- 2. Lamina propria: Loose connective tissue
- 3. Muscularis mucosae: Smooth muscle tissue
- B. **Submucosa**: Loose or dense connective tissue
- C. Muscularis externa: Skeletal and/or smooth muscle tissue; varies with the organ, part of the organ, and type of animal
- D. Serosa: An epithelium (called a mesothelium) and underlying connective tissue, OR Adventitia: A layer of connective tissue that lacks an epithelium and blends with surrounding connective tissue




Figure 13.44



×25 **Figure 13.45**



Figure 13.46 ×12.5




Figure 13.47 ×125




Figure 13.48

| KEY | |
|-------------------------------------|--|
| 1. Arteriole | 6. Lymphatic vessel |
| 2. Capillary | 7. Muscularis externa |
| 3. Connective tissue, | 8. Stomach contents |
| band of | 9. Stratified squamous epithelium, keratinized |
| Keratinized cells | 10. Submucosa |
| 5. Lamina propria | |


Figure 13.44. Rumen, Sheep. The mucosa of the rumen is differentiated into paddle-shaped papillae.

Figure 13.45. Rumen, Cow. A part of the wall from the lumen to the beginning of the muscularis externa (smooth muscle). A complete short papilla and a portion of a long papilla are included. Note the submucosa entering the long papilla. A muscularis mucosae is not present in this part of the forestomach.

Figure 13.46. Rumen, Sheep. Section shows two cuts through adjacent long papillae. These papillae are flat, paddlelike, structures (Figure 13.44). The bottom one in the photograph was cut parallel to the flat surface, and the top one was cut perpendicular to the flat surface. A dense, more darkly stained band of connective tissue mimics a muscularis mucosae.

Figure 13.47. Rumen, Sheep. Section illustrates the vacuolated, keratinized cells of the stratified squamous epithelium of a papilla. Numerous capillaries abut the epithelium.

Figure 13.48. Reticulum, Sheep. The mucosa of the reticulum is extended into intersecting folds that subdivide the surface into distinct compartments, suggesting a honeycomb. Conical papillae project from the crests of the folds and from the mucosa of the compartments.










Figure 13.51 ×25




Figure 13.52




Figure 13.53 ×25

| | KEY |
|--|------------------------------------|
| 1. Conical papilla | 5. Muscularis mucosae |
| 2. Connective tissue, band of | 6. Stratified squamous epithelium, |
| 3. Lamina propria 4. Muscularis externa | keratinized |
| 4. Muscularis externa | 7. Submucosa |

Figure 13.49. Reticulum, Cow. Short folds and the base of a long fold are visible. A muscularis mucosae occurs in the upper segment of the long fold. This is a characteristic feature of the reticulum. Sides and crests of long folds have conical papillae with keratinized tips.

Figure 13.50. Reticulum, Sheep. Section shows a long fold cut in a plane parallel to its flat surface. The apparent gaps are the result of undulations in the fold. Conical papillae are evident along the crest of the fold.

Figure 13.51. Reticulum, Goat (Masson's). The section is through the region of intersection of three long folds. The muscularis mucosae passes from one fold to another at the intersection. Two conical papillae, with keratinized tips, project from the sides of two of the folds.

Figure 13.52. Omasum, Sheep. Laminae (folds) of different sizes extend from the wall of the omasum, somewhat like the pages of a book. The mucosal surfaces of the laminae are studded with numerous conical papillae.

Figure 13.53. Omasum, Sheep. Small folds such as the one shown contain a lamina propria and muscularis mucosae, but lack an extension of smooth muscle from the muscularis externa.




Figure 13.54 ×62.5




Figure 13.55 ×25




Figure 13.56 ×125





Figure 13.57 ×12.5



igure 13.58 ×62.5

| rigule 13.30 | ×02.3 |
|---|--|
| | KEY |
| Cardiac gland region Cardiac glands Columnar epithelium, stomach Elastic fiber Fundic gland region Lamina propria Lymphatic vessel Mixed glands | 9. Muscularis mucosae 10. Papilla 11. Smooth muscle of muscularis externa 12. Stratified squamous epithelium, esophagus 13. Stratified squamous epithelium, keratinized 14. Vein |

Figure 13.54. Omasum, Sheep. The base of a long fold is shown. In addition to the muscularis mucosae, smooth muscle from the muscularis externa projects into the center of the fold.

Figure 13.55. Omasum, Sheep. Portions of two long folds are shown. Numerous small papillae cover the surface of the folds.

Figure 13.56. Omasum, Goat (Orcein). The lamina propria of a portion of a small papilla contains an extensive network of elastic fibers.

Figure 13.57. Junction, Esophagus and Cardiac Gland Region of Stomach, Dog. Numerous glands, predominantly mucous with a few serous demilunes, occupy the submucosa of the esophagus and extend into the cardiac gland region of the stomach of dogs.

Figure 13.58. Junction, Esophagus and Cardiac Gland Region of Stomach, Dog. Detail of Figure 13.57. The stratified squamous epithelium of the esophagus ends abruptly where the columnar epithelium of the stomach begins.








Figure 13.60 ×25




Figure 13.61 ×25




Figure 13.62 \times 62.5




Figure 13.63 ×250

| | KEY |
|----------------------------------|-------------------------------------|
| 1. Cardiac glands | 9. Lamina propria |
| 2. Cardiac sphincter | 10. Lymphatic nodule |
| 3. Chief cell | 11. Muscularis mucosae |
| 4. Connective tissue, papilla of | 12. Parietal cell |
| 5. Esophageal glands | 13. Simple columnar epithelium |
| 6. Fundic gland region | 14. Stratified squamous epithelium, |
| 7. Fundic glands | keratinized |
| 8. Gastric pit | 15. Submucosa |

Figure 13.59. Junction of Esophagus and Stomach, Dog. Section passes through the cardiac sphincter.

Figure 13.60. Margo Plicatus, Horse. At the margo plicatus, the keratinized stratified squamous epithelium of the forestomach ends, and the simple columnar epithelium of the cardiac gland region of the stomach begins.

Figure 13.61. Cardiac Gland Region, Stomach, Pig. Numerous lymphatic nodules characterize the mucosa of the cardiac gland region of the pig's stomach.

Figure 13.62. Light Zone, Fundic Gland Region, Stomach, Dog. The light zone of the carnivore's fundic stomach has a thinner mucosa than the more aboral dark zone. The gastric pits of the light zone are comparatively deep, extending, in some cases, to a depth equivalent to half the thickness of the mucosa. Compare with Figure 13.65. Note that smooth muscle (arrows) of the muscularis mucosae of the stomach extends into the lamina propria.

Figure 13.63. Light Zone, Fundic Gland Region, Stomach, Dog. Chief and parietal cells form the walls of the fundic glands. The glands are shown in cross section.




Figure 13.64 ×12.5




Figure 13.65 ×62.5




Figure 13.66 ×62.5




Figure 13.67 ×62.5




Figure 13.68 ×250

| | KEY | |
|--|--|--|
| 1. Chief cells 2. Fundic gland, lumen 3. Fundic glands 4. Gastric pit 5. Lamina propria 6. Mucosa 7. Mucous neck cells 8. Muscularis externa | 9. Muscularis mucosae 10. Parietal cells 11. Serosa 12. Smooth muscle 13. Stratum compactum 14. Stratum granulosum 15. Submucosa 16. Surface mucous cells | |

Figure 13.64. Dark Zone, Fundic Gland Region, Stomach, Dog. A portion of the stomach wall, including the base of a fold, is shown.

Figure 13.65. Dark Zone, Fundic Gland Region, Stomach, Dog. The mucosa of the dark zone of the fundic stomach of carnivores is thicker than that of the light zone. The gastric pits are comparatively shallow, extending no farther into the mucosa than one-third of its thickness. Compare with Figure 13.62.

Figure 13.66. Dark Zone, Fundic Gland Region, Stomach, Dog (PAS). The surface mucous cells lining the lumen, cells of the gastric pits and the mucous neck cells of the glands, both contain complex carbohydrates and are PAS positive (magenta stain).

Figure 13.67. Lamina Subglandularis, Fundic Gland Region, Stomach, Cat (Old). A thick layer of connective tissue, the stratum compactum, and an overlying layer of fibroblasts, the stratum granulosum, together form the lamina subglandularis, a structure seen consistently in cats and occasionally in dogs. Presumably, the lamina subglandularis protects the stomach from punctures by sharp objects.

Figure 13.68. Fundic Gland Region, Stomach, Cat. The fundic glands have been cut longitudinally. They are formed largely from parietal and chief cells.




Figure 13.69 ×12.5




Figure 13.70 ×62.5




Figure 13.71 ×125




Figure 13.72 ×25




Figure 13.73 ×62.5

| KEY | |
|----------------------------------|--------------------------------|
| 1. Adipose tissue | 8. Mucosa |
| 2. Cardiac glands | 9. Mucous precursor |
| 3. Chief cells | 10. Muscularis externa |
| 4. Epithelial cells, surface cut | 11. Muscularis mucosae |
| 5. Fundic glands | 12. Parietal cells |
| 6. Gastric pit | 13. Simple columnar epithelium |
| 7. Lamina propria | 14. Submucosa |

Figure 13.69. Fundic Gland Region, Stomach, Horse. Note the thick mucosa and submucosa.

Figure 13.70. Junction of the Cardiac and Fundic Gland Regions, Stomach, Horse. The mucous glands of the cardiac gland region are distinct from the parietal and chief cells of the fundic gland region of the stomach.

Figure 13.71. Fundic Gland Region, Stomach, Horse. The presence of surface mucous cells is a unique feature of the epithelium of the glandular stomach. Mucigen, the precursor of mucus, fills much of the apical portion of these cells in living tissue.

Figure 13.72. Fundic Gland Region, Abomasum, Sheep. The submucosa shows extensive infiltration by adipose tissue.

Figure 13.73. Fundic Gland Region, Abomasum, Goat. Parietal and chief cells of the glands are evident. Note the deep gastric pits.




Figure 13.74 ×250




Figure 13.75 ×62.5




Figure 13.76 ×1





Figure 13.77 ×25



| Figure 13.78 | ×62.5 |
|--|---|
| | KEY |
| 1. Chief cell 2. Gastric pit 3. Globular leukocyte 4. Lamina propria 5. Lamina subglandularis 6. Lymphocytes | 10. Parietal cell11. Pyloric gland12. Serosa13. Simple columnar epithelium14. Simple columnar epithelium, oblique section |
| 7. Mucous precursor 8. Muscularis externa 9. Muscularis mucosae | 15. Submucosa 16. Surface mucous cells |

Figure 13.74. Fundic Gland Region, Abomasum, Cow. Parietal and chief cells of the fundic glands.

Figure 13.75. Mucosa, Pyloric Gland Region, Stomach, Dog (PAS). The content of the surface mucous cells and that of the secretory units of the pyloric glands are PAS positive (magenta color).

Figure 13.76. Surface Mucous Cells, Pyloric Gland Region, Stomach, Dog. Columnar cells lining the gastric pits and bordering the gastric lumen show typical cup-shaped concentrations of mucous precursor (mucigen) in their apical ends. The epithelium contains many migrating lymphocytes.

Figure 13.77. Pyloric Gland Region, Stomach, Cat. Note the deep gastric pits. Some extend halfway through the thickness of the mucosa.

Figure 13.78. Mucosa, Pyloric Gland Region, Stomach, Cat. Note the presence of deep gastric pits. Some extend to about half the depth of the mucosa.




Figure 13.79 ×250




Figure 13.80 ×2




Figure 13.81




Figure 13.82 ×3

- ← Junction, pyloric gland region and duodenum
- 1. Arteriole
- 2. Brünner's gland
- 3. Crypt of Lieberkühn
- 4. Gastric furrow
- 5. Gastric pit
- 6. Lamina propria
- 7. Muscularis externa
- 8. Muscularis mucosae
- 9. Pyloric gland
- 10. Stratum compactum
- 11. Stratum granulosum
- 12. Submucosa
- 13. Villus of duodenum

Figure 13.79. Pyloric Gland Region, Stomach, Cat. Basal ends of pyloric glands and the stratum granulosum and stratum compactum of the lamina subglandularis are present in this section.

Figure 13.80. Pyloric Gland Region, Abomasum, Goat. Gastric furrows and gastric pits can be seen.

Figure 13.81. Junction, Pyloric Gland Region and Duodenum, I.s., Dog. Brünner's glands are located primarily in the submucosa of the duodenum. They also extend a short distance into the pyloric gland region of the stomach. They occasionally break through the muscularis mucosae and extend into the lamina propria.

Figure 13.82. Junction, Pyloric Gland Region and Duodenum, I.s., Dog. Gastric pits and mucous glands of the pyloric gland region of the stomach can be seen. Brünner's glands (mucous) occur below the interrupted muscularis mucosae. See Figure 13.83 for detail of the epithelium.




Figure 13.83 ×26

| KEY | |
|--------------------------------|----------------------------------|
| 1. Central lacteal | 5. Lymphocytes |
| 2. Goblet cell | 6. Striated border |
| 3. Intestinal absorptive cells | 7. Surface mucous cells, stomach |
| 1 Igming proprig | |

Figure 13.83. Junction, Pyloric Gland Region and Duodenum, Dog. Note the change in the epithelium when it passes from the stomach to the duodenum. Typical columnar surface mucous cells of the pyloric gland region of the stomach contrast with the columnar absorptive cells and goblet cells of the duodenum.




Figure 13.84 ×125




Figure 13.85 ×25




Figure 13.86 ×62.5




Figure 13.87 ×125

- 1. Artifact
- 2. Brünner's gland
- 3. Crypt of Lieberkühn
- 4. Duct
- 5. Goblet cell
- 6. Muscularis externa
- 7. Muscularis mucosae
- 8. Secretory unit
- 9. Serosa
- 10. Submucosa
- 11. Villus

Figure 13.84. Brünner's Gland, Duodenum, I.s., Dog. Detail of mucous secretory units. The latter, in the dog, are lined by tall columnar cells and have large lumens. Compare with Figure 13.90.

KEY

Figure 13.85. Duodenum, x.s., Cat. A segment of the wall of the duodenum is shown. The intestinal villi of carnivores tend to be longer than those of noncarnivores. Note the shrinkage artifact at the apical ends of the villi.

Figure 13.86. Duodenum, Proximal, x.s., Cat (Masson's). Ducts of Brünner's glands penetrate the muscularis mucosae. Brünner's glands are marked off into distinct lobules in the cat.

Figure 13.87. Duodenum, x.s., Cat (Masson's). The submucosa in the cat and dog is a moderately dense irregular connective tissue. In other domestic mammals, it is a loose connective tissue.

Synonyms

Brünner's gland = duodenal gland; submucosal gland Crypt of Lieberkühn = intestinal gland; intestinal crypt




Figure 13.88 ×250



Figure 13.89 ×25




Figure 13.90 ×125




Figure 13.91 ×25




Figure 13.92 ×125

| KEY | | |
|------------------------|-----------------------|--|
| 1. Adipose cell | 8. Mucous gland | |
| 2. Adipose tissue | 9. Muscularis externa | |
| 3. Crypt of Lieberkühn | 10. Paneth cell | |
| 4. Globular leukocyte | 11. Secretory unit | |
| 5. Lamina propria | 12. Serous gland | |
| 6. Lymphocyte | 13. Striated border | |
| 7. Mucosa | 14. Submucosa | |

Figure 13.88. Epithelium of Villus, Duodenum, Cat. Lymphocytes can be seen migrating through the simple columnar epithelium. Note the presence of several globular leukocytes.

Figure 13.89. Duodenum, Horse. Paneth cells are visible in the basal portions of the crypts of Lieberkühn in the small intestine of the horse.

Figure 13.90. Duodenum, Horse. In the horse, Brünner's glands have both mucous and serous components. Note that the lumens of the secretory units are small. Compare with Figure 13.84.

Figure 13.91. Duodenum, Pig. Brünner's glands fill the entire submucosa.

Figure 13.92. Duodenum, Pig. In the pig the lumens of the secretory units of Brünner's glands are very small.




Figure 13.93 ×12.5




Figure 13.94 ×125




Figure 13.95 ×125




Figure 13.96 ×125




Figure 13.97 ×125

KEY

- 1. Brünner's gland
- 2. Crypt of Lieberkühn
- 3. Mucosa
- 4. Muscularis externa, inner circular
- 5. Muscularis externa, outer longitudinal
- 6. Muscularis mucosae
- 7. Nerve fascicle, unmyelinated
- 8. Secretory unit
- 9. Serosa
- 10. Submucosa

Figure 13.93. Duodenum, I.s., Cow. Brünner's glands are present throughout much of the submucosa of an intestinal fold (plica).

Figure 13.94. Duodenum, Cow. Detail of Brünner's gland. In the cow, some gland cells have an acidophilic cytoplasm. The lumens of secretory units are large.

Figure 13.95. Duodenum, x.s., **Sheep.** The muscularis externa of the intestine is arranged into an inner circular and outer longitudinal layer of smooth muscle. Compare the appearance of the muscle layers seen in this cross section with that of the longitudinal section of the intestine in Figure 13.96.

Figure 13.96. Duodenum, I.s., Sheep. This section is through the muscularis externa. Compare the appearance of the muscle layers in this preparation with that in Figure 13.95.

Figure 13.97. Duodenum, Goat. Portions of the mucosa and submucosa. The lumens of the secretory units of Brünner's glands are large in the goat.




Figure 13.98 ×62.5




Figure 13.99 ×12



Figure 13.100 ×25




Figure 13.101 ×125




Figure 13.102 ×25

| 119010 10:102 | ×230 |
|--|---|
| | KEY |
| 1. Auerbach's plexus 2. Central lacteal, x.s. 3. Columnar epithelium 4. Crypt of Lieberkühn 5. Eosinophil 6. Globular leukocyte 7. Goblet cell 8. Intestinal lumen 9. Lamina propria | 10. Meissner's plexus 11. Muscularis externa, inner circular 12. Muscularis externa, outer longitudinal 13. Muscularis mucosae 14. Nuclei of epithelium cut obliquely 15. Stratum compactum 16. Submucosa 17. Villus |

Figure 13.98. Jejunum, x.s., Dog. A Meissner's plexus is present in the periphery of the submucosa. An Auerbach's plexus is wedged between the inner circular and outer longitudinal layers of the muscularis externa.

Figure 13.99. Jejunum, Dog. A well-developed stratum compactum is present between the lamina propria and muscularis mucosae of the small intestine in some cats and dogs.

Figure 13.100. Jejunum, x.s., Cat. Slender villi and well-defined crypts of Lieberkühn are evident.

Figure 13.101. Jejunum, Cat. Transverse sections of villi. Central lacteals are evident in two of them. Migrating lymphocytes are visible within the epithelium.

Figure 13.102. Jejunum, Cat. Globular leukocytes are present among the columnar cells lining a crypt of Lieberkühn. Numerous eosinophils are scattered through the lamina propria.

Synonyms

Auerbach's plexus = myenteric plexus Meissner's plexus = submucosal plexus




Figure 13.103 ×12.5




Figure 13.104 ×25




Figure 13.105 ×250



Figure 13.106 ×12.5




Figure 13.107 ×62.5

| | KEY |
|------------------------|---|
| 1. Crypt of Lieberkühn | 8. Muscularis externa, inner circular |
| 2. Eosinophil | 9. Muscularis externa, outer longitudinal |
| 3. Goblet cell | 10. Muscularis mucosae |
| 4. Lamina propria | 11. Serosa |
| 5. Lymphatic nodule | 12. Submucosa |
| 6. Mitotic figure | 13. Villus |
| 7. Mucosa | |

Figure 13.103. Jejunum, I.s., Horse. All layers of the wall are included in this section. The villi are shorter than those of carnivores.

Figure 13.104. Ileum, x.s., Cat. A portion of the wall from the lumen to the serosa is shown. The epithelium of the villi contain numerous goblet cells.

Figure 13.105. Ileum, Dog. Mitotic figures can be seen in the crypts.

Figure 13.106. Cecum, Tip, Dog. Large lymphatic nodules are present in the submucosa.

Figure 13.107. Cecum, Dog. Numerous goblet cells in the lining of the crypts of Lieberkühn are characteristic of the organ. The epithelial cells bordering the lumen in this preparation have undergone some autolysis and look tattered.




Figure 13.108 ×250




Figure 13.109 ×12.5




Figure 13.110 ×62.5




Figure 13.111 ×25




Figure 13.112 ×12.5

| | KEY |
|---------------------------------------|---------------------------------------|
| 1. Adipose tissue | 9. Muscularis externa, inner circular |
| 2. Cecal band | 10. Muscularis mucosae |
| Crypt of Lieberkühn | 11. Serosa |
| 4. Goblet cell | 12. Small artery |
| 5. Lamina propria | 13. Striated border |
| 6. Lymphatic nodule | 14. Submucosa |
| 7. Mucosa | 15. Taenia coli |
| 8. Muscle band | 16. Vein |

Figure 13.108. Cecum, Dog. Several crypts of Lieberkühn appear in cross section. A striated border is present on the columnar cells. Goblet cells are numerous.

Figure 13.109. Cecal Band, x.s., Horse. A cecal band consists of an admixture of smooth muscle (thickened outer longitudinal layer of the muscularis externa) and elastic fibers. Elastic fibers are predominant in the bands of both the cecum and the ventral large colon of the horse. Compare with Figure 13.112.

Figure 13.110. Cecum, Horse. Veins with bands of smooth muscle in their walls are common in the submucosa throughout the digestive tract of the horse. Similar vessels are shown at low magnification in Figure 13.109.

Figure 13.111. Cecum, Cow. In the large intestine of ruminants, the crypts of Lieberkühn are usually tortuous. Adipose tissue is abundant in the submucosa.

Figure 13.112. Taenia Coli, Small Colon, x.s., Horse. The taenia coli are thickenings of the outer longitudinal layer of the muscularis externa. In contrast to the bands of the cecum and ventral large colon, where elastic fibers are predominant, smooth muscle predominates in the bands of the small colon and dorsal large colon. Compare with Figure 13.109.




Figure 13.113 ×12.5




Figure 13.114 ×25




Figure 13.115 ×250




Figure 13.116 ×62.5




Figure 13.117

| KEY | |
|----------------------------|---|
| 1. Adipose tissue | 11. Lamina propria |
| 2. Anal canal | 12. Mucosa |
| 3. Anal glands | 13. Muscularis externa, inner circular |
| 4. Anal sac | 14. Muscularis externa, outer longitudina |
| 5. Circumanal glands | 15. Muscularis mucosae |
| 6. Crypt of Lieberkühn | 16. Rectum |
| 7. Epidermis | 17. Striated border |
| 8. External anal sphincter | 18. Submucosa |
| 9. Glands of anal sac | 19. Taenia coli |
| 10. Goblet cell | |


Figure 13.113. Taenia Coli, Colon, x.s., Pig. The muscular taenia coli is formed from the outer longitudinal layer of the muscularis externa. The submucosa is infiltrated with fat.

Figure 13.114. Spiral Colon. x.s., Goat. The mucosa contains both tortuous and straight crypts of Lieberkühn. The muscularis externa shows an abrupt thickening of its inner circular layer and outer longitudinal layer.

Figure 13.115. Spiral Colon, x.s., Goat. The columnar epithelial cells have a distinct striated border.

Figure 13.116. Rectum, x.s., Cat. The epithelium of the rectum presents a flat, uniform surface.

Figure 13.117. Rectoanal Junction, l.s., Dog. Note that the anal glands mark the junction (arrow) of the rectum and anal canal.








Figure 13.119 ×125




Figure 13.120 ×62.5



Figure 13.121 ×12.5




Figure 13.122 ×62.5

- 1. Anal gland
- 2. Apocrine tubular gland
- 3. Circumanal gland
- 4. Crypt of Lieberkühn
- 5. Débris in anal sac
- 6. External anal sphincter
- 7. Keratinized epithelium of anal sac
- 8. Mucosa
- 9. Muscularis externa, inner circular
- 10. Sebaceous gland
- 11. Stratified squamous epithelium

Figure 13.118. Rectoanal Junction, I.s., Dog. Note the change between the stratified squamous epithelium of the anal canal and the crypts of Lieberkühn of the rectal mucosa. Note also that anal glands are located in the submucosa and are scattered among the smooth muscle of the internal anal sphincter (inner circular layer of the muscularis externa). See Figure 13.117 for orientation.

Figure 13.119. Anal Glands, Dog. Section is through the secretory units of an anal gland.

Figure 13.120. Glands of the Anal Sac, Dog. A small portion of the wall of an anal sac and the secretory units of some of the glands of the anal sac are shown. See Figure 13.117 for location.

Figure 13.121. Anal Sac, Cat. About one-half of the wall of an anal sac is shown.

Figure 13.122. Glands of the Anal Sac, Cat. Portions of these glands are shown adjacent to the skeletal muscle of the external anal sphincter. The presence of sebaceous glands in this location is a characteristic of the cat.




Figure 13.123 ×125




Figure 13.124 $\times 25$




Figure 13.125



Figure 13.126




Figure 13.127

| KEY | |
|--------|---|
| 200118 | 7 |

- 1. Circumanal gland, nonsebaceous
- 2. Circumanal gland, sebaceous
- 3. Crypt of Lieberkühn
- 4. Goblet cell
- 5. Hair follicle
- 6. Keratinized epidermis
- 7. Lamina propria
- 8. Sebaceous gland
- 9. Simple columnar epithelium
- 10. Skeletal muscle
- 11. Stratified squamous epithelium
- 12. Stratum granulosum

Figure 13.123. Anal Canal, I.s., Dog. Junction of the keratinized and nonkeratinized regions of the anal canal. The stratum granulosum of the keratinized region stops abruptly at the junction.

Figure 13.124. Circumanal Glands, Dog. Numerous nonsebaceous portions of circumanal glands are present subcutaneously among the skeletal muscle of the anal sphincter. These nonsebaceous portions are often called hepatoid glands because their cells resemble hepatocytes.

Figure 13.125. Circumanal Gland, Dog. Detail of a part of one of the glands. The lower, nonsebaceous portion of these glands is more acidophilic than the upper, sebaceous portion.

Figure 13.126. Rectoanal Junction, I.s., Horse. The stratified squamous epithelium of the anal canal contrasts with the rectal mucosa.

Figure 13.127. Rectoanal Junction, Horse. The abrupt change between the stratified squamous epithelium of the anal canal and the simple columnar epithelium of the rectal mucosa is apparent.




Figure 13.129




Figure 13.130 ×125




Figure 13.131 $\times 62.5$




Figure 13.132

KEY

- 1. Bile canaliculus, I.s.
- 2. Bile canaliculus, x.s.
- 3. Bile ductule
- 4. Central vein, l.s.
- 5. Central vein, x.s.
- 6. Hepatic artery, branch
- 7. Hepatocyte

- 8. Lobule, portion of
- 9. Lymphatic vessel
- 10. Nucleus of hepatocyte
- 11. Portal vein, branch
- 12. Serosa
- 13. Sinusoid


Figure 13.128. Liver, Cat. Transverse and longitudinal sections through the central veins of two classic lobules.

Figure 13.129. Liver, Cat. Transverse section through a classic lobule. Sinusoids empty into the central vein. Hepatocytes radiate as hepatic plates from the central vein.

Figure 13.130. Liver, Cat. The portal tract in this section includes a branch of the hepatic portal vein and hepatic artery, bile ductule, and lymphatic vessel.

Figure 13.131. Liver, Horse. A portal tract is at the intersection of three classic lobules. Branches of the portal vein can be seen extending between the lobules.

Figure 13.132. Liver, Cat (Masson's). Sections through various bile canaliculi are evident in this section.








Figure 13.134 ×250




Figure 13.135 ×25




Figure 13.136 ×250




Figure 13.137 ×250

| KEY | | |
|---|---|--|
| Bile duct Binucleate hepatocyte Capsule of Glisson Central vein Connective tissue, partition of Goblet cell | 7. Hepatocyte 8. Kupffer cell 9. Mesothelium 10. Portal tract 11. Portal vein, branch 12. Sinusoid | |
| | | |

Figure 13.133. Liver, Horse. A sinusoid joins with a branch of the portal vein.

Figure 13.134. Liver, Horse. Section is through mesothelial cells and connective tissue of the capsule of Glisson surrounding the liver. Together, the mesothelial cells and the capsule of Glisson comprise the serosa.

Figure 13.135. Liver, Pig. Classic lobules are clearly separated from one another by partitions of connective tissue in the pig.

Figure 13.136. Liver, Pig. Particulate-laden Kupffer cells within sinusoids of a classic lobule are evident in this section. Binucleate hepatocytes also can be seen.

Figure 13.137. Liver, Goat. A large bile duct with columnar epithelium and goblet cells.








Figure 13.139 ×25





Figure 13.141 ×250



3010 101142

| KEY |
|-----|
| |

- 1. Adventitia
- 2. Dark cell
- 3. Epithelium
- 4. Gallbladder, lumen
- 5. Goblet cell
- 6. Lamina propria
- 7. Light cell

- 8. Liver
 - 9. Mucosa
 - 10. Mucous acinus
 - 11. Muscularis
 - 12. Serous acinus
 - 13. Striated border

Figure 13.138. Gallbladder, Dog. A portion of the wall showing the highly folded mucosa.

Figure 13.139. Gallbladder, Dog. Epithelial lining with light and dark columnar cells.

Figure 13.140. Gallbladder and Liver, Pig (Masson's). Section shows a portion of the liver and gallbladder.

Figure 13.141. Gallbladder, Goat. Portion of a mucosal fold showing goblet cells in the epithelium.

Figure 13.142. Gallbladder, Goat. Mixed glands occur within the wall of the gallbladder of ruminants.




Figure 13.143 ×125




Figure 13.144




Figure 13.145 ×125




Figure 13.146 ×250




Figure 13.147 ×250

| KEY | | |
|---------------------------------|-----------------------------|--|
| 1. A cells | 9. Dark cell | |
| 2. Acinar cells | 10. Goblet cell | |
| 3. Acinus | 11. Intercalated duct, l.s. | |
| 4. Anastomotic artery | 12. Intercalated duct, x.s. | |
| 5. B cells | 13. Interlobular duct | |
| 6. Capillary | 14. Islet of Langerhans | |
| 7. Centroacinar cell | 15. Light cell | |
| 8. Connective tissue, septum of | 16. Muscularis | |

Figure 13.143. Cystic Duct, Pig. The epithelium is composed of light cells, dark cells, and goblet cells.

Figure 13.144. Pancreas, Horse. The organ is divided into lobules by septa of connective tissue. Most of the pancreas is formed from exocrine acinar cells. Islets of Langerhans are scattered through the exocrine region of the gland.

Figure 13.145. Pancreas, Horse. Portions of two lobules showing acinar cells, interlobular ducts, and intercalated (intralobular) ducts.

Figure 13.146. Pancreas, Horse. An islet of Langerhans with some surrounding exocrine acini. In the horse, the darker A cells are located in the center of the islet, while the lighter B cells are positioned in the periphery. Compare with Figure 13.150. Note the numerous capillaries among cords of islet cells.

Figure 13.147. Pancreas, Dog. Detail of acini and intercalated ducts. Note acidophilic apical regions and basophilic basal regions of the acinar cells.

Synonyms

A cell = alpha cell
B cell = beta cell
Islet of Langerhans = pancreatic islet




Figure 13.148 ×125




Figure 13.149 ×250




Figure 13.150 ×250




Figure 13.151 ×12.5




Figure 13.152 ×62.5

| KEY | | |
|--|---|--|
| A cell Acinus B cells Centroacinar cell Duct Goblet cell | 7. Hyaline cartilage 8. Intercalated duct 9. Islet of Langerhans 10. Salivary gland 11. Stratified squamous epithelium 12. Taste bud | |
| 0. 000.0. 00 | 121 10010 200 | |

Figure 13.148. Pancreas, Pig. An interlobular duct with goblet cells interspersed among the epithelial cells of the duct.

Figure 13.149. Pancreas, Cow (Masson's). An intercalated duct enters an acinus.

Figure 13.150. Pancreas, Sheep (Masson's). In ruminants, darkly stained A cells are located at the periphery of the islets of Langerhans, while light-staining B cells are centrally located. Compare with Figure 13.146.

Figure 13.151. Tongue, Tip, I.s., Chicken. The upper surface of the tongue is covered by a thick stratified squamous epithelium, which is keratinized near the tip. The stratified squamous epithelium of the lower surface is thinner and also keratinized rostrally. The tongue is supported by hyaline cartilage rostrally. The ducts of salivary glands (mucous) open at the lower surface.

Figure 13.152. Taste Bud, Tongue, Base, Chicken. A taste bud (characteristically large and scarce in the chicken) can be seen closely associated with the duct of a salivary gland. For detail of the taste bud, see Figure 13.153.




Figure 13.153 ×250



Figure 13.154 ×25




Figure 13.155 ×25




Figure 13.156 ×12.5




Figure 13.157 ×62.5

| | KEY |
|------------------------|------------------------------------|
| 1. Gland | 10. Proventriculus, lumen |
| 2. Intercellular space | 11. Secondary duct |
| 3. Lamina propria | 12. Serosa |
| 4. Mucous gland | 13. Stratified squamous epithelium |
| 5. Muscularis externa | 14. Submucosa |
| 6. Muscularis mucosae | 15. Sulcus |
| 7. Plica | 16. Taste bud cell |
| 8. Pore | 17. Tertiary duct |
| 9. Primary duct | , |


Figure 13.153. Taste Bud, Chicken. Detail of the taste bud seen in Figure 13.152. Its cells stain lightly with eosin. Numerous spaces occur between cells. A taste bud pore is visible.


Figure 13.154. Esophagus, x.s., Chicken. The esophagus is lined by a thick nonkeratinized stratified squamous epithelium. Mucous glands occur in the lamina propria. The submucosa is sparse.


Figure 13.155. Crop, Chicken. The crop is a diverticulum of the esophagus. Unlike the latter, it lacks mucous glands, except close to its junction with the esophagus.

Figure 13.156. Proventriculus (Glandular Stomach), x.s., Chicken. The submucosa contains lobules of compound tubular glands that are arranged around a central, secondary duct. A primary duct, which drains several lobules, opens through a raised mucosal papilla.

Figure 13.157. Proventriculus, x.s., Chicken. A magnified view of Figure 13.156 shows that the mucosa of the papilla is arranged into folds (plicae) covered by columnar cells, and depressions (sulci) lined by shorter cells. A primary duct, lined by columnar cells, joins the lumen of the proventriculus.










Figure 13.161 ×25


| KEY | | |
|---|--|--|
| 1. Basal cell 2. Chief cell 3. Gastric gland 4. Gastric pit 5. Gland cell | 8. Muscularis externa 9. Secondary duct 10. Submucosa 11. Surface epithelium 12. Tertiary duct | |
| 6. Keratinoid 7. Lamina propria | 13. Ventriculus, lumen | |


Figure 13.158. Proventriculus, x.s., Chicken. A tertiary duct branching from the secondary duct leads into a glandular unit. The glandular epithelial cells vary from simple cuboidal to low columnar and contain a grainy secretory material. These cells secrete both pepsinogen and HCl.


Figure 13.159. Ventriculus (Gizzard), Chicken. The pink, thick layer of keratinoid that lines the ventriculus is produced by branched tubular gastric glands in the lamina propria. The submucosa and a portion of the smooth muscle layers of a thick muscularis externa are shown.

Figure 13.160. Ventriculus, Chicken. Keratinoid lines the mucosal surface and fills the lumen of the gastric pits and glands. The cells of the surface epithelium are low to tall columnar. The cells decrease in height as they extend into the gastric pits. Flattened cells (chief cells) line the upper and mid-regions of the tubular gastric glands seen in this micrograph. Branching of some of the glands is evident.

Figure 13.161. Ventriculus, Chicken. The flattened chief cells lining the mid-portion of the gastric glands become cuboidal to low columnar in the fundus of the gland. A few large basal cells with pale nuclei and pale cytoplasm occur in the fundus of the glands.










Figure 13.165 ×62.5




Figure 13.166 $\times 25$

| KEY | | |
|--|--|--|
| 1. Crypt of Lieberkühn 2. Lymphatic tissue 3. Muscularis externa | 5. Serosa 6. Submucosa 7. Villus | |
| 1 Muscularis mucosae | | |

Figure 13.162. Duodenum, x.s., Chicken. Lymphatic tissue (diffuse and nodular) in the duodenum close to the stomach.

Figure 13.163. Duodenum, x.s., Chicken (Mallory's). The serosa is thick. The submucosa is characteristically sparse.

Figure 13.164. Ileum, x.s., Chicken. Villi are long and slender with numerous goblet cells. The submucosa is thin.

Figure 13.165. Cecum, Tip, I.s., Chicken. In the tip of the cecum, villi are short and broad. Compare Figure 11.51 of the cecal

Figure 13.166. Large Intestine, x.s., Chicken. Villi are present in the chicken's large intestine.

Figure 13.164




Figure 13.167



Figure 13.168 ×62.5



Figure 13.169 ×62.5



Figure 13.170 ×62.5

KEY

- 1. Cloacal lip, dorsal
- 2. Cloacal lip, ventral
- 3. Coprodeum
- 4. Crypt of Lieberkühn
- 5. Herbst corpuscle
- 6. Lymphatic nodule
- 7. Lymphoglandular ridge
- 8. Proctodeum


- 9. Simple columnar epithelium
- 10. Skeletal muscle
- 11. Smooth muscle
- 12. Stratified squamous epithelium
- 13. Tubular gland
- 14. Urodeum
- 15. Villus


Figure 13.167. Cloaca, I.s., Chicken. The cloaca is subdivided into three regions: the coprodeum, urodeum, and proctodeum. The large intestine is continuous with the coprodeum. The ureters and genital ducts terminate in the urodeum. The terminal proctodeum opens to the exterior through the cloacal lips.

Figure 13.168. Coprodeum, Cloaca, Chicken. The mucosa of the coprodeum is thrown into short, flat villi. Shallow crypts of Lieberkühn open at their bases. Simple columnar epithelium covers their surface.

Figure 13.169. Cloaca, l.s., Chicken. Branched tubular glands (modified crypts) form a part of the lymphoglandular ridge of Jolly located in the dorsal proctodeum. The epithelium of the ridge consists of tall columnar cells. A portion of the stratified squamous epithelium of the inner surface of the ventral cloacal lip can be seen.

Figure 13.170. Cloaca, I.s., Chicken. A large Herbst corpuscle lies beneath the stratified squamous epithelium of the cloacal lip. These corpuscles typically occur close to the latter's point of juncture with the simple columnar epithelium of the proctodeum.








Figure 13.173 ×250




Figure 13.174 ×125




Figure 13.175 ×250

| KEY | |
|-----------------|----------------------|
| 1. A cells | 6. Erythrocytes |
| 2. Acinus | 7. Hepatocyte |
| 3. B cells | 8. Lobule, margin of |
| 4. Capillary | 9. Reticular fibers |
| 5. Central vein | 10. Sinusoid |

Figure 13.171. Liver, Chicken. Central veins of several lobules are evident. One lobule is indicated by a dashed line.

Figure 13.172. Liver, Chicken. Sinusoids can be seen entering a central vein. Radiating plates of hepatocytes are two cells thick in the chicken.

Figure 13.173. Liver, Chicken (Silver). The wall of a central vein and surrounding plates of hepatocytes (whose cellular features are indistinct in this preparation) are supported by a network of reticular fibers that have been blackened with silver.

Figure 13.174. Pancreas, Chicken. Numerous portions of tubuloacinar serous glands surround an alpha islet of Langerhans. Alpha islets contain columnar A cells and are larger than beta islets. Erythrocytes in capillaries can be seen between cords of islet cells.

Figure 13.175. Pancreas, Chicken. Beta islets contain polygonal B cells and are smaller than alpha islets.

URINARY SYSTEM

MAMMALS

he urinary system of mammals is comprised of the paired kidneys, renal pelvises, ureters, urinary bladder, and urethra.

Kidneys

The kidneys are highly vascularized, compound tubular glands that function to maintain the composition of body fluids at a constant level and remove excretory wastes. Each kidney is surrounded by a capsule of connective tissue, which may contain a distinct layer of smooth muscle in its deepest portion, as in the cow, sheep, and goat. Both the cortex and medullary regions of the kidney are formed principally of numerous closely packed, uriniferous tubules. The spaces between tubules are mainly occupied by an extensive capillary network. The cortex and medulla are arranged into one or more pyramidal configurations called renal pyramids; the apex of each pyramid is called a renal papilla. In the cortex, groups of radially arranged tubules form the pars radiata (cortical ray or medullary rays), consisting of collecting tubules and straight portions of nephrons. The pars convoluta (cortical labyrinth) are located between the rays and consist of renal corpuscles and numerous proximal and distal convoluted tubules. The proximal convoluted tubules are longer than the distal convoluted tubules and comprise the major portion of the cortex. Proximal convoluted tubules are distinguished by the brush borders of their epithelial cells and the somewhat scalloped appearance of the apical surface of their cells when the latter are seen in profile. Distal convoluted tubules have a smooth internal surface, and their cells lack a brush border.

Filtrate processed by the nephrons is passed to collecting tubules, which open either directly or indirectly via calyces into the renal pelvis through papillary ducts at the tip of a renal papilla. The epithelial cells of the collecting tubules are pale and vary from

cuboidal near the distal tubules to columnar close to the papilla. Cell boundaries are normally clearly defined compared with the cells of the proximal and distal convoluted tubules. As they progress toward the renal papilla, the collecting tubules become wider. The terminal portion of these tubules is lined by a columnar or pseudostratified epithelium and is called the papillary duct.

Each renal corpuscle consists of a Bowman's capsule and glomerulus. The outer layer of Bowman's capsule is the parietal layer, a simple squamous layer. The inner layer is the visceral layer. It is formed of highly branched podocytes that surround the capillary loops of the glomerulus. In most histologic preparations made for light microscopy, it is not possible to distinguish podocytes from the adjacent endothelial cells of the capillary loops. The cavity between the visceral and parietal layers is the urinary space. The latter is continuous with the lumen of a proximal convoluted tubule at the urinary pole of each corpuscle. At the opposite, vascular pole, an afferent arteriole and efferent arteriole unite with the capillaries of the glomerulus. A portion of the distal convoluted tubule is positioned between the afferent and efferent arterioles. The macula densa of the juxtaglomerular apparatus forms a part of the wall of the distal convoluted tubule in this region. Each macula is composed of closely grouped epithelial cells and is easily identified by the tightly packed nuclei of these cells. **Juxtaglomerular cells** are modified cells of smooth muscle in the walls of afferent arterioles close to the glomerulus. They have an epithelioid appearance when seen in cross section.

The medulla of each kidney is formed from collecting tubules, thick and thin segments of the loops of Henle, and numerous vasa rectae. Thick descending portions of Henle's loop are continuations of the proximal convoluted tubules and are located close to the corticomedullary junction. They are straight tubules whose cells are lined by a brush border. Each thick descending tubule joins abruptly with a thin segment whose wall is formed from flattened cells with round, bulging nuclei. The straight, thick ascending portion of each loop resembles the distal convoluted tubule with which it is continuous.

Ureters, Urinary Bladder, and Urethra

The walls of the renal pelvis, ureter, urinary bladder, and urethra include a mucosa, muscularis of smooth muscle, and adventitia. A submucosa may be present. The lining of the mucosa is almost exclusively transitional epithelium. The hilus region, between the capsule of the kidney and the outer wall of the renal pelvis, contains loose connective tissue and adipose tissue.

The mucosa of the **ureter** presents a folded appearance. Its transitional epithelium is separated from the muscularis by a lamina propria. Tubuloalveolar mucous glands occur in the lamina propria of the first several centimeters of the ureter of the horse. A submucosa is lacking in the ureter. The muscularis consists of inner longitudinal, middle circular, and outer longitudinal layers. An adventitia of loose connective tissue surrounds the muscularis.

The transitional epithelial cells of the **urinary bladder** become flattened when the bladder is distended with urine.

A lamina propria and submucosa are present. Usually, there is a thin muscularis mucosae between these layers. The muscularis, external to the submucosa, is composed of an outer and inner longitudinal and a thick middle circular layer. The inner and outer longitudinal layers may be incomplete in some areas. Much of the bladder (body and apex) is covered by a serosa. An adventitia of loose connective tissue is present at the neck of the bladder.

CHICKEN

The urinary system of the chicken consists of large, elongated, paired kidneys. Ureters drain each kidney and open into the urodeum of the cloaca. There is no renal pelvis or urinary bladder in the bird.

Kidney

Each kidney of the chicken is partitioned into a cranial, middle, and caudal subdivision. Each subdivision is comprised of lobules. A lobule consists of a large cortical and a smaller medullary component. All of the lobules that drain into a single branch of the ureter constitute a lobe.

There are two types of nephrons. The cortical (reptilian) type is more numerous and lacks a loop of Henle. It is located entirely within the cortex. The other is the less numerous medullary (mammalian) type. It has a loop of Henle (also called a medullary loop), which extends into the medulla. Cortical nephrons are arranged radially around central (intralobular) veins of the cortex. Their renal corpuscles lie approximately midway between the intralobular vein and a peripheral interlobular vein. The cortical nephron has a smaller renal corpuscle than the medullary nephron. The large renal corpuscles of medullary nephrons lie close to the medulla. Other than size, there is no structural difference between small and large renal corpuscles. Each glomerulus contains a compact mass of mesangial cells (small cells with large nuclei) at its center. The mass appears basophilic because of the relatively high concentration of nuclear material. A layer of podocytes, with large round or oval nuclei, covers the surface of the glomerular capillaries, forming the visceral layer of Bowman's capsule. The parietal layer of Bowman's capsule consists of a simple squamous epithelium. Juxtaglomerular cells and a macula densa are associated with the renal corpuscle at its vascular pole.

Generally, cortical tissue located between renal corpuscles and interlobular veins consists mainly of proximal convoluted tubules, and that located between renal corpuscles and intralobular veins is comprised of distal convoluted tubules. Cells of proximal convoluted tubules are low columnar and have a well-developed brush border. Distal convoluted tubules are shorter than proximal convoluted tubules. Their cuboidal cells lack a brush border, but the apex may form a projecting bleb of clear cytoplasm that fills much of the lumen. In cortical nephrons a short intermediate tubule (which is without a brush border, and which is about half the diameter of a distal convoluted tubule) connects proximal convoluted tubules to distal convoluted tubules. In medullary nephrons long or short


Henle's (medullary) loops connect proximal convoluted tubules to distal convoluted tubules. The thin segment of a medullary loop forms only a part of the descending limb. Hence, thin segments are less numerous than either thick descending or thick ascending portions of the loop. The diameter of a thin segment is about one-half that of a thick segment. The cells of the thin segment are cuboidal and do not stain as intensely as the cuboidal cells of the thick segments. Apical cytoplasmic blebs of the cells of the thick segments project into the lumen. Collecting tubules occur in the more peripheral parts of the cortex. They are lined by pale cells with cuboidal to low columnar shape and are intermediate in size between proximal convoluted and distal convoluted tubules. Collecting tubules join distal convoluted tubules to perilobular collecting ducts. The latter unite with those of other lobules to form medullary tracts, each of which is surrounded by a thin capsule of connective tissue. Tracts group together to form a medullary cone. Each cone terminates in a single branch of the ureter. Cones and tracts contain thin and thick segments of medullary loops, in addition to collecting ducts. The


lining epithelium of the smallest collecting ducts is simple cuboidal. It gradually becomes simple columnar and finally changes to pseudostratified columnar in the proximity of the ureteral branch.

Ureter

The ureter of the chicken is a muscular duct about 2 mm in diameter. Its wall consists of a mucosa, muscularis, and adventitia. It is generally lined by a pseudostratified columnar epithelium. The majority of cells are tall with a lesser number of cuboidal cells lying close to the basement membrane. The apices of the columnar cells contain numerous vacuoles filled with mucus. Beneath the epithelium is a thick layer of loose connective tissue containing varying amounts of diffuse lymphatic tissue and, sometimes, a lymphatic nodule. The muscularis consists of an inner longitudinal and outer circular layer of smooth muscle. A third outer longitudinal layer is present near the cloaca. The adventitia consists of a layer of loose connective tissue.

| Word Roots | | |
|-------------|-------------------------------|---|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Convolve | Rolled together, coiled up | The proximal <i>convoluted</i> tubule is highly coiled. |
| Glomer | A ball of yarn, a ball | The <i>glomerulus</i> of a renal corpuscle is formed of loops of capillaries resembling a ball of yarn. |
| Juxta | Near | Juxtaglomerular cells are close to the glomerulus. |
| Macul | Spot, spotted | The nuclei of the closely grouped cells of the <i>macula</i> densa give it a spotted appearance. |
| Mes Angi | Middle Vessel | Mesangial cells are evident in the center of the glomerular capillaries of the chicken. |
| Neph | Kidney | The dog has about 400,000 nephrons per kidney. |
| Ren | Kidney | <i>Renal</i> corpuscles are in the cortex of the kidney. |
| Ure | Urine | The <i>ureters</i> and <i>urethra</i> convey urine. |





KEY

- 1. Adipose tissue
- 2. Capsule
- 3. Medulla


- 4. Pars convoluta
- 5. Pars radiata
- 6. Renal corpuscle


Figure 14.1. Cortex and Portion of Medulla, Kidney, Dog. Renal corpuscles are limited to the cortex.


Figure 14.2. Cortex, Kidney, Dog (Masson's). Pars radiata alternate with the pars convoluta.

Regions of the kidney and their major components:

- A. Cortex of the Kidney
 - 1. Pars convoluta (cortical labyrinth): Renal corpuscles, proximal and distal convoluted tubules, peritubular capillaries, portions of collecting tubules
- Pars radiata (medullary rays): Collecting tubules and portions of the loops of Henle
 Medulla: Collecting tubules, thick and thin segments of
- the loops of Henle, and vasa recta










Figure 14.6 \times 250

KEY

- 1. Bowman's capsule, parietal layer 7. Macula densa
- 2. Capsule of kidney
- 3. Collecting tubule
- 4. Connective tissue
- 5. Distal convoluted tubule
- 6. Fat vacuole

- 8. Nucleus of podocyte
- 9. Proximal convoluted tubule
- 10. Smooth muscle tissue
- 11. Subcapsular vein
- 12. Urinary space

Figure 14.3. Capsule and Superficial Cortex, Kidney, Cat. The capsule of the kidney consists entirely of connective tissue in the


Figure 14.4. Capsule, Kidney, Sheep. The inner portion of the capsule of the kidney of ruminants contains a distinct layer of smooth muscle. Smooth muscle is also present in the capsule of the dog, horse, and pig.


Figure 14.5. Renal Corpuscle, Kidney, Puppy. In young animals, the podocytes of the visceral layer of Bowman's capsule have round to oval nuclei.


Figure 14.6. Renal Corpuscle, Kidney, Cat. Cells of the proximal convoluted tubules of the cat contain numerous fat vacuoles. A macula densa, consisting of closely packed cells, forms a portion of the wall of the distal convoluted tubule adjacent to the vascular pole of the renal corpuscle.

Synonyms

Bowman's capsule = glomerular capsule Parietal layer of Bowman's capsule = capsular epithelium Urinary space = capsular space Viscerál layer of Bowman's capsule = glomerular epithelium










Figure 14.10 ×250

KEY

- 1. Afferent arteriole
- 2. Bowman's capsule, parietal layer
- 3. Brush border
- 4. Collecting tubule
- 5. Distal convoluted tubule
- 6. Juxtaglomerular cells
- 7. Macula densa
- 8. Proximal convoluted tubule
- 9. Urinary space

Figure 14.7. Cortex, Kidney, Horse. Portions of three renal corpuscles, each with an accompanying macula densa, are present. In the horse, the macula densa commonly consists of a stratified layer of cells.

Figure 14.8. Cortex, Kidney, Horse. A collecting tubule with clearly defined cells and a smooth lining can be contrasted with proximal convoluted tubules whose cells possess a brush border of microvilli.

Figure 14.9. Cortex, Kidney, Pig. The junction of a proximal convoluted tubule with the Bowman's capsule of a renal corpuscle is shown.

Figure 14.10. Afferent Arteriole, Kidney, Pig. An afferent arteriole, with juxtaglomerular cells, is entering a glomerulus. The juxtaglomerular cells are epithelioid. Note that a macula densa borders the afferent arteriole.

Summary of histologic features of the nephron and collecting tubule

I. Nephron

A. Renal corpuscle

- 1. Bowman's capsule
 - a. Parietal layer: Simple squamous epithelium
 - b. **Urinary space:** The cavity between the parietal and visceral layers
 - c. **Visceral layer:** Formed of podocytes, which are cells with numerous processes that surround capillaries of the glomerulus; usually it is difficult to distinguish between podocytes and the endothelial cells of the glomerular capillaries
- thelial cells of the glomerular capillaries

 2. **Glomerulus:** A tuft of capillaries within the Bowman's capsule
- B. **Proximal convoluted tubule:** Lined by plump simple cuboidal cells that have brush borders of microvilli, which give the lumen a somewhat scalloped appearance; cell boundaries are indistinct; proximal convoluted tubules are longer than distal convoluted tubules and therefore more obvious in sections through the cortex.

C. Loop of Henle

- Thick descending portion: Lined by cells that resemble the proximal convoluted tubule with which it is continuous
- Thin segment: Formed of a single layer of squamous cells with round, bulging nuclei
- Thick ascending portion: Epithelial cells resemble the distal convoluted tubule with which it is continuous
- D. Distal convoluted tubule: Lined by simple cuboidal cells smaller than those of the proximal convoluted tubule and without a brush border, so that the lumen has a smoother internal surface; cell boundaries are indistinct
- II. Collecting tubule: Epithelial cells vary from simple cuboidal to simple columnar; cell boundaries between the cells are evident

Papillary duct: Terminal portion of the collecting tubule lined by simple columnar or pseudostratified epithelium




Figure 14.11 ×250




Figure 14.12 ×250




Figure 14.13 ×12.5




Figure 14.14 ×250




Figure 14.15 ×250

KEY

- 1. Adipose tissue
- 2. Afferent arteriole
- 3. Cavity of renal pelvis
- 4. Collecting tubule
- 5. Collecting tubule, surface cut
- 6. Efferent arteriole
- 7. Macula densa
- 8. Papillary duct

- 9. Proximal convoluted tubule
- 10. Renal papilla
- 11. Thick ascending, Henle's loop
- 12. Thick descending, Henle's loop
- 13. Thin segment, Henle's loop
- 14. Urinary space
- 15. Vasa recta

Figure 14.11. Afferent and Efferent Arterioles, Kidney, Pig. Junction of a glomerulus with an afferent arteriole. A macula densa parallels the afferent arteriole.

Figure 14.12. Pars Radiata, I.s., Cortex, Kidney, Horse. The component tubules of a medullary ray include collecting tubules as well as thick descending and thick ascending portions of the loop of Henle.

Figure 14.13. Renal Papilla, Kidney, Dog. Papillary ducts open onto the tip of a renal papilla.

Figure 14.14. Medulla, Kidney, Dog. Nuclei of the cells lining the thin segment of Henle's loop are rounded; those of endothelial cells of the vasa recta are flattened and more darkly stained.

Figure 14.15. Medulla, Kidney, Horse. Various portions of uriniferous tubules appear in transverse section.




Figure 14.16 ×180




Figure 14.17 ×90

| KEY | |
|---------------------------|----------------------------------|
| 1. Cavity of renal pelvis | 4. Thick ascending, Henle's loop |
| 2. Collecting tubule | 5. Thin segment, Henle's loop |
| 3. Papillary duct | 6. Vasa recta |

Figure 14.16. Medulla, Kidney, Cow (Trichrome). Longitudinal sections of vasa rectae and portions of uriniferous tubules. The vasa rectae are filled with red blood cells (stained orange).

Figure 14.17. Renal Papilla, Kidney, Goat. Papillary ducts near the tip of a renal papilla are lined by transitional epithelium.



Figure 14.18 ×250




Figure 14.19 ×250




Figure 14.20 ×12.



Figure 14.21 ×62.5

KEY

- 1. Adventitia
- 2. Collecting tubule
- 3. Erythrocytes
- 4. Lamina propria
- 5. Mucous connective tissue
- 6. Muscularis


- 7. Papillary duct
- 8. Reticular fiber
- 9. Thin segment, Henle's loop
- 10. Transitional epithelium
- 11. Vasa recta

Figure 14.18. Medulla, Kidney, Goat. A papillary duct, l.s., some distance away from the apex of the papilla, is lined by columnar cells. A thin segment of Henle's loop parallels the duct.

Figure 14.19. Medulla, Kidney, Goat (Silver). The collecting tubules are encircled by reticular fibers (stained black). The fibers provide a supportive framework for other portions of the uriniferous tubules as well.

Figure 14.20. Urachus, Umbilical Cord, Cow. The urachus (allantoic stalk) is lined by a transitional epithelium. A portion of the lining is shown.

Figure 14.21. Ureter, x.s., Cat (Masson's). The middle circular layer of smooth muscle of the muscularis is most evident. Inner and outer longitudinal layers are present but sparse in this section.








Figure 14.23 ×12.5




Figure 14.24 ×125




Figure 14.25 ×12.5




Figure 14.26 ×250

| KEY | | |
|--|---|--|
| Capillary Lamina propria Mucous gland Muscularis | 5. Muscularis mucosae6. Submucosa7. Transitional epithelium | |
| | | |


Figure 14.22. Ureter, x.s., Horse (Masson's). The proximal (anterior) portion of the horse's ureter contains tubuloalveolar mucous glands.


Figure 14.23. Ureter, x.s., Horse. Distally (posteriorly), the horse's ureter lacks mucous glands. The muscularis consists of an inner longitudinal, middle circular, and outer longitudinal layer of smooth muscle.

Figure 14.24. Urinary Bladder, Pig. The mucosa to a portion of the muscularis is shown. Scattered muscle cells of the muscularis mucosae are located adjacent to the lamina propria.

Figure 14.25. Urinary Bladder, Cow. The bladder contains a muscularis mucosae between the lamina propria and submucosa. Only a portion of the thick muscularis is shown.

Figure 14.26. Urinary Bladder, Goat. Numerous capillaries are located beneath the transitional epithelial lining of the bladder of ruminants.








Figure 14.29 ×12


| KEY | |
|--|---|
| Cortex Distal convoluted tubule Intralobular vein Medullary cone | 5. Proximal convoluted tubule6. Renal corpuscle, large7. Renal corpuscle, small |

Figure 14.27. Kidney, Chicken. Cortical parenchyma and portion of a medullary cone are shown. An intralobular vein and both cortical (small) and medullary (large) renal corpuscles are apparent.

Figure 14.28. Kidney, Chicken. A portion of a medullary cone is surrounded by cortical lobules. The intralobular veins of two cortical lobules are clearly represented.

Figure 14.29. Cortex, Kidney, Chicken. An intralobular vein is surrounded by cortical tissue. Distal convoluted tubules are located mainly in the region of the intralobular vein.

Figure 14.28 ×36 URINARY SYSTEM 193








Figure 14.31 ×250



Figure 14.32 ×90

| K | Y |
|-------------------------------------|-----------------------------------|
| 1. Bowman's capsule, parietal layer | 9. Muscularis, outer layer |
| 2. Collecting duct | 10. Nucleus of podocyte |
| 3. Collecting tubule | 11. Proximal convoluted tubule |
| 4. Distal convoluted tubule | 12. Pseudostratified epithelium |
| 5. Erythrocytes | 13. Serosa |
| 6. Lymphatic tissue | 14. Thick ascending, Henle's loop |
| 7. Mesangial cells | 15. Thin segment, Henle's loop |
| 8. Muscularis, inner layer | 16. Urinary space |

Figure 14.30. Cortex, Kidney, Chicken. In the chicken, the visceral layer of Bowman's capsule is composed of podocytes that have large round or oval nuclei. The center of the glomerulus contains a compact mass of mesangial cells.

Figure 14.31. Medullary Cone, Kidney, Chicken. Various portions of uriniferous tubules (medullary type) are evident. Cells lining the thick ascending portions of Henle's loop show characteristic clear cytoplasmic blebs. A small portion of the cortex containing a proximal convoluted tubule can be seen on the upper right side.

Figure 14.32. Ureter, x.s., Chicken. The lamina propria is infiltrated with lymphocytes. Outer circular and inner longitudinal layers of the muscularis are distinguishable. The epithelium is pseudostratified columnar.

RESPIRATORY SYSTEM

MAMMALS

Air flows from the nostrils through a system of passages to the respiratory surfaces of the lungs. As it progresses, it becomes warmed, humidified, and cleansed of some of its particulate matter. Dust, which finds its way to the alveoli, is ultimately consumed by macrophages patrolling the tiny cul-de-sacs. The major components of the air-passage system are the **nasal cavity**, **pharynx**, **larynx**, **trachea**, **bronchi**, and the various smaller subdivisions of the **bronchial tree** leading to the **alveoli**.

Nasal Cavity

Air from the naris enters the **vestibule**, the first part of the nasal cavity. The vestibule is lined by a stratified squamous epithelium, which is continuous with the skin externally and with the respiratory portion of the nasal cavity internally. In the horse, hairy skin continues into the vestibule. A lamina propria and underlying submucosa support the vestibular epithelium.

The respiratory portion of the nasal cavity is lined by a ciliated, pseudostratified columnar epithelium with goblet cells. The lamina propria contains tubuloalveolar glands. The latter are mainly serous, but mucous and mixed glands do occur. Glands are sparse in carnivores. A submucosa supports the lamina propria.

The olfactory epithelium (pseudostratified columnar) is composed of olfactory (sensory), supporting, and basal cells. Bowman's glands, tubular and mucoserous, occur within the lamina propria. They open to the surface through ducts lined by cuboidal or flattened cells. A submucosa lies below the lamina propria.

Pharynx

The nasopharynx and oropharynx are subdivisions of the pharynx. The former is lined by a ciliated, pseudostratified columnar epithelium with goblet cells, whereas the latter is covered by a stratified squamous epithelium. The lamina propria contains tubular mixed glands in the nasopharynx and mucous glands in the oropharynx. In carnivores, the glands of the oropharynx are mixed. A network of elastic fibers separates the mucosa from an underlying sheet of skeletal muscle consisting of circularly and longitudinally arranged cells. The musculature is separated from an adventitia of loose connective tissue by a layer of connective tissue containing elastic networks.

Larynx

The larynx is lined in part by a stratified squamous epithelium and partly by a ciliated, pseudostratified columnar epithelium. Numerous elastic fibers are present in the lamina propria. Glands (serous, mucous, and mixed) occur in the lamina propria and submucosa, but are lacking in the vocal and vestibular folds. Hyaline and elastic cartilage provide support for the laryngeal wall. The elastic cartilage of the epiglottis may be partially or completely replaced by adipose tissue, as in carnivores. Skeletal muscles are an integral part of the laryngeal structure.

Trachea

The trachea is lined by a ciliated, pseudostratified columnar epithelium with goblet cells. A lamina propria and submucosa lie below the epithelium, but are not clearly demarcated from one another. Glands, mostly mixed, occur in the deeper layers of the lamina propria and within the submucosa. Rings of hyaline cartilage, which are incomplete dorsally, support the tracheal wall. A layer of smooth muscle, the trachealis muscle, is located dorsally in the trachea. It is positioned internal to the gap in the tracheal cartilages in the horse, pig, and ruminants. It lies external to the gap in the cat and dog. An adventitia of connective tissue completes the wall of the trachea.

Bronchial Tree and the Lungs

The trachea bifurcates into bronchi, which enter the lungs, where they branch extensively. The lungs are covered by a visceral pleura, which is thick in large mammals and thinner in carnivores. Connective tissue and some smooth muscle form a part of the visceral pleura. The interior of the lungs contains a framework of connective tissue, rich in elastic fibers, which supports the bronchial tree and divides the lungs into lobules. The interlobular connective tissue is sparse in carnivores.

A ciliated, pseudostratified columnar epithelium with goblet cells lines the bronchi. The epithelium becomes reduced in height as the caliber of the bronchi diminishes. The lamina propria is surrounded by a layer of obliquely arranged smooth muscle. The connective tissue external to the musculature contains mixed glands and plates of hyaline cartilage. In the cat, the bronchial cartilages may contain elastic fibers. When seen

in histologic sections, the mucosa of large bronchi has few folds. Folds increase as the bronchi decrease in diameter.

The smallest bronchi give rise to suborders of bronchioles. The smallest of the latter, the terminal bronchioles, branch into two or more respiratory bronchioles, which divide into alveolar ducts that, in turn, empty into alveolar sacs.

Bronchioles lack cartilage and glands. Glands, however, may extend into bronchioles from bronchi in cats. Spirally or obliquely arranged smooth muscle forms part of the wall of a bronchiole. The amount of smooth muscle is proportional to the size of the bronchiole. Large bronchioles are lined by ciliated columnar cells, whereas the smallest (terminal) bronchioles are lined proximally by ciliated cuboidal cells and, distally, by nonciliated cells. The mucosa of the bronchioles is folded, unless the lungs were inflated at the time when the tissue was processed.

Respiratory bronchioles branch from the ends of terminal bronchioles. They are lined by a cuboidal epithelium, which becomes flattened distally, and their wall contains some smooth muscle. Alveoli are scattered within the epithelium. Respiratory bronchioles are best developed in the cat and dog.

Alveolar ducts branch from respiratory bronchioles. Their thin walls are constructed entirely of alveoli. The lip of each alveolus of an alveolar duct contains smooth muscle arranged circumferentially. The presence of the muscle gives the lip of the alveolus a knoblike appearance when histologic sections occur at right angles to the long axis of the muscle cells.

Ultimately, each alveolar duct branches into three or more alveolar sacs. No smooth muscle is present in the sacs. Therefore, the alveoli, which form the walls of the sacs, do not have lips with knoblike expansions as do those of the alveolar ducts.

Alveoli are lined mainly by exceedingly thin squamous epithelial cells (type I alveolar cells) and fewer type II alveolar cells, which produce surfactant. Alveoli are separated from one another by a thin, highly vascularized layer of fine collagenous and elastic fibers. This layer, together with the squamous cells lining the adjacent alveoli, forms an alveolar septum.

CHICKEN

The nostrils, nasal cavity, pharynx, trachea, syrinx, bronchi, air capillaries, and air sacs comprise the respiratory system of the bird.

Nasal Cavity

The skin enters the nostrils to the first part of the nasal cavity, the vestibule, which is lined by a modified, keratinized, stratified squamous epithelium. It is characterized by epithelial cells that are organized into columns, giving the surface a wavy appearance. The respiratory region of the nasal cavity is lined by a ciliated, pseudostratified columnar epithelium. Mucous glands occur within the respiratory epithelium. The olfactory epithelium is pseudostratified columnar. It is located in the upper portions of the respiratory regions. Its structure, like that of mammals, is composed of basal, sensory, and supporting cells. Bowman's glands are present.

Pharynx

The pharynx is lined by a stratified squamous epithelium. A dense lamina propria and less dense submucosa lie below the epithelium. Salivary glands (mucous) occur within the lamina propria or submucosa. Bundles of skeletal muscle occur below the floor of the pharynx.

Larynx, Trachea, and Syrinx

At the anterior end of the trachea is a cranial larynx, which is reinforced by a cartilaginous ring. A caudal larynx (syrinx) is located at the posterior end of the trachea. The trachea is supported by overlapping, complete cartilaginous rings. It is lined by a ciliated, pseudostratified columnar epithelium containing numerous, simple alveolar mucous glands. In the posterior portion of the trachea, the glands are replaced by goblet cells. A lamina propria and submucosa are present. Each consists of dense connective tissue. The submucosa is rich in elastic fibers.

The syrinx, or voice box, is located in the thoracic cavity at the point of tracheal bifurcation into two bronchi. Internal and external tympanic membranes, located in the region of the tracheal bifurcation, characterize the wall of the syrinx. Intersyringeal cartilages and a bony wedge, the pessulus, provide support in the region of the syrinx.

Bronchi and Air Capillaries

Each extrapulmonary primary bronchus enters a lung as an intrapulmonary primary bronchus (mesobronchus). Secondary bronchi stem from the primary bronchi and branch into numerous parabronchi (tertiary bronchi) within the lung. The latter anastomose with each other. Tiny, respiratory air capillaries form extensive networks interconnecting the tertiary bronchi.

Primary bronchi are lined by a ciliated pseudostratified columnar epithelium with mucous glands and goblet cells. Extrapulmonary primary bronchi have C-shaped cartilages, while the walls of intrapulmonary primary bronchi contain cartilaginous plates, which become scarce distally. Bundles of smooth muscle, mainly circular, occur below the lamina propria. Numerous elastic fibers are found throughout the connective tissue of the bronchi.


Secondary bronchi are lined by a ciliated, columnar epithelium with mucous cells. There is a lamina propria and a well-developed muscularis.


Parabronchi are lined by a cuboidal epithelium. A thin layer of connective tissue lies below the epithelium. Bundles of smooth muscle cells lie below the connective tissue layer. The inner wall of each tertiary bronchus is pierced by numerous openings, each of which leads into a cavity called an atrium (air vesicle). Atria are lined by a squamous to cuboidal epithelium. Air capillaries, lined by squamous cells, open into atria. Their simple squamous lining is a respiratory surface and is analogous to the lining epithelium of the alveoli of the mammalian lung. Numerous vascular capillaries surround the air capillaries and are separated from the latter by a basement membrane.

Air Sacs

Air sacs are paired or unpaired, thin-walled structures occurring in the cervical, clavicular, thoracic, and abdominal regions of the body. They connect to the lungs by bronchi. Many of the hollow bones of the fowl contain extensions of the air sacs. Among others, these bones include the sternum, humerus, pelvic girdle, and most of the thoracic and cervical vertebrae. The air sacs are lined by squamous, ciliated cuboidal, and ciliated columnar cells. The epithelium is supported by a thin layer of connective tissue consisting of collagenous and elastic fibers. The sacs are poorly vascularized and do not participate in gas exchange.

| Word Roots | | |
|------------|--------------------------|---|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Alveol | A cavity, pit, socket | The bronchial tree of mammals ends in tiny, thin walled outpocketings called <i>alveoli</i> . |
| Bronch | Windpipe | The bronchi and bronchioles conduct air. |
| Olfact | Smell | Olfactory cells of the olfactory epithelium play a role in the sense of smell. |
| Trache | Windpipe | Air from the larynx is conveyed to bronchi by the <i>trachea</i> . |
| Viscera | The organs of the body | The <i>visceral</i> pleura covers the lungs. |








Figure 15.3 ×180




Figure 15.4 ×125

| rigure 15.4 | ×123 |
|-------------------|------------------------------------|
| | KEY |
| 1. Basal cells | 8. Olfactory cells, nuclei |
| 2. Bone | 9. Pseudostratified epithelium |
| 3. Bowman's gland | 10. Sensory hairs |
| 4. Cartilage | 11. Serous gland |
| 5. Cavernous vein | 12. Stratified squamous epithelium |
| 6. Goblet cell | 13. Supporting cells, nuclei |
| 7. Nasal cavity | |
| | |

Figure 15.1. Nasal Cavity, Vestibule, Dog. This portion of the vestibule is supported by hyaline cartilage and lined by a stratified squamous epithelium. Numerous cavernous veins occur throughout the connective tissue of the mucosa. In addition, there are tubular serous glands within the connective tissue.

Figure 15.2. Portion of Nasal Concha, Dog. The scroll-like nasal conchae are supported by spongy bone and are covered by a mucous membrane with a ciliated, pseudostratified columnar epithelium.

Figure 15.3. Olfactory Epithelium, Nasal Cavity, Dog (Masson's). This thick, pseudostratified columnar epithelium is composed of three types of cells. Basal cells are located at the level of the basement membrane. The nuclei of olfactory cells form a broad

three types of cells. Basal cells are located at the level of the basement membrane. The nuclei of olfactory cells form a broad band in the central portion of the epithelium. The nuclei of supporting cells are pale and form the uppermost level of nuclei. The apices of olfactory cells bear sensory hairs.

Figure 15.4. Respiratory Epithelium, Nasal Concha, Dog. The ciliated, pseudostratified columnar epithelium with goblet cells and underlying, vascular, loose connective tissue and bone are shown.




Figure 15.5 ×250




Figure 15.6 ×62.5




Figure 15.7 ×25




Figure 15.8 ×12.5




Figure 15.9 ×250




Figure 15.10 ×12.5

- Adipose tissue
 Arytenoid cartilage
- 3. Duct
- 4. Elastic cartilage
- 5. Intercalated duct
- 6. Lamina propria7. Mixed gland
- 8. Pseudostratified epithelium

KEY

- 9. Serous acinus
- 10. Skeletal muscle
- 11. Stratified squamous epithelium, nonkeratinized
- 12. Striated duct
- 13. Submucosa
- 14. Taste bud

Figure 15.5. Lateral Nasal Gland, Dog (Masson's). This serous gland is located in the maxillary sinus in carnivores.


Figure 15.6. Nasopharynx, Dog. This portion of the pharynx is lined by a ciliated, pseudostratified columnar epithelium with goblet cells. Mixed glands are present. The muscularis consists of skeletal muscle.

Figure 15.7. Epiglottis, Dog. The supporting elastic cartilage of the epiglottis is heavily infiltrated in its mid-region by adipose tissue in carnivores.

Figure 15.8. Epiglottis, l.s., Sheep. Blocklike chunks of elastic cartilage, without infiltrating adipose tissue, are found in the epiglottis of the sheep and goat.

Figure 15.9. Epiglottis, Sheep. Occasionally, taste buds are found in the epithelium of the laryngeal surface of the epiglottis.

Figure 15.10. Glottis, x.s., Goat. The glottis is supported by the arytenoid cartilages (elastic) and is lined by a nonkeratinized stratified squamous epithelium.








Figure 15.12 ×25



Figure 15.13 ×12.5




Figure 15.14 ×250




Figure 15.15 ×62.5

| | ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,, |
|------------------------|--|
| | KEY |
| 1. Adipose tissue | 10. Plasma cell |
| 2. Arytenoid cartilage | Pseudostratified epithelium |
| 3. Elastic fibers | 12. Serous gland |
| 4. Esophagus | 13. Skeletal muscle |
| 5. Goblet cell | 14. Stratified squamous epithelium |
| 6. Hyaline cartilage | 15. Submucosa |
| 7. Lamina propria | 16. Trachea |
| 8. Mixed gland | 17. Trachealis muscle |
| 9. Muscularis externa | 18. Vocal ligament |
| | |

Figure 15.11. Vocal Fold, l.s., Cat. Junction of the vocal fold with the arytenoid cartilage.

Figure 15.12. Vocal Fold, Goat (Orcein). The vocal fold consists of a fold of the mucous membrane. The vocal fold encloses the vocal ligament, which is a band of elastic fibers.

Figure 15.13. Trachea and Esophagus, x.s., Cat. Note that the trachealis muscle (smooth) lies external to the gap in the C-shaped cartilage in carnivores.

Figure 15.14. Trachea, x.s., Cat. The trachea is lined by a ciliated, pseudostratified columnar epithelium with goblet cells. The lamina propria contains a band of longitudinally oriented elastic fibers

Figure 15.15. Trachea, x.s., Horse. A thick band of longitudinally arranged elastic fibers extends from the lamina propria into the submucosa.



Figure 15.16 ×25




Figure 15.17 ×250




Figure 15.18 ×12.5




Figure 15.19 ×25




Figure 15.20 ×25

| KEY | | |
|--------------------------|---|--|
| 1. Adipose tissue | 7. Mixed glands | |
| 2. Elastic fibers | 8. Muscularis | |
| 3. Fibroelastic membrane | 9. Perichondrium | |
| 4. Goblet cell | Pseudostratified epithelium | |
| 5. Hyaline cartilage | 11. Tracheal glands | |
| 6. Lamina propria | 12. Trachealis muscle | |

Figure 15.16. Trachea, x.s., Cow. In noncarnivores, the trachealis muscle attaches to the perichondrium on the inside of the tracheal cartilage. The lamina propria and submucosa are both very rich in elastic fibers.

Figure 15.17. Trachea, x.s., Cow. The trachea is lined by a ciliated, pseudostratified columnar epithelium with goblet cells.

Figure 15.18. Trachea, x.s., Sheep (Orcein). A fibroelastic membrane surrounds the C-shaped tracheal cartilage and also spans the gap in the cartilage.

Figure 15.19. Trachea, x.s., Goat (Orcein). Numerous elastic fibers occur below the epithelium. Elastic fibers are also present where the trachealis muscle joins with the perichondrium.

Figure 15.20. Primary Bronchus, Extrapulmonary, x.s., Dog (Mallory's). Plates of hyaline cartilage support the wall of the bronchus. Smooth muscle bundles of the muscularis occur between the plates and internal to them.




Figure 15.21 ×25




Figure 15.22 ×62.5




Figure 15.23 ×25




Figure 15.24 ×62.5




Figure 15.25 ×250

| KEY | | |
|----------------------------------|---------------------------------|--|
| 1. Alveolus | 7. Lamina propria | |
| 2. Bronchiole | 8. Lymphatic nodule | |
| 3. Columnar epithelium, ciliated | 9. Mixed gland | |
| 4. Elastic band | 10. Muscularis | |
| 5. Goblet cell | 11. Pseudostratified epithelium | |
| 6. Hyaline cartilage | 12. Serous gland | |

Figure 15.21. Bronchus, x.s., Cow (Masson's).

Figure 15.22. Bronchus, x.s., Cow. Detail of the wall of a bronchus. Numerous lymphocytes are present below the epithelium.

Figure 15.23. Small Bronchus, x.s., and Bronchioles, Cat. Bronchioles lack cartilaginous plates and possess a simple epithelium.

Figure 15.24. Large Bronchiole, x.s., Cat. In cats, submucosal serous glands extend from bronchi into the bronchioles.

Figure 15.25. Large Bronchiole, x.s., Cat. Detail of a portion of the bronchiole shown in Figure 15.24.




Figure 15.26 ×125




Figure 15.27 ×125




Figure 15.28 ×12




Figure 15.29 ×62..




Figure 15.30 ×125

| KEY | | |
|------------------------|---------------------------|--|
| 1. Alveolar duct | 6. Elastic fibers | |
| 2. Alveolar sac | 7. Macrophage | |
| 3. Alveolus | 8. Muscularis | |
| 4. Bronchiole | 9. Respiratory bronchiole | |
| 5. Cuboidal epithelium | 10. Smooth muscle | |
| 5. Cuboladi epimelium | 10. Smooth muscle | |


Figure 15.26. Bronchioles, Pig (Orcein). The tissues of the lung are heavily infiltrated with elastic fibers.


Figure 15.27. Respiratory Bronchiole, x.s., Cat. Respiratory bronchioles are lined by a cuboidal epithelium and have alveoli in their walls.


Figure 15.28. Respiratory Bronchiole, I.s., Sheep. Note the cuboidal epithelium, alveoli, and the small amount of smooth muscle tissue in the wall of the respiratory bronchiole.

Figure 15.29. Alveolar Ducts and Alveolar Sacs, Sheep. An alveolar duct is characterized by the presence of smooth muscle arranged circumferentially in the lips of the alveoli that form its wall. Conversely, the alveoli of alveolar sacs lack smooth muscle.

Figure 15.30. Alveolar Duct, x.s., Sheep. Detail of an alveolar duct. The smooth muscle associated with the entrance of the alveoli that form the wall of the duct is evident.







11-

Figure 15.34 ×12.5

| KEY | | |
|-----------------------------|---------------------------------------|--|
| 1. Alveolar septum | 7. Macrophage | |
| 2. Alveolus | 8. Mesothelium | |
| 3. Bronchiole | 9. Septum | |
| 4. Bronchus | 10. Type II alveolar cell | |
| 5. Elastic band | 11. Visceral pleura | |
| 6. Erythrocyte in capillary | , , , , , , , , , , , , , , , , , , , | |

Figure 15.31. Alveoli, Cat. Detail of alveolar septa.

Figure 15.32. Visceral Pleura, Dog. The visceral pleura of carnivores is relatively thin.

Figure 15.33. Visceral Pleura, Horse. The visceral pleura of domestic mammals, except carnivores, is thick. Incomplete septa extend inward from the visceral pleura in the horse.

Figure 15.34. Visceral Pleura, Pig. Lungs are highly lobulated in the pig and ruminants. Unlike those of the horse, the septa are complete.



Figure 15.35 ×62.5




Figure 15.36 ×125




Figure 15.37 ×62.5




Figure 15.38 × 250




Figure 15.39 ×123

| | KEY |
|--------------------------|---------------------------------|
| 1. Alveolus | 9. Parietal pleura |
| 2. Basal cells | 10. Pseudostratified epithelium |
| 3. Epidermis | 11. Pulmonary vein |
| 4. Intercostal muscle | 12. Pyknotic nucleus |
| 5. Lamina propria | 13. Septum |
| 6. Mucous gland | 14. Smooth muscle |
| 7. Nerve, unmyelinated | 15. Turbinate cartilage |
| 8. Neuromuscular spindle | 16. Vestibular epithelium |
| | |

Figure 15.35. Parietal Pleura, Cat. Parietal pleura and intercostal muscle. The parietal pleura lines the wall of the pleural cavity. It consists of a mesothelium and underlying connective tissue.

Figure 15.36. Lung, Cow (Masson's). In the cow and pig, pulmonary veins have thick bands of circularly arranged smooth muscle.

Figure 15.37. Nasal Cavity, Chicken. The vestibule is lined by a uniquely structured, keratinized, stratified squamous epithelium (see Figure 15.38). The vestibular epithelium blends with the epidermis on the inner side of each nostril. In this micrograph, these epithelia lie to either side of a turbinate cartilage.

Figure 15.38. Vestibular Epithelium, Chicken. This keratinized, stratified squamous epithelium is characterized by the presence of columns of cells. The uppermost cells in each column have pyknotic nuclei. One or two layers of basal cells are present. The outer surface of the epithelium presents a corrugated appearance.

Figure 15.39. Respiratory Epithelium, Chicken (Masson's). This ciliated, pseudostratified columnar epithelium is interrupted by simple, branched, alveolar mucous glands.





Figure 15.40 ×250



×12.5 **Figure 15.41**




Figure 15.42 ×62.5




Figure 15.43 $\times 125$



Figure 15.44 $\times 25$

KEY

- 1. Basal cell
- 2. Bony tracheal ring
- 3. Bowman's gland
- 4. Cartilaginous tracheal ring
- 5. Mucous gland
- 6. Nerve, unmyelinated
- 7. Olfactory cells, nuclei
- 8. Pseudostratified epithelium
- 9. Skeletal muscle
- 10. Stratified squamous epithelium
- 11. Supporting cell, nucleus

Figure 15.40. Olfactory Epithelium, Chicken (Masson's). This pseudostratified columnar epithelium is similar to that found in mammals (see Figure 15.3). It is composed of basal, olfactory, and supporting cells.

Figure 15.41. Trachea, I.s., Chicken. Cartilaginous tracheal rings are complete and overlap each other. When the trachea is cut longitudinally, as in this preparation, the rings are cut transversely and present a lenticular profile.

Figure 15.42. Trachea, x.s., Chicken. Simple alveolar mucous glands occur in the ciliated, pseudostratified columnar epithelium. Portions of two overlapping tracheal rings are present. The inner ring was cut through its thin edge, while the outer one was cut through its thick middle region. See Figure 15.41 for an example of the shape of a tracheal ring that has been cut transversely.

Figure 15.43. Trachea, x.s., Chicken. Intraepithelial mucous glands are abundant in the trachea of the chicken. Also see Figure

Figure 15.44. Trachea, Near Syrinx, Chicken. The majority of the posterior, complete rings of the trachea shown here are bony. The ciliated pseudostratified columnar epithelium of the trachéa is followed in the syrinx by a stratified squamous epithelium.





Figure 15.45






Figure 15.47 ×125

KEY

- 1. Bony tracheal ring
- 2. Bronchial rings
- 3. Cartilaginous tracheal rings
- 4. External tympanic membrane
- 5. Intermediate syringeal cartilage

- 6. Internal tympanic membrane
- 8. Pessulus
- 7. Medial bronchial wall
- 9. Stratified squamous epithelium

Figure 15.45. Trachea, Syrinx, and Primary Bronchi, l.s., Chicken. As this drawing indicates, the syrinx is an upside-down-Y-shaped structure positioned between the trachea and the primary bronchi. Two pairs of thin membranes, the external and internal tympanic membranes, produce sound. The syrinx is supported by cartilages and a wedge-shaped cartilaginous or bony pessulus.

Figure 15.46. Syrinx, I.s., Chicken. The external and internal tympanic membranes of the syrinx, as well as the bony pessulus and an intermediate syringeal cartilage, are shown.

Figure 15.47. Syrinx, I.s., Chicken. A portion of the external tympanic membrane and intermediate syringeal cartilage.




Figure 15.48 ×62.5



Figure 15.49 ×125




Figure 15.50 ×12.5




Figure 15.51 ×25

| KEY | | |
|-------------------------------------|------------------------------------|--|
| 1. Atrium | 9. Pessulus | |
| 2. Bronchial cartilage | 10. Primary bronchus | |
| 3. Elastic fibers | 11. Pseudostratified epithelium | |
| 4. External tympanic membrane | 12. Secondary bronchus | |
| 5. Intermediate syringeal cartilage | 13. Smooth muscle | |
| 6. Internal tympanic membrane | 14. Stratified columnar epithelium | |
| 7. Lymphatic tissue | 15. Stratified squamous epithelium | |

16. Syrinx

Figure 15.48. Syrinx, I.s., Chicken (Orcein). Intermediate syringeal cartilages are connected by numerous elastic fibers.

8. Parabronchus

Figure 15.49. Syrinx, l.s., Chicken. Portion of pessulus and internal tympanic membrane. The latter is covered by both a ciliated, stratified columnar epithelium and by a stratified squamous epithelium.

Figure 15.50. Tympanic Membrane and Primary Bronchus, Chicken. Three transected bronchial cartilages. Bronchial cartilages are incomplete (C-shaped). They do not extend to the medial side of the bronchus.

Figure 15.51. Lung, Chicken. Longitudinal section through a secondary bronchus and parabronchi. The presence of numerous cup-shaped atria in the parabronchus distinguish this part of the bronchial tree from the secondary bronchus.




Figure 15.52 ×62.5




Figure 15.53




Figure 15.54 ×250




Figure 15.55 ×125

| KEY | | |
|-----------------------------|------------------------|--|
| 1. Air capillary | 8. Granulocyte | |
| 2. Atrium | 9. Macrophage | |
| 3. Bone | 10. Mucous cells | |
| 4. Cavity of air sac | 11. Parabronchus | |
| 5. Connective tissue lamina | 12. Secondary bronchus | |

13. Simple squamous epithelium

Figure 15.52. Lung, Chicken. Cross section of a parabronchus and a portion of an adjacent secondary bronchus.

7. Erythrocytes in vascular capillary 14. Smooth muscle

6. Epithelium

Figure 15.53. Lung, Chicken. Detail of the wall of a parabronchus. Note the continuity of the air capillaries with the atria. The latter are lined by an epithelium that varies from simple cuboidal to simple squamous.

Figure 15.54. Abdominal Air Sac, Chicken. The wall of the air sac consists of a connective tissue lamina and an epithelium that may be simple squamous, cuboidal, or ciliated columnar. Air sacs are not well vascularized.

Figure 15.55. Humerus, x.s., Chicken (Masson's). Many bones of the chicken contain extensions of air sacs.

ENDOCRINE SYSTEM

MAMMALS

The pituitary, pineal, thyroid, parathyroid, and adrenal glands possess certain features that distinguish them as organs of the endocrine system. They are very rich in wide, thin-walled vessels called sinusoids. The sinusoids are intimately associated with parenchymal cells, whose secretory products (hormones) pass directly into the circulatory system. Endocrine glands lack ducts. In contrast, exocrine glands convey their secretions (e.g., enzymes, mucus, and bile) through ducts to a mucosal or skin surface.

Endocrine cells are not limited to the glands presented in this chapter. For example, hormones are secreted by interstitial cells of the testes, corpora lutea and ovarian follicles, islets of Langerhans (pancreatic islets), and enterochromaffin cells of the gastrointestinal epithelium.

Pituitary Gland

The pituitary gland (hypophysis) is a major endocrine gland that is suspended from the hypothalamus of the brain. It releases several hormones, many of which influence the activity of other endocrine glands. The glandular portion, the adenohypophysis, forms from an outpocketing of the embryonic ectoderm of the dorsal portion of the oral cavity, called Rathke's pouch. The pars distalis, pars tuberalis, and pars intermedia constitute the adenohypophysis. The neural part of the pituitary gland, the neurohypophysis, is derived from a ventral outpocketing of the diencephalon. It is divisible into a median eminence, infundibular stalk, and pars nervosa.

The pars distalis is the largest portion of the pituitary gland. The parenchyma consists of irregular cords of cells separated by sinusoids and sparse connective tissue. There are two main types of parenchymal cells: chromophobes, characterized by a small

amount of cytoplasm that stains poorly, and chromophils, with more abundant cytoplasm that is readily stained. The chromophils are classified as acidophils (alpha cells) and basophils (beta cells). Basophils tend to be larger than acidophils. Chromophobes are smaller than chromophils and are most evident in groups, appearing as clusters of closely packed nuclei in tissue sections.

The pars intermedia is situated between the pars distalis and the pars nervosa. These regions are closely apposed in the horse. In other domestic mammals, the pars intermedia and pars distalis are partially separated by a small cleft, the hypophyseal cavity, which is the vestigial cavity of Rathke's pouch. The pars intermedia consists predominantly of basophilic cells. Follicles filled with colloid are often present.

The pars tuberalis is mainly located around the infundibular stalk. It is composed primarily of cords, clusters, and follicles of small, faintly basophilic cells.

The neurohypophysis contains numerous unmyelinated nerve fibers whose cell bodies are located in the supraoptic and paraventricular nuclei of the hypothalamus. Their axons converge at the median eminence (ventral boundary of the third ventricle) and form the hypothalamohypophyseal tract. They pass through the narrow infundibular stalk to the pars nervosa (infundibular process). The neurosecretions of these cells move along within the axons and accumulate at the terminal regions of the nerve fibers as Herring bodies, which are best demonstrated with special staining methods. Overall, the pars nervosa has an unorganized, fibrous appearance, and individual axons are indistinct. Numerous pituicytes (neuroglial cells) are scattered among the nerve fibers. They possess round to oval nuclei and long cytoplasmic processes. Their cytoplasm cannot be distinguished from nerve fibers in routine histologic preparations.

The infundibular cavity, which is continuous with the third ventricle and lined by ependymal cells, extends deep into the pars nervosa in the cat and pig and to a lesser extent in the dog and horse. The cavity does not reach beyond the infundibular stalk in ruminants. These relationships are evident in midsagittal sections of the pituitary gland.

Pineal Gland

The pineal gland (pineal body; epiphysis cerebri) is a dorsal evagination of the roof of the diencephalon. It is covered by connective tissue of the pia mater and divided into lobules by septa of connective tissue. The parenchyma is composed predominantly of pinealocytes, which are arranged as clusters, cords, or follicles. These epithelioid cells have a round nucleus and acidophilic cytoplasm. Neuroglial cells are also present.

Thyroid Gland

Each lobe of the **thyroid gland** is surrounded by a thin capsule of connective tissue and divided into lobules by

thin trabeculae. The latter are continuous with sparse intralobular connective tissue that contains numerous sinusoids. The connective tissue is abundant in the pig and cow. Each lobule consists of numerous follicles of various sizes that are frequently filled with colloid. The follicular cells vary in height, depending on the state of activity of the follicle. Their appearance changes from squamous or low cuboidal in the resting stage to cuboidal or columnar in the active stage. In an active follicle, the periphery of the colloid adjacent to the apical surface of the follicular cells is vacuolated. In an inactive follicle, the colloid has a smoother peripheral surface and vacuoles are not present.

Parafollicular (C) cells occur among the cells that line the thyroid follicles and also between the follicles. They are larger and have a paler cytoplasm than the follicular cells. Their nuclei are relatively large and pale. Parafollicular cells usually occur singly, but may also appear in groups. These cells are particularly abundant in the dog.

Parathyroid Glands

The parathyroid glands are classified as internal and external. Those that are adjacent to or embedded in the thyroid gland are the internal parathyroids. The external parathyroids lie a variable distance away from the thyroid gland. The parathyroid glands are surrounded by a thin capsule of connective tissue, which may be absent where the glands are deeply embedded within the thyroid gland. A stroma of connective tissue is well developed in the pig and cow, but is sparse in other domestic mammals.

The parenchyma of the parathyroid gland consists primarily of clusters and cords of principal (chief) cells. There are two different functional stages of the principal cell. The light principal cell is inactive and has a large, pale nucleus and pale, acidophilic cytoplasm. The dark principal cell is a smaller, active cell with a small, dark nucleus and a deeply acidophilic cytoplasm. In the sheep and goat, light cells tend to be located peripheral to the more central, dark cells. These cells are distributed randomly in the other domestic mammals.

Oxyphilic cells are large cells with an acidophilic cytoplasm and a pyknotic nucleus. They have been reported to occur in small numbers in the parathyroid glands of the horse and cow, particularly in older animals.

Adrenal Glands

The paired adrenal glands are situated close to the anterior end of the kidneys. The glands are covered by a capsule of dense irregular connective tissue that sometimes contains smooth muscle. Clusters of epithelioid cortical cells also occur in the capsule. Thin trabeculae project partially into the parenchyma.

Each adrenal gland is organized into a peripheral cortex and a central medulla. The adrenal cortex is

divided into four zones. The zona glomerulosa (zona multiformis) is the outermost zone. In the carnivore, horse, and pig, the parenchymal cells of this region are columnar and arranged into arcs. In the horse, the columnar cells are especially tall. In ruminants, the zona glomerulosa contains polyhedral cells that form irregular clusters or cords.

The zona intermedia lies between the zone glomerulosa and the zona fasciculata. It consists of small, closely packed cells. This zone is seen more often in the horse and carnivore than in the other domestic mammals.

The zona fasciculata, the widest zone of the adrenal cortex, is formed by radially arranged cords of cuboidal or polyhedral cells. The cords are one or two cells thick and separated by sinusoids. The cytoplasm of the cells in this zone frequently appears foamy because of the presence of numerous lipid vacuoles.

The zona reticularis is the innermost zone of the adrenal cortex. It is arranged as an irregular network of anastomosing cords of cells surrounded by sinusoids.

The adrenal medulla is composed mostly of columnar or polyhedral chromaffin cells, which form clusters and anastomosing cords separated by sinusoids. In domestic mammals, an outer and inner zone of the medulla can often be distinguished. The former consists of larger, more darkly stained cells, and the latter contains smaller, more lightly stained cells. Ganglion cells, either individually or in clusters, are scattered through the medulla. Because the cortex and medulla interdigitate at their junction, projections of the zona reticularis may appear within the medulla.

CHICKEN

Pituitary Gland

As in mammals, the pituitary gland (hypophysis) of the chicken is attached to the base of the brain below the diencephalon and is encapsulated by the dura mater. The adenohypophysis is composed of the pars distalis and pars tuberalis. A pars intermedia is absent. The pars distalis is divided into a cephalic region and a caudal region. Both regions contain cords of acidophils and basophils, and clusters of chromophobes. The acidophils of the cephalic region are pale, and those of the caudal zone are more darkly stained. Thus, the cephalic zone appears more basophilic, and the caudal zone appears more acidophilic. The cords of cells of the former are more closely packed than those of the latter. Some parenchymal cells of the pars distalis may be arranged around a lumen filled with colloid, especially in older birds. Cysts lined by ciliated cells and mucous cells also occur in this part of the pituitary gland.

The pars tuberalis surrounds the infundibulum and spreads dorsally over the ventral surface of the brain for a short distance. Ventrally, it extends to the posterior margin of the cephalic zone of the pars distalis. The pars tuberalis

contains small, round to elongated, slightly basophilic cells that are arranged in several layers.

The neurohypophysis includes the median eminence of the tuber cinereum, the infundibular stalk, and the pars nervosa (infundibular process). The median eminence and the infundibular stalk consist primarily of nerve fibers, neuroglial cells, and ependymal cells that line the infundibular cavity. The pars nervosa has an irregular surface and consists of numerous lobules. Each lobule contains a diverticulum of the infundibular cavity that is lined by ependymal cells. The latter are surrounded by irregular masses of tissue consisting of pituicytes and other neuroglial cells, nerve fibers, and Herring bodies.

Pineal Gland

The pineal gland (epiphysis cerebri) is a small, conical body that is situated between the cerebral hemispheres and the cerebellum. It is surrounded by connective tissue and is composed of a body and a narrow, ventral stalk that is attached to the roof of the third ventricle. The parenchyma of the gland is arranged into lobules separated by thin septa of connective tissue. The lobules contain cells, predominantly pinealocytes, that form rosettes or follicles.

Thyroid Glands

The **thyroid glands** are composed of numerous colloidfilled follicles, as in mammals. Cells that are similar in function to the parafollicular cells of mammals, however, occur in the **ultimobranchial bodies**, rather than the thyroid glands, of the chicken.

Parathyroid Glands

The parathyroid glands are each surrounded by a capsule of connective tissue. The parenchyma is composed of irregular cords of chief cells, separated by connective tissue and numerous sinusoids.

Adrenal Glands

The adrenal glands are enclosed within a capsule of dense connective tissue. Unlike mammals, the parenchyma is not organized into a distinct cortex and medulla. Instead, it is composed of intermingled cortical (interrenal) tissue and medullary (chromaffin) tissue. The cortical cells are arranged as irregular cords. These cells have dark nuclei and appear columnar when the cords are sectioned longitudinally. In a cross section of a cord, the cells appear tall and pyramidal with several cells arranged radially. Medullary tissue is composed of polygonal cells. They are

larger than cortical cells and possess a large, round nucleus and basophilic cytoplasm. Ganglion cells occur among the medullary cells. Two ganglia (the cranial and caudal suprarenal ganglia) are apposed to the surface of the adrenal glands and are frequently included in histologic sections of this gland.

| Word I | Roots | |
|----------------|----------------------|--|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Ad Ren | To, toward Kidney | The adrenal glands are near the kidneys. |
| Adeno | Gland | The <i>adenohypophysis</i> is the glandular portion of the pituitary gland. |
| Hypo Physis | Down Growth | The pituitary gland, which is also called the <i>hypophysis</i> , develops in part as a downgrowth from the brain. |
| Para | Beside, near | The <i>parathyroid</i> glands are beside the thyroid glands. |
| Pars | Part | The <i>pars</i> distalis is the distal part of the pituitary gland. |
| Pinea | A pine cone | The <i>pineal</i> gland resembles a small pine cone. |
| Pituita | Mucus or phlegm | The <i>pituitary</i> gland was thought to produce mucus that was expelled through the nostrils. |
| Rete | A net | The zona <i>reticularis</i> of the adrenal gland consists of an irregular network of cells. |
| Zona | A zone | The zona intermedia is an area between the zona glomerulosa and zona fasciculata. |






Figure 16.1 ×18



×62.5 Figure 16.2



×62.5 Figure 16.3

KEY 6. Median eminence 1. Cavity of Rathke's pouch 2. Ependymal cells 3. Follicle 7. Pars distalis 8. Pars intermedia 4. Infundibular cavity 9. Pars nervosa 5. Infundibular stalk 10. Pars tuberalis

Figure 16.1. Pituitary Gland, Cat. Parasagittal section showing all major components. In domestic mammals, except the horse, the cavity of Rathke's pouch persists in the adult. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 16.2. Pituitary Gland, Cat. Detail of the infundibular stalk and pars tuberalis. Note the presence of small follicles in the pars tuberalis lined by faintly basophilic epithelial cells. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 16.3. Pituitary Gland, Cat. Pars intermedia, pars distalis, and pars nervosa. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)




Figure 16.4 ×125




Figure 16.5




Figure 16.6 ×12.5




Figure 16.7 ×125

KEY

- 1. Acidophils
- 2. Blood vessel3. Cavity of Rathke's pouch
- 4. Chromophobes 5. Follicle


- 6. Infundibular cavity
- 7. Pars distalis
- 8. Pars intermedia
- 9. Pars nervosa
- 10. Pars tuberalis


Figure 16.4. Pituitary Gland, Dog. Detail of the pars distalis, pars intermedia, and pars nervosa.

Figure 16.5. Pituitary Gland, Parasagittal Section, Horse. Although present in other domestic animals, the cavity of Rathke's pouch is lacking in the horse.

Figure 16.6. Pituitary Gland, Horse (Alcian Blue, Orange G, Schiff's Reagent). Pars distalis, pars intermedia, and pars nervosa. Note the presence of follicles in the pars intermedia.

Figure 16.7. Pituitary Gland, Horse (Alcian Blue, Orange G, Schiff's Reagent). Detail of the pars intermedia and pars nervosa. The latter has a distinctive fibrous appearance.









Figure 16.10 ×25



| KEY | | |
|------------------------|-----------------------|--|
| 1. Acidophil | 5. Infundibular stalk | |
| 2. Basophil | 6. Pars distalis | |
| 3. Chromophobes | 7. Pars tuberalis | |
| 4. Infundibular cavity | 8. Sinusoid | |

Figure 16.8. Pituitary Gland, Horse (Alcian Blue, Orange G, Schiff's Reagent). Detail of the pars distalis. In this preparation, acidophils are orange, whereas basophils vary from blue to red. Chromophobes are small and pale.

Figure 16.9. Pituitary Gland, Horse. Detail of the pars distalis. Chromophobes appear in clusters and have closely spaced nuclei.

Figure 16.10. Pituitary Gland, Horse. A portion of the infundibular stalk, pars distalis, and pars tuberalis.




Figure 16.11 ×25




Figure 16.12 ×125




Figure 16.13 ×25




Figure 16.14 ×125




Figure 16.15 ×250

| | KEY |
|---|---|
| Adipose tissue Colloid Fibers of neuroglial cells Follicle Follicle, tangential cut Follicular cell | 7. Parafollicular cell 8. Parafollicular cells 9. Parathyroid gland 10. Pineal gland 11. Pineal stalk 12. Pinealocytes |

Figure 16.11. Pineal Gland, Dog. This gland consists primarily of pinealocytes and is located in the midline of the epithalamus.

Figure 16.12. Pineal Gland, Dog. Detail of the gland.

Figure 16.13. Thyroid and Parathyroid Glands, Dog. The basophilic, highly cellular parathyroid gland contrasts with the numerous colloid-filled follicles of the thyroid gland.

Figure 16.14. Thyroid Gland, Inactive, Dog. Parafollicular cells (C-cells) have pale-staining cytoplasm. In the dog they are particularly numerous and frequently occur in groups.

Figure 16.15. Thyroid Gland, Inactive, Dog. Large, pale-staining parafollicular cells often form a part of the cellular lining of a thyroid follicle.








Figure 16.17 ×25




Figure 16.18 ×125




Figure 16.19 ×250




Figure 16.20 ×62.5

| 1. Cilia 7. Follicular cell, pigmented 2. Colloid 8. Light cell 3. Cyst 9. Principal cells 4. Dark cell 10. Space artifact | | |
|--|-----------------------|---|
| 2. Colloid 8. Light cell 3. Cyst 9. Principal cells 4. Dark cell 10. Space artifact | | KEY |
| 5. Erythrocytes in sinusoid 11. Stroma | 2. Colloid 3. Cyst | 8. Light cell 9. Principal cells 10. Space artifact |

Figure 16.16. Thyroid Gland, Active, Horse. Active thyroid follicles are characterized by tall follicular cells and vacuolated colloid. Compare with Figures 16.15 and 16.17.

Figure 16.17. Thyroid Gland, Inactive, Goat. The high degree of vascularity of the thyroid gland is well illustrated in this micrograph. Pigment granules accumulate in the follicle cells of older animals.

Figure 16.18. Parathyroid Gland, Dog. Cysts containing colloid frequently occur in the parathyroid gland. Such cysts are lined by a ciliated columnar epithelium.

Figure 16.19. Parathyroid Gland, Dog. Both light and dark principal cells are visible. The active, dark cells have a nucleus with condensed chromatin and a dark, acidophilic cytoplasm. The inactive, light cells have a larger and paler nucleus and a lighter, acidophilic cytoplasm.

Figure 16.20. Parathyroid Gland, Cow. The stroma of connective tissue of the parathyroid gland of cows and pigs is abundant.




Figure 16.21 ×62.5




Figure 16.22 ×250




Figure 16.23 ×62.5




Figure 16.24 ×62.5




Figure 16.25 ×25

| | KEY |
|---|--|
| 1. Capsule 2. Epithelioid cells 3. Ganglion cell 4. Medulla 5. Medulla, inner region 6. Medulla, outer region | 7. Sinusoid 8. Zona fasciculata 9. Zona glomerulosa 10. Zona intermedia 11. Zona reticularis, cells of |


Figure 16.21. Adrenal Gland, Dog. Adrenal cortex and capsule. The cells of the zona glomerulosa are arranged into arclike formations in carnivores, horses, and pigs.

Figure 16.22. Adrenal Gland, Cat. Detail of a portion of the cortex. A zona intermedia occurs between the zona glomerulosa and zona fasciculata. It is especially well developed in carnivores and horses. It consists of small, polyhedral cells. Cells of the zona fasciculata are characteristically highly vacuolated.

Figure 16.23. Adrenal Gland, Horse. Adrenal cortex and capsule. The zona glomerulosa consists of high arcs composed of especially tall epithelial cells in the horse. Clusters of epithelioid cortical cells frequently occur in the capsule of the adrenal gland. A distinct intermediate zone separates the zona glomerulosa from the zona fasciculata.

Figure 16.24. Adrenal Gland, Horse. An autonomic ganglion, partly surrounded by cells extending from the zona reticularis, is situated in the medulla.

Figure 16.25. Adrenal Gland, Cow. Portions of the cortex and medulla. The medulla is subdivided into an outer region of darkly stained cells and an inner portion of lightly stained cells.








Figure 16.27 ×125




Figure 16.28 ×250




Figure 16.29 ×12.5




Figure 16.30 ×250

| KEY | | | | |
|---|---|--|--|--|
| 1. Capsule 2. Chromaffin cells 3. Chromophils 4. Chromophobes 5. Colloid 6. Cyst 7. Diencephalon 8. Ganglion cell 9. Infundibular cavity 10. Infundibular stalk 11. Medulla, inner region | 12. Medulla, outer region 13. Pars distalis, caudal 14. Pars distalis, cephalic 15. Pars nervosa 16. Pars tuberalis 17. Sinusoid, with erythrocytes 18. Skull 19. Zona fasciculata 20. Zona glomerulosa 21. Zona reticularis | | | |


Figure 16.26. Adrenal Gland, Cow. Portion of the adrenal cortex. The cells of the zona glomerulosa are arranged into irregular groups and cords in ruminants. Compare with Figures 16.21, 16.22, and 16.23.

Figure 16.27. Adrenal Gland, Cow. The basophilic cells of the outer region of the adrenal medulla contrast with the paler cells of the inner region.

Figure 16.28. Adrenal Gland, Sheep (Masson's). Adrenal medulla with ganglion cells and cells of the zona reticularis amid the chromaffin cells.

Figure 16.29. Pituitary Gland, Parasagittal Section, Chicken. In the chicken, the pars distalis is divisible into a cephalic zone and a caudal zone. The cephalic zone is more basophilic. Ciliated cysts commonly occur within the pars distalis. See Figure 16.32 for detail of a cyst.

Figure 16.30. Pituitary Gland, Chicken. The cephalic zone of the pars distalis consists of closely packed cords of chromophils and chromophobes. Some cords have a lumen filled with colloid.








Figure 16.32 ×250




Figure 16.33 ×250




Figure 16.34 ×125




Figure 16.35 ×25

| | KEY | |
|---|--|--|
| Acidophil Basophil Capsule Cerebellum Ciliated cell Granulocytes Infundibular stalk | 8. Mucous cell 9. Pars distalis, caudal 10. Pars tuberalis 11. Pineal gland 12. Pituicyte 13. Sinusoid 14. Skull | |
| | | |

Figure 16.31. Pituitary Gland, Chicken. In the caudal zone of the pars distalis, the cells of the cords are more loosely arranged than those of the cephalic zone. Acidophils have a more intensely stained cytoplasm than those of the cephalic zone, and they can be readily distinguished from basophils.

Figure 16.32. Pituitary Gland, Chicken. Portion of a cyst, in the pars distalis, lined by ciliated cells and mucous cells.

Figure 16.33. Pituitary Gland, Chicken. Portions of the pars tuberalis and adjacent infundibular stalk. The cells of the pars tuberalis are rounded to elongated with a finely granular, slightly basophilic cytoplasm and a round to oval nucleus.

Figure 16.34. Pituitary Gland, Chicken. A group of pituicytes within the pars nervosa. Pituicytes have a clear cytoplasm and a large, vesicular nucleus.

Figure 16.35. Pineal Gland, Parasagittal Section, Chicken. The body of the pineal gland, portion of the overlying skull, and the cerebellum.




Figure 16.36 ×125




Figure 16.37 ×250




Figure 16.38 $\times 62.5$




Figure 16.39 ×250

- 1. Adipose tissue
- Capsule
 Chief cells
- 4. Cortical cells 5. Follicle

- 6. Ganglion7. Medullary cells
- 8. Rosette
- 9. Sinusoid

Figure 16.36. Pineal Gland, Chicken. The parenchymal cells of the pineal gland are arranged as compact masses (rosettes) or as round to oval follicles with distinct lumens.

Figure 16.37. Parathyroid Gland, Chicken. This gland consists of chief cells arranged into a feltwork of anastomosing cords. The cords are surrounded by strands of connective tissue and numerous sinusoids.

Figure 16.38. Adrenal Gland, Chicken. Cords of cortical cells are interwoven between clumps and irregular masses of medullary cells throughout the gland.

Figure 16.39. Adrenal Gland, Chicken. Detail of cortical and medullary cells. Cortical cells are columnar. When longitudinal cuts have been made through cords of cortical cells, the cells form a bilayer. When cords are cut transversely, the cells are seen to be arranged radially. Medullary cells are polygonal, are larger than cortical cells, and possess basophilic cytoplasm. They are arranged as clumps or irregular masses.

MALE REPRODUCTIVE SYSTEM

MAMMALS

he male reproductive system includes the testes, the system of ducts that leads from them, the penis, and accessory glands.

Testes

The testes are contained in the scrotum and are compound tubular glands that are invested by a thick capsule of dense irregular connective tissue, the tunica albuginea. This capsule is rich in smooth muscle in the stallion. The tunica albuginea is covered by a peritoneum, the visceral layer of the tunica vaginalis. The latter is composed of a mesothelium and underlying connective tissue that blends with that of the tunica albuginea. Septa of connective tissue extend from the tunica albuginea into the testis, partially or completely dividing the testis into lobules. These septa are thin in ruminants and thicker in the carnivore, stallion, and boar. Centrally, the septa may merge with the loose connective tissue of the mediastinum testis.

Within each lobule of the testis there are convoluted seminiferous tubules. These are lined by a stratified epithelium of spermatogenic cells and Sertoli cells. The spermatogenic cells give rise to spermatozoa. Spermatogonia, the most immature spermatogenic cells, are small, round cells with dark, round nuclei that lie adjacent to the basement membrane. These undergo mitotic divisions and produce primary spermatocytes, larger cells whose nuclei often show distinct chromatin. Primary spermatocytes undergo the first meiotic division, giving rise to smaller, secondary spermatocytes. Secondary spermatocytes are rarely observed because they undergo the second meiotic division shortly after they arise, forming haploid spermatids. Early spermatids are round cells with pale nuclei that occur in clusters toward the lumen of the seminiferous tubule. Late spermatids are

characterized by small, oval to elongated, dark heads, and long, faint tails that project into the lumen. They are eventually released from the seminiferous epithelium as spermatozoa.

Various combinations of developing spermatogenic cells occur within the epithelium of a seminiferous tubule. These cell associations (stages) are unique and occupy only a portion of the length of each tubule. The total number of different stages varies with different animals; for example, rats are known to have 14 different stages, mice have 12, and pigs have eight.

The cell mix within each stage can be observed while examining a histologic section of the testis. In sections through some seminiferous tubules, for example, spermatogonia, two layers of primary spermatocytes, and numerous early spermatids may be apparent; in another segment there may be spermatogonia, a single layer of primary spermatocytes, and numerous early and late spermatids. Other spermatogenic cell combinations, characteristic of the animal, become apparent as more tubules are examined.

Sertoli cells (sustentacular cells) are fewer in number than the spermatogenic cells. They are distinguished by a pale oval or triangular nucleus that has a prominent nucleolus and occasional cleftlike infoldings. They are tall cells that extend from the basement membrane to the lumen of the tubule, but their boundaries are indistinct in routine histologic preparations. Numerous lateral and apical invaginations of their cell membranes embrace the differentiating spermatogenic cells.

Flattened, contractile myoid cells lie just outside the basement membrane of each seminiferous tubule. The connective tissue between adjacent tubules contains polyhedral interstitial (Leydig) cells. These produce testosterone and are particularly abundant in the stallion and boar. They are recognized by their small, round nucleus and an acidophilic, often foamy cytoplasm.

Near the terminal segment of a seminiferous tubule, the spermatogenic cells decrease in number and the Sertoli cells become more numerous. A **transitional zone**, lined by Sertoli cells, joins a seminiferous tubule to a **straight tubule**. The latter may be lined by simple columnar, cuboidal, or squamous cells and is continuous with a network of anastomosing channels that form the **rete testis**. The rete testis possesses a simple squamous or cuboidal epithelium that may be bistratified cuboidal in the bull. It is surrounded by the loose connective tissue of the **mediastinum testis**.

Efferent Ductules and Duct of the Epididymis

Efferent ductules, lined by a simple columnar or a pseudostratified epithelium with some ciliated cells, lead from the rete testis and pass through the tunica albuginea to join the duct of the epididymis in the head of the epididymis. In the stallion, the tubules of the rete testis penetrate the tunica albuginea and form an extratesticular rete testis, which is then joined to the duct of the epididymis by efferent ductules. The coiled duct of the epididymis varies in structure from the head to the tail region of the epi-

didymis. Its pseudostratified columnar epithelium, with stereocilia, is thickest in the head region and is encircled by some smooth muscle. In the body (mid) region there is less smooth muscle, and the epithelium is thinner. In the tail region of the epididymis, the pseudostratified epithelium is thinnest and the surrounding smooth muscle is most abundant. In the stallion, the lining of the duct in the tail region of the epididymis forms short, villuslike projections.

Vas Deferens

The vas deferens (ductus deferens) leads from the duct of the epididymis and joins with the urethra. The vas deferens is lined by a pseudostratified columnar epithelium (some cells with stereocilia) that may become simple columnar distally. The smooth muscle of its thick muscularis presents a variety of arrangements. It may form an inner circular and an outer longitudinal layer, and each of these layers may contain interwoven fibers of smooth muscle. In contrast, the entire muscularis may be interwoven with no distinct layers of smooth muscle. No particular arrangement has been observed to be consistent within a species.

Accessory Glands

The male accessory glands include the glands of the ampulla, seminal vesicles, the bulbourethral glands, and the prostate gland. They are composed of branched tubular or tubuloacinar secretory units that often have vesicular dilations. The secretory epithelium of these glands is classified as pseudostratified because, although consisting primarily of columnar cells (or sometimes cuboidal cells such as in the prostate), occasional basal cells are present.

Near its junction with the urethra, the vas deferens forms a dilated **ampulla** whose lamina propria and submucosa are filled with glandular secretory units. The ampulla is absent in the tomcat, and the ampullary glands are not well developed in the boar.

The prostate gland is a seromucous gland except in the dog, where it is entirely serous. In the boar and ruminants, the prostate gland consists mostly of a disseminate portion (pars disseminata) in the form of a glandular layer in the submucosa of the pelvic urethra. In the stallion and carnivores, the disseminate portion is represented only by scattered glands. The body of the prostate gland is well developed in the stallion and carnivore and is absent in the ram and billy goat (buck). It is an encapsulated, lobulated gland that partially or completely surrounds a part of the pelvic urethra.

The seminal vesicles (vesicular glands) are absent in carnivores. In the stallion they are true vesicular outpocketings in the form of bladderlike sacs with wide central lumens into which the glands open. In the boar and ruminants they are compact glands with a lobulated surface.

The mucous-secreting **bulbourethral** (Cowper's) **glands** are present in all domestic mammals except the dog. The columnar cells of the pseudostratified epithelium are tall and pale and possess basally displaced nuclei.

Urethra

The male urethra, which carries both urine and semen, can be divided into a pelvic and a penile portion. The pelvic urethra is lined by a transitional epithelium, which may become stratified columnar or cuboidal distally. Along the entire length of the urethra, the connective tissue below the mucosa contains erectile tissue with thin-walled cavernous spaces (veins). In the pelvic urethra this erectile tissue forms the stratum cavernosum (vascular stratum). Peripheral to this stratum are the glands of the disseminate portion of the prostate gland. The muscularis of the urethra near the bladder consists of an inner and outer longitudinal layer and a middle circular layer of smooth muscle. In the vicinity of the prostate gland, most of the smooth muscle is replaced by skeletal urethral muscle. However, some longitudinal smooth muscle remains. The muscularis of the pelvic urethra is surrounded by an adventitia.

The penile urethra, which courses through the ventral region of the penis, is lined by a mixture of transitional, stratified cuboidal, stratified columnar, or simple columnar epithelium. The larger, more abundant cavernous spaces of the penile urethra form the corpus spongiosum (corpus cavernosum urethra), which is surrounded by a tunica albuginea. Except for an occasional cell, the wall of the penile urethra lacks a muscularis of smooth muscle.

In the stallion and ruminants, the terminal portion of the urethra extends beyond the penis, forming a **urethral process**. It is covered by a cutaneous membrane and lined by transitional or stratified squamous epithelium. In the stallion the urethral process contains well-developed erectile tissue. In the ram and billy goat the urethral process contains small cavernous spaces and two cords of fibrocartilage that parallel the urethra.

Penis

The penis can be divided into the body and glans penis. Both regions contain the penile urethra with its erectile tissue, the corpus spongiosum. The body of the penis (corpus penis) is characterized by two additional masses of erectile tissue called the corpora cavernosa. Each corpus cavernosum is enclosed by the dense connective tissue and elastic fibers of the tunica albuginea. The tunic is especially thick in the boar and ruminants and contains smooth muscle in the stallion. It extends inward to form a network of trabeculae between which lies the spongy erectile tissue. The latter contains cavernous spaces, lined by endothelium and surrounded by various proportions of smooth muscle and fibroelastic connective tissue. The smooth muscle is predominant in the vascular penis of the stallion. In the fibroelastic penis of the boar and ruminants, the cavernous spaces are surrounded mainly by connective tissue that is rich in elastic fibers and contains little or no smooth muscle. In the intermediate type of penis of the carnivore, both smooth muscle and connective tissue fill the spaces between the cavernous vessels. The corpus cavernosum of all domestic mammals contains scattered adipose tissue in the connective tissue between the cavernous vessels. This is abundant in the tomcat, especially toward the tip of the

corpus cavernosum, where adipose tissue nearly replaces the erectile tissue.

The cavernous spaces receive their blood supply from groups of helicine arteries. The walls of these tortuous vessels have cushionlike thickenings, formed from longitudinal bundles of smooth muscle, epithelioid cells, and abundant elastic tissue.

The expanded, distal portion of the penis, called the glans penis, is best developed in the stallion and dog. It contains erectile tissue, which is continuous with that of the corpus spongiosum. In carnivores the glans contains an os penis. This bone is small in the tomcat. It is well developed in the dog and possesses a fibrocartilaginous tip. The surface (visceral prepuce) of the glans penis of the tomcat bears small, keratinized epidermal spines. Small epidermal projections also occur in the stallion and billy goat.

Prepuce

The prepuce is a tubelike reflection of skin that covers the distal, free portion of the penis. It is composed of an external, parietal, and visceral layer. The external layer is typical skin and is continuous with the abdominal skin. The external layer turns inward at the preputial opening to form the parietal prepuce (internal layer). This, in turn, reflects at the fornix and continues onto the end of the penis as the visceral prepuce. The stallion has an additional outer fold called the sheath. Hair, sweat glands, and sebaceous glands occur over a variable distance from the external layer to the parietal prepuce. Glands may occur occasionally in the visceral prepuce of the stallion.

CHICKEN

Testes

The testes are situated in the abdominal cavity of the rooster. They are surrounded by a thin connective tissue capsule, the tunica albuginea, which is covered by a peritoneum. There are no well-developed septa to divide the testes into lobules.

The epithelial cells of the convoluted seminiferous tubules are like those of mammals: Sertoli cells, spermatogonia, primary spermatocytes, secondary spermatocytes, spermatids, and spermatozoa. Unlike those in mammals, various cell associations do not occur in segments along the length of the seminiferous tubules. Instead, the seminiferous epithelium is arranged into narrow columns of cells that undergo spermatogenesis independently. There is very little connective tissue between adjacent seminiferous tubules, and interstitial cells are sparse. They occur singly or in small clusters, primarily in the larger interstitial spaces. They are flattened to polyhedral cells with a relatively large, round nucleus and cytoplasm that is often vacuolated.

The seminiferous tubules are continuous with straight tubules, which are lined by Sertoli cells. Straight tubules lead into the anastomosing channels of the rete testis,

which is lined by a simple cuboidal to squamous epithelium. The rete testis lies outside the tunica albuginea below the epididymis.

Epididymis

Three types of tubules occur within the epididymis: efferent ductules, connecting ducts, and the duct of the epididymis. The numerous, convoluted efferent ductules join the rete testis to the connecting ducts. They are lined by a simple epithelium of intermittent groups of tall and low columnar cells, as well as patches of cells that appear to be pseudostratified. The epithelial cells are arranged into folds, and many of the cells bear tufts of cilia. The connecting ducts (excretory canals) are smaller in diameter than the other tubules of the epididymis and are lined by a pseudostratified columnar epithelium. The epithelial cells are rarely ciliated and are not arranged into numerous folds, as are the cells that line the efferent ductules. Thus, the luminal surface of the connecting ducts has a smooth appearance. The single, convoluted duct of the epididymis is similar in structure to the connecting ducts, except that it is much larger in diameter. All of the tubules of the epididymis are surrounded and bound by connective tissue.

Vas Deferens and Ejaculatory Duct

The duct of the epididymis joins the vas deferens at the terminal portion of the epididymis. The latter is a convoluted duct with a pseudostratified columnar epithelium,

underlying smooth muscle, and a more peripheral layer of dense connective tissue. Each vas deferens merges with a small, conical ejaculatory duct, whose submucosa contains erectile tissue. The ejaculatory duct protrudes and opens into the urodeum of the cloaca, marking the termination of the duct system of the male.

| Word Roots | | |
|--------------------|---|---|
| | | |
| ROOT | MEANING | EXAMPLE SENTENCE |
| Corp Spongiosum | A body Like a sponge | Having numerous cavernous spaces, the corpus spongiosum is a spongelike mass. |
| Epi Didym | Upon Testis | The epididymis is on the testis. |
| Inter Sistum | Between Set | Interstitial cells are situated in the interstitial tissue between seminiferous tubules of the testes. |
| Муо | Muscle | Myoid cells are contractile, like muscle cells. |
| Rete | A net | The <i>rete testis</i> consists of a network of channels in the testis. |
| Semin Fer | Seed, sperm Carry | Sperm are produced in the seminiferous tubules of the testes. |
| Sperm Genesis | Seed, male reproductive cells Origin, birth | Spermatogenic cells give rise to sperm. |
| Tunic Vagin | A covering or cloak A sheath | The tunica albuginea is covered by the <i>tunica</i> vaginalis. |
| | | |






Figure 17.1 ×18



| | KEY |
|---|---|
| Gonocyte Interstitial cells Rete testis | Sex cord Straight tubule Supporting cells |

Figure 17.1. Testis, x.s., Baby Boar. Developing sex cords in the testis of a two-day-old boar.

Figure 17.2. Testis, x.s., Baby Boar. Detail of developing sex cords and interstitial cells. Two types of cells can be distinguished in the sex cord. Supporting cells (small with dark nuclei) are positioned along the edges of the cords. They will develop into Sertoli cells. Gonocytes, precursors of spermatogonia, are located in the interior of the cords. They have large, pale nuclei.






Figure 17.4




Figure 17.5 \times 360

KEY

- 1. Interstitial cells
- 2. Lumen
- 3. Myoid cell, nucleus
- 4. Primary spermatocyte
- 5. Seminiferous tubulé
- 6. Sertoli cell, nucleus
- 7. Spermatid, early
- 8. Spermatid, late 9. Spermatogonia
- 10. Tunica albuginea
- 11. Tunica albuginea, smooth muscle
- 12. Tunica albuginea, vascular layer
- 13. Tunica vaginalis

Figure 17.3. Testis, Stallion (Mallory's). The tunica albuginea of the stallion is characterized by the presence of smooth muscle.

Figure 17.4. Testis, Boar (Mallory's). The tunica albuginea consists of dense irregular connective tissue. It lacks smooth muscle in domestic mammals, except the stallion.

Figure 17.5. Seminiferous Tubules, Testis, Dog. A portion of each of three adjacent seminiferous tubules is shown.

Summary of cells of the seminiferous tubule

Spermatogenic cells

Spermatogonia: Small, round cells with dark, round nuclei that lie on the basement membrane

Primary spermatocytes: Larger cells whose large nuclei often show distinct chromatin


Secondary spermatocytes: Smaller and closer to the lumen than primary spermatocytes; rarely seen because they undergo the second meiotic division soon after they form


Early spermatids: Small, round cells with pale nuclei that occur in clusters near the lumen of the seminiferous tubule


Late spermatids: Small, oval to elongated dark heads, tucked in the apical surface of Sertoli cells, with long tails that project into the lumen of the tubule

Spermatozoa: May be present in the lumen of the seminiferous tubule; they have elongated dark heads and long tails

Sertoli cells (sustentacular cells): Tall cells with poorly defined outlines, as they have numerous apical and lateral pockets that envelop the spermatogenic cells; have an elongated, sometimes triangular, nucleus with fine chromatin and a prominent nucleolus












Figure 17.8 ×125




Figure 17.9 $\times 12.5$

KEY

- Efferent ductule
- 2. Interstitial cell
- 3. Mediastinum testis
- 4. Myoid cell, nucleus
- 5. Primary spermatocyte6. Rete testis, channel
- 7. Seminiferous tubules
- 8. Sertoli cell, nucleus
- 9. Sertoli cells 10. Spermatid, early 11. Spermatid, late
- 12. Spermatogonium13. Straight tubule
- 14. Transitional zone
- 15. Tunica albuginea
- Figure 17.6. Seminiferous Tubules, Testis, Stallion. Portions of four seminiferous tubules are visible. Note the numerous intersti-

tial cells (abundant in the boar and stallion) and the section through a straight tubule. Figure 17.7. Interstitial Tissue, Testis, Ram. Interstitial tissue and

portions of three seminiferous tubules are shown. Interstitial cells

are relatively sparse in carnivores and ruminants.

Figure 17.8. Transitional Zone and Straight Tubule, Testis, Stallion. A transitional zone joins a seminiferous tubule to a straight tubule. Sertoli cells line this zone and protrude into the lumen of the straight tubule.

Figure 17.9. Rete Testis, Stallion. Anastomosing channels of the rete testis lie within the loose connective tissue of the mediastinum testis. In the stallion, the rete testis extends through the tunica albuginea and becomes extratesticular, as in this micrograph. Junctions of rete channels and efferent ductules can be seen. See Figure 17.10 for a magnified view of a junction.

Synonyms:

Interstitial cells = Leydig cells Sertoli cells = sustentacular cells




Figure 17.10 ×250




Figure 17.11 ×25




Figure 17.12 ×250




Figure 17.13 ×25




Figure 17.14 ×62.5

| | KEY |
|--|--|
| Columnar epithelium Cuboidal epithelium, rete testis Duct of the epididymis Efferent ductule Loose connective tissue Lymphocyte, migrating | Pseudostratified epithelium Pseudostratified epithelium, efferent ductule Smooth muscle Spermatozoa Tunica albuginea, smooth muscle Tunica vaginalis, mesothelium |
| | |


Figure 17.10. Junction of Rete Testis and Efferent Ductule, Stallion. The rete testis is lined by cuboidal cells, whereas the efferent ductule is lined by a ciliated, pseudostratified columnar epithelium.

Figure 17.11. Efferent Ductules, Stallion. Various cuts through the tortuous efferent ductules are surrounded by loose connective tissue.

Figure 17.12. Efferent Ductules, Stallion. Efferent ductules are lined by a ciliated, pseudostratified columnar epithelium. However, the epithelium may be simple columnar in some places.

Figure 17.13. Head of Epididymis, Stallion (Masson's). The epididymis is surrounded by a tunica albuginea of dense, irregular connective tissue, which contains smooth muscle in the stallion. Portions of the coiled duct of the epididymis are shown.

Figure 17.14. Head of Epididymis, Stallion. In this region the pseudostratified columnar epithelium of the duct of the epididymis is thickest. Smooth muscle is scarce. Compare with Figures 17.15 and 17.16.








Figure 17.16 ×62.5




Figure 17.17 ×250




Figure 17.18 ×25




Figure 17.19 ×62.5

- 1. Basal cell
- 2. Lamina propria
- 3. Loose connective tissue
- 4. Muscularis
- 5. Pseudostratified epithelium
- 6. Serosa
- 7. Smooth muscle
- 8. Spermatozoa
- 9. Stereocilia
- 10. Villus-like projection

Figure 17.15. Body of Epididymis, Stallion. The duct of the epididymis in this region is surrounded by more smooth muscle than in the head of the epididymis, and the pseudostratified columnar epithelium is not as thick as in the head of the epididymis (see Figure 17.14).

Figure 17.16. Tail of Epididymis, Stallion. A low, pseudostratified columnar epithelium and abundant circular smooth muscle characterize the duct of the epididymis in this region. In the stallion, the caudal segment of the duct of the epididymis has villuslike projections.

Figure 17.17. Tail of Epididymis, Stallion. Detail of the duct of the epididymis lined by low, pseudostratified columnar epithelium and surrounded by abundant, circular smooth muscle.

Figure 17.18. Vas Deferens, x.s., Dog. The bulk of the wall consists of smooth muscle, which forms an inner circular and an outer predominantly longitudinal layer with some randomly arranged cells.

Figure 17.19. Vas Deferens, x.s., Stallion (Masson's). This section of the vas deferens, taken from near the epididymis, has long mucosal folds. The inner layer of the muscularis contains interwoven bundles of smooth muscle. Although out of the field of view in this micrograph, the smooth muscle of the outer layer of the muscularis is mostly arranged longitudinally.




Figure 17.20 ×62.5



Figure 17.21 ×12.5




Figure 17.22 ×62.5




Figure 17.23 ×125




Figure 17.24 ×25

KEY

- 1. Adventitia
- 2. Ampulla, lumen
- 3. Basal cell, nucleus
- 4. Gland
- 5. Interlobular septum
- 6. Lamina propria
- 7. Muscularis
- 8. Pseudostratified epithelium
- 9. Secretion
- 10. Serosa
- 11. Spermatozoa

Figure 17.20. Vas Deferens, x.s., Boar (Masson's). The muscularis consists of an admixture of longitudinally and randomly arranged smooth muscle. The epithelium is pseudostratified columnar with stereocilia present intermittently.

Figure 17.21. Ampulla, x.s., Ram. The terminal segment of the vas deferens, the ampulla, contains branched tubuloalveolar glands in the dog, stallion, and ruminants. The glands are poorly developed in the boar. The ampulla is absent in the tomcat.


Figure 17.22. Ampulla, Ram. Detail of the mucosa. Spermatozoa are stored in the glands close to their openings into the lumen of the ampulla.

Figure 17.23. Ampulla, Ram. The secretory alveoli are lined by a pseudostratified epithelium composed of cuboidal to columnar cells and occasional basal cells.


Figure 17.24. Seminal Vesicle, Castrated Billy Goat. In the castrated male, the glandular tissue of the accessory glands is greatly reduced. Compare with Figure 17.25.



Figure 17.25 ×25



×250 **Figure 17.26**



×12.5 **Figure 17.27**




Figure 17.28 $\times 62.5$




Figure 17.29 $\times 12.5$

KEY

- 1. Basal cell
- Capsule
 Duct
- 4. Gland
- 5. Interlobular septum
- 6. Prostate gland
- 7. Pseudostratified epithelium
- 8. Secretion
- 9. Stratum cavernosum
- 10. Trabecula
- 11. Transitional epithelium
- 12. Urethra, lumen


Figure 17.25. Seminal Vesicle, Ram. Lobules of tubuloalveolar glands are divided by interlobular septa, which contain an abundance of smooth muscle in ruminants. In the stallion and boar, the septa consist predominantly of connective tissue with some smooth muscle. Seminal vesicles are absent in carnivores.

Figure 17.26. Seminal Vesicle, Ram. The pseudostratified glandular epithelium is characterized by sparse basal cells. Note the muscular septum.

Figure 17.27. Body of the Prostate, Dog. The body of the prostate, which is well developed in carnivores and the stallion, is surrounded by a capsule of dense connective tissue and smooth muscle. Trabeculae from the capsule divide the gland into lobules.

Figure 17.28. Body of the Prostate, Dog. In the dog, this is a serous gland. Compare with Figure 17.31.

Figure 17.29. Disseminate Portion of the Prostate, x.s., Ram (Masson's). This portion of the prostate is well developed in the boar and ruminants. The glands are located within the submucosa of the pelvic urethra. The stratum cavernosum of the pelvic urethra contains cavernous spaces that are smaller and less numerous than those of the corpus spongiosum of the penile urethra.






Figure 17.31 ×125



Figure 17.32 ×3a




Figure 17.33 ×12.5

Figure 17.34 ×62.5

| KEY | | |
|----------------------|-----------------------------|--|
| 1. Cavernous space | 7. Secretory cells | |
| 2. Corpus spongiosum | 8. Serous cells | |
| 3. Duct | 9. Smooth muscle | |
| 4. Lamina propria | 10. Spongy bone | |
| 5. Mucous cells | 11. Transitional epithelium | |
| 6. Os penis | 12. Urethra, lumen | |

Figure 17.30. Disseminate Portion of the Prostate, x.s., Ram (Masson's). The transitional epithelium of both the pelvic urethra and a duct of the prostate gland is shown.

Figure 17.31. Disseminate Portion of the Prostate, x.s., Ram (Masson's). The prostate is a mixed gland except in the dog, where there are no mucous secretory units.

Figure 17.32. Bulbourethral Gland, Boar. This gland is a compound tubular gland in the boar, tomcat, and billy goat. It is a tubuloalveolar gland in the stallion, bull, and ram. It is absent in the dog. The pale-staining secretory cells are columnar or pyramidal and have basally displaced nuclei.

Figure 17.33. Penis, x.s., Puppy. Section is through the developing os penis, which is present in carnivores.

Figure 17.34. Penis, x.s., Puppy. Detail of the urethra and portion of the os penis.




Figure 17.35 ×12.5



Figure 17.36 ×62.5




Figure 17.37 ×12.5




Figure 17.38 ×62.5




Figure 17.39 ×12:

- 1. Cavernous space
- 2. Connective tissue
- 3. Corpus cavernosum
- 4. Corpus spongiosum
- 5. Dermis
- 6. Endothelium
- 7. Epidermal spine
- 8. Lamina propria
- 9. Os penis
- 10. Smooth muscle
- 11. Stratified columnar epithelium
- 12. Tunica albuginea
- 13. Urethra, lumen


Figure 17.35. Penis, I.s., Tomcat. In the tomcat, the distal portion of the corpus cavernosum consists largely of nonerectile, adipose tissue. A small os penis is present in the glans, and small spines are present on the surface of the glans of the tomcat. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 17.36. Glans Penis, I.s., Tomcat. Detail of a keratinized epidermal spine. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 17.37. Penile Urethra, x.s., Stallion. Portion of the penile urethra with abundant cavernous spaces of the corpus spongiosum.

Figure 17.38. Corpus Spongiosum, Body of Penis, Stallion. The cavernous spaces of the stallion and carnivore are surrounded by connective tissue rich in elastic fibers and by many bundles of smooth muscle.

Figure 17.39. Penile Urethra, x.s., Stallion. The epithelial lining of the urethra in this section is stratified columnar. The epithelium, however, is variable in the penile urethra and in places may also be simple columnar, transitional, or stratified cuboidal.








Figure 17.41 ×12.5




Figure 17.42




Figure 17.43 ×62.5




Figure 17.44 ×25

- 1. Adipose tissue
- 2. Cavernous space
- 3. Connective tissue
- 4. Connective tissue trabecula
- 5. Corpus cavernosum
- 6. Elastic fibers
- 7. Retractor penis muscle 8. Smooth muscle

- 9. Tunica albuginea 10. Urethra, lumen


Figure 17.40. Body of Penis, Stallion. Large masses of smooth muscle surround the cavernous spaces of the corpus cavernosum of the stallion.


Figure 17.41. Body of Penis, x.s., Boar. A portion of the sigmoid flexure, including the retractor penis muscle.


Figure 17.42. Body of Penis, x.s., Boar (Orcein). The connective tissue surrounding the cavernous spaces of the corpus cavernosum is rich in elastic fibers in the boar and ruminants.

Figure 17.43. Body of Penis, Boar. In the boar and ruminant, the cavernous spaces of the corpus cavernosum are invested largely by connective tissue and only a smattering of smooth


Figure 17.44. Penile Urethra, x.s., Ram. The distribution of the cavernous spaces of the corpus spongiosum of the penile urethra is especially well represented in this section.








×25 **Figure 17.46**



×62.5 **Figure 17.47**



Figure 17.48 $\times 62.5$



×12.5 **Figure 17.49**

- Cavernous space
 Elastic fibers
 Epidermis

- 4. Fibrocartilaginous cord
- 5. Helicine artery
- 6. Lamina propria
- 7. Smegma

- 8. Smooth muscle
- 9. Stratified squamous epithelium
- 10. Trabecula
- 11. Transitional epithelium
- 12. Tunica albuginea
- 13. Urethra, lumen


Figure 17.45. Body of Penis, x.s., Ram (Orcein). The cavernous spaces of the corpus spongiosum are surrounded by connective tissue rich in elastic fibers in boars and ruminants.

Figure 17.46. Helicine Artery, Body of Penis, Ram (Orcein). The corpus cavernosum contains helicine arteries, which are tortuous vessels with an abundance of elastic fibers throughout their walls.

Figure 17.47. Helicine Artery, Body of Penis, Ram (Masson's). Junction of a helicine artery with a cavernous space in the corpus

Figure 17.48. Urethral Process, x.s., Ram. The urethral process is a tortuous, wormlike extension of the urethra in the ram and billy goat. One of the two fibrocartilaginous cords that parallel the urethra is shown.

Figure 17.49. Urethral Pouch, Stallion. The urethral pouch, found only in the stallion, is filled with smegma, which is composed of desquamated epithelial cells and the secretion of the preputial glands.








Figure 17.51 ×25




Figure 17.52 ×18




Figure 17.53 ×250


| KEY | | | | |
|---|---|--|--|--|
| 1. Connecting duct 2. Dermal papilla 3. Efferent ductule 4. Hair follicle 5. Interstitial cell 6. Lymphatic nodule 7. Parietal prepuce 8. Preputial gland 9. Primary spermatocyte | 10. Rete testis 11. Sebaceous gland 12. Seminiferous tubule 13. Sertoli cell, nucleus 14. Spermatid, early 15. Spermatid, late 16. Spermatogonium 17. Stratified squamous epithelium 18. Visceral prepuce | | | |

Figure 17.50. Parietal Prepuce, Stallion. The dermis contains sebaceous glands and tubular preputial (sweat) glands.

Figure 17.51. Prepuce, Boar. The parietal and visceral prepuce are shown.

Figure 17.52. Testis and Epididymis, x.s., Rooster. Seminiferous tubules, the rete testis, and portions (efferent ductules and connecting ducts) of the epididymis.

Figure 17.53. Interstitial Tissue, Testis, Rooster. Interstitial (Leydig) cells are found principally in the larger intertubular spaces. These cells are either polyhedral or elongated and may contain vacuoles.








Figure 17.56




Figure 17.57

- Cilia
 Efferent ductule
- 3. Primary spermatocyte4. Rete testis5. Seminiferous tubule

- 6. Sertoli cell, nucleus
- 7. Spermatid, early
- 8. Spermatid, late 9. Spermatogonium
- 10. Spermatogonium, dividing
- 11. Spermatozoa 12. Straight tubule
- 13. Tunica albuginea

Figure 17.54. Seminiferous Tubules, Testis, Rooster. Detail of portions of adjacent seminiferous tubules. Note that the seminiferous epithelial cells are organized into narrow columns.

Figure 17.55. Testis, Rooster. A straight tubule, lined by Sertoli cells, connects a seminiferous tubule with the rete testis.

Figure 17.56. Straight Tubule, Testis, Rooster. Sertoli cells form the epithelium of straight tubules.

Figure 17.57. Junction of Rete Testis and Efferent Ductule, Rooster. The epithelial cells lining efferent ductules vary in shape and many possess cilia. The rete testis is lined by squamous epithelial cells.




Figure 17.58 ×250




Figure 17.59 ×250




Figure 17.60




Figure 17.61 ×125

- 1. Cilia
- Connecting duct
 Duct of the epididymis
- 4. Efferent ductule
- 5. Pseudostratified epithelium
- 6. Smooth muscle
- 7. Spermatozoa

Figure 17.58. Efferent Ductule, x.s., Rooster. Detail of an efferent ductule. The epithelial cells vary in shape, and many bear cilia. The epithelium is folded and is surrounded by loose connective tissue. Occasionally, smooth muscle may be present.

Figure 17.59. Connecting Duct and Efferent Ductule, Rooster. Connecting ducts present a smooth inner surface and are lined by a pseudostratified columnar epithelium with occasional basal cells. The epithelial cells are generally without cilia. In contrast, the efferent ductule has a folded lining, and its epithelial cells are mostly ciliated.

Figure 17.60. Duct of the Epididymis, Rooster. The duct of the epididymis has a larger diameter than a connecting duct, but otherwise is comparable in structure with the latter.

Figure 17.61. Vas Deferens, Distal, Rooster. The lining epithelium is similar to that of the epididymis. A layer of smooth muscle separates the epithelium from the surrounding connective tissue.

FEMALE REPRODUCTIVE SYSTEM

MAMMALS

he ovaries, oviducts, uterus, vagina, and vulva are the major components of the mammalian female reproductive system.

Ovaries

A simple squamous or cuboidal epithelium, germinal epithelium, often missing from histologic preparations, covers the cortex of the ovary. Beneath the epithelium is a layer of dense connective tissue, the tunica albuginea. A cortical stroma, containing ovarian follicles in various stages of development, lies internal to the tunica albuginea. In bitches and queens, but ordinarily not in other domestic mammals, cords of epithelioid cells called interstitial glands occur throughout the stroma. The epithelioid cells are derived from the theca interna of atretic, antral follicles or from the granulosa cells of atretic, preantral follicles.

A medulla consisting of richly vascularized loose connective tissue lies internal to the ovarian cortex. In the mare, the medullary tissue is located external to the cortex. Channels, lined by a cuboidal epithelium and called the rete ovarii, are conspicuous components of the medulla in carnivores and ruminants. Hilus cells (groups of epithelioid cells) may be found close to the rete ovarii in the region of the hilus in some mammals.

Primordial follicles are the least developed and most numerous follicles of the ovary. They lie just below the tunica albuginea. Each consists of a **primary oocyte** surrounded by a layer of simple squamous **follicle cells**. In response to periodic hormonal stimulation, growth is initiated in some of the primordial follicles. The earliest growing follicle,

the primary follicle, consists of an enlarging oocyte surrounded by a layer of cuboidal cells. Proliferation of the follicle cells results in the formation of a multilaminar (late primary) follicle. Fluid-filled spaces appearing between the follicle cells gradually coalesce, forming an antrum. Concomitantly, an acidophilic, translucent membrane, the zona pellucida, appears around the oocyte. Further growth results in the formation of a secondary follicle with a C-shaped antrum. Its follicle cells are now called the membrana granulosa. A sheath of stromal cells, the theca folliculi, forms around the follicle. The theca differentiates into a cellular, vascular inner layer, the theca interna, and an outer layer of connective tissue, the theca externa. The boundary between the stroma and the theca externa and that between the theca externa and theca interna is often indistinct. Continued growth results in the formation of a large tertiary (Graafian) follicle whose oocyte is surrounded by a multilayer of membrana granulosa cells, the cumulus oophorus. The columnar cells of the innermost portion of the latter constitute the corona radiata, which is separated from the oocyte by the zona pellucida.

Ordinarily, each mature tertiary follicle contains a single oocyte. However, the follicles of certain animals (carnivores, sows, and ewes) may contain as many as six oocytes.

Mature follicles vary widely in size. They are about 2 mm in diameter in the bitch and queen, 15 mm in the cow, and as large as 70 mm in the mare. Maximum size is reached just prior to ovulation. Following ovulation, granulosa cells and cells of the theca interna of most species multiply, hypertrophy, and differentiate into granulosa lutein cells and smaller, more peripheral theca lutein cells, respectively, of the corpus luteum. A yellow pigment (lutein) is formed by the luteal cells of the cow, mare, and carnivores, but is lacking in ewes, nanny goats (does), and sows. Luteal cells produce progesterone. Regression of the corpus luteum occurs during late diestrus, leaving scar tissue, the corpus albicans.

Although many primordial follicles begin the process of growth and differentiation, few become mature. The majority undergo a degenerative regression, called atresia. The oocyte and membrane granulosa degenerate first. Cells of the theca interna hypertrophy, and the zona pellucida becomes swollen. Eventually, the entire follicle is resorbed.

Oviduct

The **oviduct** is a muscular tube consisting of an **isthmus**, which arises from the uterus, a middle segment, the **ampulla**, and a funnel-shaped **infundibulum**, which lies next to the ovary. From the outside inward, the wall of the oviduct is comprised of a serosa, muscularis, lamina propria, and epithelium. The muscularis, which is thickest in the isthmus, is formed mainly from circular smooth muscle with a modicum of longitudinally arranged smooth muscle external to it. Many of the epithelial cells lining the cavity of the oviduct are ciliated. In part, the epithelium of ruminants and sows is pseudostratified. The mucosa is thrown into longitudinal folds, with less folding occurring in the isthmus than in the ampulla.

Uterus

The wall of the bicornuate uterus of domestic mammals has three layers: the outer perimetrium (serosa), middle myometrium, and inner endometrium (mucosa). The myometrium is divisible into a thick, inner circular layer and a thin, outer longitudinal layer. A richly vascularized and well-innervated stratum vasculare usually separates the muscle layers. The stratum vasculare, however, is indistinct in the sow and may be located in the outer half of the circular layer in the cow.

The epithelium of the endometrium is simple cuboidal or columnar in the bitch, queen, and mare, but may be stratified or pseudostratified in ruminants and sows. Simple, branched **uterine** (endometrial) **glands** extend into the lamina propria. These may be considerably coiled in the mare, sow, and ruminants. Nonglandular regions of the endometrium, called **caruncles**, occur in ruminants.

The mucosa of the uterine **cervix** is elevated into longitudinal folds that may become subdivided into secondary and tertiary folds. The epithelial lining is simple columnar with goblet cells. In the bitch, however, it is stratified squamous. Glandular tissue fades in the cervix, extending to the cervical os only in carnivores. An inner circular and an outer longitudinal layer of smooth muscle form the muscularis.

Estrous Cycle

The estrous cycle consists of a succession of stages. The first stage, proestrus, is characterized by endometrial growth. It is followed by estrus, or the period when the female is receptive to the male. In most species, ovulation occurs during estrus. The development of the corpus luteum occurs during the next stage, metestrus. Diestrus follows metestrus and coincides with the presence of a fully functional corpus luteum. The development and secretory activity of the endometrial glands peak during this time. Anestrus, a period of sexual inactivity, follows diestrus.

Placenta

The placenta is derived from the endometrium and the chorioallantoic membrane (CAM). The degree of intimacy between these two components varies and is a basis for classifying placentas. A placenta is **indeciduate** when these two membranes are in contact but are not intimately fused. The placenta is **deciduate** when the membranes have become fused. Little or no endometrium is lost during the birth process in animals having an indeciduate placenta (mare, ruminants, and sow). Conversely, considerable mucosa is lost at parturition in animals with deciduate placentas (carnivores).

The extent to which the CAM contributes to the placenta varies. If most of the CAM contributes, as in the mare and sow, the placenta is **diffuse**; if numerous but isolated areas contribute, as in ruminants, the placenta is **cotyledonary**; when a beltlike portion of the CAM contributes, as in carnivores, the placenta is **zonary**.

The surfaces of the chorioallantoic membrane and the endometrium may contact one another in three different ways. These types of contact are designated as folded, villous, and labyrinthine. In the sow, both surfaces are folded and are closely applied to each other. In the mare and ruminants, chorioallantoic villi insert into pockets (crypts) in the endometrium. In carnivores, the apposed surfaces form a complex, interlinked, fused labyrinth.

Classification of the placenta can also be based on the number of tissue layers separating the fetal and maternal blood. In the mare and sow, six layers intervene: the endothelium, connective tissue, and epithelium of the CAM; and the epithelium, connective tissue, and endothelium of the endometrium. This configuration characterizes the epitheliochorial placenta. In ewes and nanny goats (does), the epithelium of the caruncles (endometrial elevations where functional contact with the CAM is made) is lost, thereby reducing the number of tissue layers to five (syndesmochorial placenta). In the cow, the epithelium of the caruncle remains intact (epitheliochorial), but portions of the intercaruncular epithelium degenerate. In carnivores, both the endometrial epithelium and the endometrial connective tissue are lost, bringing the epithelium of the CAM and the endothelium of the endometrium into contact. Only four tissue layers separate the fetal and maternal blood in this type, the endotheliochorial placenta.

Vagina

A mucosa, muscularis, and adventitia or serosa (cranial region only) form the wall of the vagina. The mucosa is lined, throughout, by a stratified squamous epithelium in all species except the cow. In the anterior portion of the vagina of the cow, the epithelium is stratified columnar with goblet cells. In carnivores, the epithelial cells become keratinized during estrus. A lamina propria and submucosa are present. Usually, the inner layer of the muscularis is thick and consists of circularly arranged smooth muscle, while the outer layer is thin and consists of longitudinally organized smooth muscle. In some animals (bitch, queen, and sow), a thin layer of longitudinal muscle occurs internal to the circular layer. An adventitia or serosa is present.

Vaginal Cytology

Vaginal cytology provides a way of determining stages of the estrous cycle of the bitch or queen and therefore can be helpful to the practitioner who is trying to determine the best time to breed an animal. In the bitch, for example, proestrus, estrus, diestrus, and anestrus are stages of the estrous cycle. The formation of the corpus luteum occurs during late estrus in the bitch. Therefore, there is no metestrus. Proestrus lasts an average of nine days and is characterized by a watery, bloody discharge and swollen vulva. Estrus is evidenced when a bitch is willing to stand for mating, and ordinarily lasts about nine days. A clear or bloody discharge is present. Diestrus lasts for about two

months and begins on the day when the bitch no longer tolerates a male's advances. Anestrus follows diestrus and may last from two to 10 months.

Various types of epithelial cells are found in vaginal smears taken during the estrous cycle. Parabasal cells are the smallest. They are round cells with round nuclei, and have the highest nucleocytoplasmic ratio of any of the sloughed cells. Intermediate cells are larger than parabasal cells. Their nuclei are similar in size and shape to those of the latter. The corners of intermediate cells are rounded. Superficial intermediate cells (transitional cells) are bigger than intermediate cells and have angular edges. Their nuclei resemble those of parabasal and intermediate cells. Superficial cells are similar in size to superficial intermediate cells. Their edges are angular and may be folded. Their nuclei are pyknotic, faded, or lacking.

Smears taken during proestrus (early to mid) may contain erythrocytes and neutrophils as well as parabasal, intermediate, superficial intermediate, and superficial cells. During late proestrus, superficial intermediate and superficial cells are the most numerous, and neutrophils decline.

The vast majority (90% or more) of cells found in smears taken during estrus are superficial cells. Similar smears may be obtained during late proestrus. Ordinarily, during estrus, neutrophils are not observed. Erythrocytes show a reduction in number, but in many bitches they can be found throughout estrus and into early diestrus. Bacteria may be found in estrous smears.

During diestrus, superficial cells decrease by a minimum of 20%. Parabasal and intermediate cells, which may have been absent or very sparse, increase to more than 10% and frequently rise to more than 50%. Although neutrophils reappear during diestrus, smears from some bitches contain few or none. Because erythrocytes may be present in smears from early diestrus, it is not possible to distinguish proestrus from diestrus without taking more than one smear.

During anestrus, parabasal and intermediate cells predominate in smears. Bacteria may be found, but will be less abundant than in proestrus or estrus. Neutrophils may occur, but are ordinarily less abundant than in early diestrus.

Vulva

In domestic mammals, the vulva includes the vestibule, labia, and clitoris. The mucosal epithelium is stratified squamous. The major vestibular glands are bilateral, mucus-secreting, tubuloacinar glands in the submucosa, found in ruminants and the queen. Minor vestibular glands occur in the mucosa of most domestic animals. They are small, branched, tubular, mucous glands distributed through the vestibular mucosa.

The integument of the labia (lips of the vulva) has a structure like that of the external skin. It is well endowed with both sebaceous and tubular apocrine glands.

The clitoris consists of **erectile tissue** (corpus cavernosum clitoridis), a **glans**, and a **prepuce**. The amount of erectile tissue varies. The prepuce has parietal and visceral components as in males.

CHICKEN

Ovary

The left **ovary** and oviduct represent the reproductive organs of the hen. The ovary consists of an outer **cortex** that envelops a vascular **medulla**. Ovarian follicles of various sizes occur within the cortex. A layer (germinal epithelium) of cuboidal or flattened cells covers the cortex. The **tunica albuginea**, composed of dense connective tissue, lies below the epithelium. A **stroma** of loose connective tissue occurs below the tunica albuginea.

Developing follicles occur throughout the stroma of the cortex. Large follicles are suspended from the surface of the ovary by stalks of cortical tissue. Each follicle consists of a growing, yolk-laden oocyte with a rounded nucleus (germinal vesicle). The oocyte is surrounded by several layers. These are, from the outside inward, the theca externa, theca interna, membrana granulosa, and perivitelline membrane. The latter abuts the surface membrane of the oocyte.

The theca externa is formed from a compact connective tissue that contains groups of pale interstitial (luteal) cells. The latter may also be found, in groups, in the cortical stroma and medulla. The theca interna is only about one quarter as thick as the externa. It is formed from a compact layer of spindle-shaped cells. The membrana granulosa consists of a single layer of cuboidal cells in the smallest and largest follicles, but in those of intermediate size, the epithelium is pseudostratified columnar.

The cortex of the mature ovary also contains concentrations of fat-filled **vacuolar cells**. Numerous fat vacuoles occur throughout the cytoplasm of these cells, and their nuclei are pyknotic. Collections of these cells are believed to represent regressing postovulatory follicles.

Atretic follicles are commonly found in normal active ovaries. In the most common type of atretic follicle, cells of the membrana granulosa proliferate, forming a number of irregular layers around the oocyte. The oocyte becomes smaller and is eventually replaced by granulosa cells. Ultimately, scar tissue replaces the granulosa cells. In older birds, the oocyte becomes surrounded by hyperplastic and hypertrophied interstitial (luteal) cells during atresia. Both the oocyte and the cells of the membrana granulosa eventually degenerate.

Oviduct

The oviduct of the chicken is tortuous and muscular. It consists, in anterioposterior sequence, of the following five regions: infundibulum, magnum, isthmus, uterus (shell gland), and vagina. From the outside inward, the wall of the oviduct consists of a serosa, muscularis (outer longitudinal and inner circular smooth muscle), lamina propria, and epithelium. In most regions, the lamina propria contains glands.

The infundibulum is composed of a thin-walled funnel and a neck region. Scattered bundles of smooth muscle lie within the connective tissue between the serosa and ciliated, simple columnar epithelium. Longitudinal folds are present in the mucosa within the interior of the funnel near the neck. The folds increase in depth within the neck, and

secondary folds appear. The muscularis becomes sorted out into circular and longitudinal layers in the neck.


The magnum is the longest part of the oviduct. Its well developed tubular glands produce albumin. Its mucosal folds are more numerous and taller than those of the infundibulum. Tertiary folds are present. The muscularis is better developed than in the infundibulum. The pseudostratified epithelium is composed of ciliated columnar cells and secretory (goblet) cells.

The **isthmus** is a relatively short region with a diameter less than that of the magnum. Its longitudinal mucosal folds possess numerous secondary folds. The muscularis is better developed than the magnum's. The epithelium is ciliated, pseudostratified columnar with secretory cells. Its numerous tubular glands secrete the shell membranes.

The uterus (shell gland) is an expanded portion of the oviduct. Its walls are not as thick as those of the preceding segments. The muscularis is well developed, especially the longitudinal layer. The mucosa is thrown into longitudinal, leaf-shaped folds that are covered by a ciliated, pseudostratified, columnar epithelium. The shell of the egg is produced from secretions of its tubular glands.

The vagina is a short, narrow duct. Its muscularis is well developed, especially the circular layer. Its mucosa is thrown into numerous tall, narrow folds bearing many small secondary folds. The surface is covered by a ciliated, pseudostratified columnar epithelium with mucous cells. Sperm storage occurs in the sperm-host glands. These tubular glands occur within the connective tissue of the mucosa of the vagina near the junction between the uterus and vagina. After insemination, sperm appear in compact masses within the glands. The vagina of the oviduct opens into the urodeum of the cloaca.

| Word Roots | | |
|-----------------------------|-------------------------------|---|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Corona | A crown | The cells of the <i>corona</i> radiate enclose the oocyte. |
| Corpus Lute | Body Yellowish | A yellow pigment forms in the cells of the <i>corpus luteum</i> in some mammals. |
| Infundibulum | A funnel | The <i>infundibulum</i> is the funnel shaped portion of the oviduct. |
| Isthm | A narrow passage | The <i>isthmus</i> is the relatively short and narrow part of the oviduct of the chicken. |
| Magna | Great or large | The <i>magnum</i> is the longest part of the oviduct of the chicken. |
| Metri | Uterus | The <i>myometrium</i> is the middle, muscular layer of the wall of the uterus. |
| Multi Lamina Follicle | Many Layer A little bag | A <i>multilaminar follicle</i> in the ovary has many layers of cells that surround the oocyte. |
| Oo Cyte | An egg Cell | An <i>oocyte</i> is an immature egg cell. |
| Ovi | Egg | The <i>oviduct</i> is a passageway for eggs. |
| Pellucid | Transparent | The zona <i>pellucida</i> is a transparent membrane that develops around the oocyte in the ovary. |
| Stratum Vas | Layer A vessel | The stratum vasculare is a richly vascularized layer within the myometrium. |
| Theca | A capsule | A sheath of cells, the <i>theca</i> folliculi, forms ground the ovarian follicle |






Figure 18.2 ×125




Figure 18.3 ×12




Figure 18.4 ×62.5


| KEY | | | |
|---|--|--|--|
| 1. Antrum | 11. Oocyte, nucleus | | |
| 2. Corona radiata | 12. Primary follicle 13. Primordial follicles | | |
| 3. Corpus luteum | 13. Primordial follicles | | |
| 4. Cortex | 14. Stroma | | |
| Cumulus oophorus Granulosa cells | 15. Theca externa | | |
| | 16. Theca folliculi | | |
| 7. Growing follicles | 17. Theca interna | | |
| 8. Medulla | 18. Tunica albuginea | | |
| 9. Membrana granulosa | 19. Zona pellucida | | |
| 10. Oocyte, cytoplasm | ' | | |


Figure 18.1. Ovary, Queen. Follicles of various ages and a corpus luteum can be seen in the cortex. A portion of the vascular medulla is present.

Figure 18.2. Ovary, Queen. Early follicles in the outer region of the cortex.

Figure 18.3. Ovary, Bitch. A multilaminar, primary follicle.

Figure 18.4. Ovary, Queen. A young, tertiary follicle.








Figure 18.7 ×62.5




Figure 18.8 ×25


| KEY | | | |
|---|---|--|--|
| Antrum Corpus albicans Germinal epithelium Granulosa lutein cell Hypertrophied theca cells Interstitial gland | 7. Membrana granulosa 8. Theca externa 9. Theca interna 10. Theca lutein cell 11. Young tertiary follicle 12. Zona pellucida | | |


Figure 18.5. Ovary, Bitch. Portion of the wall of a tertiary follicle, and part of an adjacent corpus luteum.

Figure 18.6. Corpus Luteum, Ovary, Sow. Peripheral region of a corpus luteum showing theca lutein cells (small) and granulose lutein cells (large).

Figure 18.7. Ovary, Bitch. Atretic follicles, each with a swollen zona pellucida.

Figure 18.8. Corpus Albicans, Ovary, Cow (Masson's). The scar tissue of the corpus albicans is stained bright blue-green in this preparation.








Figure 18.10 ×250




Figure 18.11 ×62.5



Figure 18.12 ×62.5




Figure 18.13 ×25

| KEY | | | |
|---|---|--|--|
| Corpus luteum Folds Granulosa lutein cell | 4. Hilus cells 5. Interstitial gland 6. Rete ovarii | | |


Figure 18.9. Ovary, Bitch. Several interstitial glands are visible within the stroma between two corpora lutea.

Figure 18.10. Interstitial Glands, Ovary, Bitch. Cords of epithelioid cells form the parenchyma of interstitial glands. These glands are well developed in queens and bitches.

Figure 18.11. Hilus Cells, Ovary, Cow. Clusters of epithelioid cells, located in the vicinity of the hilus, are called hilus cells. They resemble the epithelioid cells of the interstitial glands (see Figure 18.10).

Figure 18.12. Rete Ovarii, Ovary, Cow. Cords of cells, or channels lined by cuboidal epithelial cells, located in the medulla of the ovary are called the rete ovarii. They are considered to be homologous to the rete testis.

Figure 18.13. Fimbria of Infundibulum, Oviduct, Mare. The mucosa of the fimbria is highly folded.








Figure 18.15 ×125




Figure 18.16 ×250



Figure 18.17 ×12.5




Figure 18.18 ×12.5

| | KEY | |
|--|---|--|
| Adipose tissue, mesosalpinx Cilia Columnar epithelium, ciliated Extruded nucleus Folds | 6. Lamina propria7. Muscularis8. Secretory bleb9. Serosa10. Submucosa | |

Figure 18.14. Fimbria of Infundibulum, Oviduct, Mare. Detail of the wall. Note the smooth muscle of the thin muscularis.

Figure 18.15. Fimbria of Infundibulum, Oviduct, Cow. Portions of the fimbria may lack smooth muscle, as in this example.

Figure 18.16. Infundibulum, Oviduct, Cow (Masson's). The epithelium consists of ciliated, columnar epithelial cells and nonciliated, secretory cells. Extruded nuclei, which appear to arise from epithelial cells, are common.

Figure 18.17. Infundibulum, Neck, Oviduct, x.s., Cow. The mucosa is highly folded, and the muscularis is thin.

Figure 18.18. Ampulla, Oviduct, x.s., Cow. The mucosa is highly folded. The muscularis is relatively thick. Compare with Figure 18.17.




Figure 18.19



Figure 18.20 ×25




Figure 18.21 $\times 250$




Figure 18.22 ×12.5



Figure 18.23 ×180

- 1. Circular muscle
- Columnar epithelium
 Cuboidal epithelium
- 4. Endometrial gland

- 6. Longitudinal muscle
- 7. Myometrium, circular 8. Myometrium, longitudinal
- 9. Stratum vasculare

Figure 18.19. Isthmus, Oviduct, x.s., Mare. The mucosa of the isthmus has fewer folds than any other part of the oviduct. The muscularis is thickest in this part of the oviduct.

Figure 18.20. Uterine Horn, x.s., Anestrus, Bitch. The endometrium is thin and the glands are sparse in anestrus.

Figure 18.21. Uterine Horn, x.s., Anestrus, Queen. The lumen of the anestrus uterus is lined by a simple cuboidal epithelium.

Figure 18.22. Uterine Horn, x.s., Proestrus, Bitch. In proestrus, the endometrium becomes thicker and the glands enlarge.

Figure 18.23. Uterine Horn, x.s., Proestrus, Bitch. Luminal epithelial cells become columnar during proestrus and estrus.




Figure 18.24 ×12.5




Figure 18.25 ×12.5




Figure 18.26 ×12.5




Figure 18.27 ×25




Figure 18.28 ×250

- 1. Artery
- 2. Endometrial gland
- 3. Mesometrium
- 4. Myometrium
- 5. Myometrium, circular
- Myometrium, longitudinal
- 7. Stratum vasculare
- 8. Tunica adventitia
- 9. Tunica intima
- 10. Tunica media
- 11. Vein

Figure 18.24. Uterine Horn, x.s., Estrus, Bitch. A thick endometrium and highly developed glands are characteristic of the estrous uterus.

Figure 18.25. Uterine Horn, x.s., Diestrus, Bitch. The endometrium and its glands become fully developed during diestrus.

Figure 18.26. Mesometrium, Bitch. The mesometrium contains an abundance of smooth muscle and numerous blood vessels. Smooth muscle of the mesometrium is continuous with the outer, longitudinal layer of the myometrium.

Figure 18.27. Uterine Horn, Brood Mare (Orcein). There is an abundance of elastic fibers (red-brown in this micrograph) in the intima and adventitia of blood vessels of the uterus of animals that have been through a pregnancy. The section is from the midregion of the myometrium.

Figure 18.28. Uterine Horn, Brood Mare. Detail of a portion of an artery in the myometrium. The intima becomes thickened with elastic fibers and smooth muscle in animals that have experienced a pregnancy. The adventitia also becomes heavily infiltrated with elastic fibers.




Figure 18.29 ×25





Figure 18.30 ×125





×12.5 **Figure 18.32**




Figure 18.33

- 1. Blood vessels
- 2. Caruncle
- 3. Cilia
- 4. Cuboidal epithelium
- 5. Endometrial gland
- 6. Hemorrhagic region
- 7. Myometrium, circular
- 8. Myometrium, longitudinal
- 9. Pseudostratified epithelium

Figure 18.29. Uterine Horn, x.s., Metestrus, Cow. Metestrous bleeding occurs in the cow. Numerous erythrocytes of hemorrhagic regions can be seen beneath the surface epithelium (see Figure 18.30).

Figure 18.30. Uterus, Metestrus, Cow. Detail of Figure 18.29. Hemorrhagic regions are evident in the endometrium beneath the surface epithelium. The epithelial cells are cuboidal during met estrus in the cow.

Figure 18.31. Uterine Horn, Cow (Masson's). The epithelial cells lining the uterine glands are sometimes ciliated, as in this section.

Figure 18.32. Caruncle, Uterus, x.s., Cow. The endometrium of the uterus of ruminants contains nonglandular, highly cellular prominences called caruncles. Uterine glands that lie deep to the caruncle open near its base.

Figure 18.33. Caruncle, Uterus, Cow. The caruncle consists of highly cellular (mostly fibroblasts) connective tissue and numerous blood vessels located beneath the epithelium.




Figure 18.34




Figure 18.35 ×25




Figure 18.36 ×62.5

- 1. Allantoic blood vessel
- 2. Allantoic epithelium
- 3. Chorioallantoic membrane
- 4. Chorion laeve
- 5. Deep glandular layer
- 6. Marginal hematoma
- 7. Myometrium
- 8. Placental labyrinth
- 9. Spongy layer
- 10. Supraglandular layer

Figure 18.34. Placenta (Zonary and Endotheliochorial), Bitch (Drawing). Carnivores have a zonary placenta that appears, grossly, as a beltlike band around the middle of the chorionic sac. The chorionic (fetal) tissue penetrates to the endothelium of the maternal blood vessels. A placenta with this type of fetal-maternal junction is called an endotheliochorial placenta. The association between the maternal endothelium and syntrophoblast can be seen in Figures 18.37 and 18.39.

Note: This drawing is of a section through a portion of one of the edges of the zonary placenta of a bitch.

Figure 18.35. Placenta (Zonary and Endotheliochorial), Bitch (Trichrome). Section through the deepest layers of the placenta. See Figure 18.34 for location. The spongy layer is formed by the occluded uterine glands in the mid-region of the endometrium. The deep glandular layer consists of the bases of the uterine glands. The supraglandular layer is a sheet of connective tissue between the deep glandular and spongy layers. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.36. Placenta (Zonary and Endotheliochorial), Bitch (Trichrome). Portion of the chorioallantoic membrane and the placental labyrinth. See Figure. 18.34 for location. The chorioallantoic membrane in this micrograph appears thicker than normal because of the presence of extensive space artifact. Note the presence of fetal blood vessels in the chorioallantoic membrane. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)




Figure 18.37 ×12:




Figure 18.38 ×250




Figure 18.39 ×62




Figure 18.40 ×125


- 1. Allantoic blood vessel
- 2. Chorioallantoic connective tissue
- 3. Cytotrophoblast
- 4. Eroded endometrial epithelium
- 5. Maternal blood vessel
- 6. Necrotic endometrial tissue
- 7. Syntrophoblast
- 8. Trophoblastic projection, vacuolated cells

Figure 18.37. Placenta (Zonary and Endotheliochorial), Bitch (Trichrome). Section is through the placental labyrinth. Both maternal and fetal components of this endotheliochorial placenta can be seen. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.38. Placenta (Zonary and Endotheliochorial), Bitch (Trichrome). Detail of the placental labyrinth. The maternal blood vessels are lined by endothelial cells with bulging nuclei. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.39. Placenta (Zonary and Endotheliochorial), Bitch (Trichrome). Trophoblastic projections, lined by large, pale, vacuolated cells, protrude into spaces (areolae) in the region where the placental labyrinth is forming. A portion of the maternal tissue, which has been partially destroyed by the invading trophoblast, is represented by the red-stained, necrotic tissue seen in the lower left quadrant of the micrograph. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, lowa State University.)

Figure 18.40. Placenta (Zonary and Endotheliochorial), Bitch (Trichrome). Detail of Figure 18.39. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)








Figure 18.42 ×62.5




Figure 18.43 ×25




Figure 18.44 ×125


- 1. Allantoic blood vessel
- 2. Chorioallantoic membrane
- 3. Chorioallantoic villus
- 4. Chorioallantoic villus, epithelium
- 5. Chorion laeve, epithelium
- 6. Crypt
- 7. Crypt, epithelium
- 8. Marginal hematoma
- 9. Maternal blood vessel
- 10. Microplacentome
- 11. Placental labyrinth
- 12. Spongy zone
- 13. Uterine gland
- 14. Uterus, epithelium

Figure 18.41. Placenta (Zonary and Endotheliochorial), Bitch (Trichrome). A portion of a marginal hematoma, consisting of large compartments filled with blood derived from hemorrhaging uterine blood vessels, is shown. See Figure 18.34 for location. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)


Figure 18.42. Chorion Laeve, Bitch (Trichrome). The chorion laeve is the part of the chorioallantoic sac that is not involved in the formation of the placenta. Its surface is smooth and is apposed to the uterine epithelium. This section is from the region adjacent to the hematoma of the placenta. See Figure 18.34 for location. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, lowa State University.)

Figure 18.43. Placenta (Diffuse and Epitheliochorial), Mare. In the horse, small tufts of branched chorioallantoic villi interdigitate with crypts of the endometrium. Together, the tufts and the crypts form structures called microplacentomes. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.44. Placenta (Diffuse and Epitheliochorial), Mare. Detail of a microplacentome. Longitudinal and cross sections of chorioallantoic villi are surrounded by endometrial crypts. The epithelium of the crypts, which may vary in height, is flattened in this region. The villi contain blood vessels and connective tissue and are covered by pink-stained trophoblast cells. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)









 $\times 180$ **Figure 18.47**




Figure 18.48 $\times 12.5$

- 1. Allantoic blood vessel
- Chorioallantoic membrane
- Chorioallantoic membrane, primary fold
- 4. Chorioallantoic membrane, secondary fold
- Chorioallantoic villus, epithelium 17. Space artifact
 Crypt, epithelium 18. Stalk of placentome
- 6. Crypt, epithelium7. Cuboidal cells
- 8. Endometrium, connective tissue
- 9. Endometrium, primary fold
- 10. Endometrium, secondary fold

- 11. High columnar cells
- 12. Intercotyledonary endometrium
- 13. Low columnar cells
- 14. Maternal blood vessel
- 15. Myometrium
- 16. Placentome

- 19. Uterine gland

Figure 18.45. Placenta (Diffuse and Epitheliochorial), Mare. Detail of a placentome adjacent to the endometrium. The epithe-

lium of the endometrial crypt consists of pale, cuboidal cells. The epithelium of the chorioallantoic villi is formed from pink-stained cuboidal and low columnar cells. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.46. Placenta (Diffuse and Epitheliochorial), Sow (Trichrome). The placenta of the sow is folded, diffuse, and epitheliochorial. Folds of the chorioallantoic membrane interdigitate with folds of the uterus. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.47. Placenta (Diffuse and Epitheliochorial), Sow (Trichrome). Interdigitating secondary folds of the chorioallantoic membrane and endometrium. The bases of the folds of the chorioallantoic membrane are lined by high columnar epithelial cells, whereas the crests of the maternal folds are covered by shorter columnar cells. The remainder of both epithelial surfaces is lined by cuboidal or flattened cells. Note that the chorioallantoic epithelial surface is invaded by capillaries. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.48. Placenta (Cotyledonary and Epitheliochorial), Cow. A section through a placentome formed from the association of a cotyledon (clumps of chorioallantoic villi) with a uterine caruncle (endometrial elevation). Note that the endometrial epithelium of the intercotyledonary region is discontinuous.




Figure 18.49




Figure 18.50





Figure 18.51



×12.5




Figure 18.53

| KEY | | | |
|--|---|--|--|
| Allantoic blood vessel Chorioallantoic villus, branch Chorioallantoic villus, main stem Cilia Crypt Cryptal epithelium | 7. Diplokaryocyte 8. Lamina propria 9. Primary fold 10. Secondary fold 11. Stratified squamous epithelium 12. Tertiary fold | | |


Figure 18.49. Placenta (Cotyledonary and Epitheliochorial), Cow. Detail of Figure 18.48. Highly branched chorioallantoic villi interdigitate with uterine crypts.

Figure 18.50. Placenta (Cotyledonary and Epitheliochorial), Cow. Detail of a portion of a placentome adjacent to the stalk. Note that the cryptal epithelium is cuboidal or flattened. The epithelium of the chorioallantoic villus consists of irregularly shaped cells and binucleate giant cells (diplokaryocytes).

Figure 18.51. Cervix of Uterus, Bitch. The mucosa of the cervix is thrown into folds. The epithelium of the bitch's cervix is stratified squamous. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.52. Cervix of Uterus, Mare. Cervical folds are evident.

Figure 18.53. Cervix of Uterus, Mare (Masson's). The cervical epithelium is simple columnar, except in the bitch (see Figure 18.51). Epithelial cells may be ciliated.








Figure 18.55 ×62.5




Figure 18.56 ×62.5



Figure 18.57 ×12.5




Figure 18.58 ×62.

- 1. Adventitia
- 2. Cavernous spaces
- Columnar cells
 Duct
- 5. Keratinized cells
- 6. Lamina propria
- 7. Minor vestibular gland
- 8. Muscularis
- 9. Stratified epithelium
- 10. Stratified squamous epithelium
- 11. Transitional epithelium
- 12. Urethral epithelium
- 13. Vestibular epithelium


Figure 18.54. Vagina, Anestrus, Bitch. The epithelium of the carnivore's anestrous vagina is stratified squamous to stratified cuboidal. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.55. Vagina, Estrus, Queen. In carnivores, the vagina is lined by a thickened, keratinized, stratified squamous epithelium during estrus. Exfoliated keratinized cells are visible in the vaginal lumen in this micrograph.

Figure 18.56. Urethra, x.s., Queen. Section was taken from the region close to the bladder. This portion of the urethra is lined by a transitional epithelium. (Photomicrograph of a histologic section borrowed from the College of Veterinary Medicine, Iowa State University.)

Figure 18.57. Junction of Vestibule and Urethra, x.s., Queen. Near the vestibule, the urethra is lined by a stratified squamous epithelium. Note the presence of cavernous spaces in the connective tissue adjacent to the muscularis. Such spaces occur only in the distal two-thirds of the urethra in the doe, ewe, and queen. In all other domestic mammals, cavernous spaces occur throughout the entire length of the urethra.

Figure 18.58. Vestibule, Queen. Detail of a minor vestibular gland. The secretory tubules of these branched, tubular glands are lined by columnar cells. Their ducts are lined by stratified squamous epithelium.








Figure 18.60 ×125




Figure 18.61 ×125




Figure 18.62 ×125




Figure 18.63 ×125

| KEY | | | |
|---|---|--|--|
| Columnar cells Duct Erythrocyte Intermediate cell | 5. Neutrophil 6. Parabasal cell 7. Superficial cell 8. Superficial intermediate cell | | |


Figure 18.59. Vestibule, Queen. Detail of a major vestibular gland. These glands are found in queens and ruminants. They are compound tubular glands with secretory units like those of the minor vestibular glands (see Figure 18.58).

Figure 18.60. Vaginal Smear, Anestrus, Bitch (Hema-3). During anestrus, parabasal and intermediate cells are the predominant epithelial cells present. See the introduction for a description of cell types. Neutrophils and bacteria may be present in limited numbers.

Figure 18.61. Vaginal Smear, Early Proestrus, Bitch (Hema-3). During early to mid proestrus, smears may contain neutrophils, erythrocytes, and various epithelial cell types (parabasal, intermediate, superficial intermediate, and superficial cells).

Figure 18.62. Vaginal Smear, Mid to Late Proestrus, Bitch (Diff-Quik). In late proestrus, superficial intermediate and superficial cells are predominant. Neutrophils decrease in number at this time.

Figure 18.63. Vaginal Smear, Estrus, Bitch (Diff-Quik). Most (90% or more) of the epithelial cells from a bitch in estrus are superficial cells. Erythrocytes may be present in small numbers. Some estrous smears may contain large numbers of bacteria. Neutrophils are not normally present.








Figure 18.65 ×12.5




Figure 18.66 ×62.5




Figure 18.67




Figure 18.68 ×62.5

- 1. Follicle, early
- 2. Follicle, late 3. Granulocytes
- 4. Intermediate cell
- 5. Medulla
- 6. Neutrophil
- 7. Oocyte

- 8. Oocyte, nucleus
 9. Oocyte, yolk-laden cytoplasm
 10. Parabasal cell
- 11. Superficial cell
- 12. Superficial intermediate cell
- 13. Vacuolar cells

Figure 18.64. Vaginal Smear, Diestrus, Bitch (Diff-Quik). There is a significant change in the numbers of epithelial-cell types during diestrus. Superficial cells decrease, and parabasal and intermediate cells increase. Neutrophils usually reappear during diestrus. Because erythrocytes may be present in smears from bitches in early diestrus, it is not possible to distinguish proestrus from diestrus on the basis of a single smear.

Figure 18.65. Ovary, Hen. A portion of the ovarian cortex with developing follicles.

Figure 18.66. Ovary, Vacuolar Cells, Hen. A portion of the cortex with a mass of fat-laden vacuolar cells. The latter may represent regressing postovulatory follicles.

Figure 18.67. Ovary, Vacuolar Cells, Hen. Detail of Figure 18.66. Vacuolar cells have pyknotic nuclei and contain numerous fat vacuoles. Cell boundaries are often indistinct.

Figure 18.68. Ovary, Granulocytes, Hen. Granulocytes are often found in the cortex of mature ovaries. The acidophilic granules of these cells impart a red tinge to a large area of the cortex in this micrograph.




Figure 18.69 ×250




Figure 18.70 ×25




Figure 18.71 ×125




Figure 18.72 ×12.5




Figure 18.73

| KEY | | | | |
|---|--|--|--|--|
| Atretic follicle Cloaca Developing follicle Infundibulum Isthmus Interstitial cells Magnum Membrana granulosa Membrana granulosa, thickened | 10. Oocyte, yolk-laden cytoplasm 11. Ovary 12. Perivitelline membrane 13. Theca externa 14. Theca interna 15. Uterus (shell gland) 16. Vagina 17. Yolk sphere | | | |

Figure 18.69. Ovary, Developing Follicle, Hen. A portion of the wall of a developing follicle. Note the flattened cells of the theca interna and the presence of interstitial cells in the theca externa.

Figure 18.70. Ovary, Atretic Follicle, Hen. In some atretic follicles, interstitial (luteal) cells proliferate, hypertrophy, and migrate inward (see Figure. 18.71).

Figure 18.71. Ovary, Atretic Follicle, Hen. Detail of Figure 18.70.

Figure 18.72. Ovary, Atretic Follicle, Hen. Cells of the membrane granulosa have proliferated, forming a thick layer characteristic of many atretic follicles.

Figure 18.73. Oviduct, Diagrammatic Drawing, Hen. The oviduct of the hen is divisible into an infundibulum, magnum, isthmus, uterus, and vagina.




Figure 18.74 ×62.5




Figure 18.75 ×62.5




Figure 18.76 ×250




Figure 18.77 ×12.5




Figure 18.78 ×25

| | KEY |
|--|--|
| Ciliated epithelium Glandular groove Lamina propria Mucosa Mucosal fold, infundibulum Mucosal fold, magnum Muscularis Muscularis, circular | 9. Primary fold 10. Pseudostratified columnar epithelium 11. Regenerating gland 12. Resting gland 13. Secondary fold 14. Secretory gland 15. Serosa 16. Tertiary fold |

Figure 18.74. Funnel of Infundibulum, Oviduct, x.s., Hen. The mucosa is thrown into shallow ridges that increase in height as the funnel narrows toward the neck region. The epithelium is ciliated, simple columnar. Scattered bundles of smooth muscle form the muscularis. A serosa covers the funnel externally.

Figure 18.75. Neck of Infundibulum, x.s., Oviduct, Hen. Tall primary mucosal folds bear secondary and tertiary folds.

Figure 18.76. Neck of Infundibulum, x.s., Oviduct, Hen. Detail of mucosa showing folds lined by ciliated columnar cells. The bottoms of the grooves between the folds are lined by nonciliated secretory cells.

Figure 18.77. Neck of Infundibulum and Magnum, x.s., Oviduct, Hen. The primary mucosal folds of the magnum are taller and broader vis-à-vis the infundibulum, because of the presence of numerous tubular glands.

Figure 18.78. Magnum, x.s., Oviduct, Hen. Portion of a fold. Depending on their activity, the tubular glands of the magnum exhibit distinctive features. Three morphologic phases of activity can be recognized (regenerating, secretory, resting). See Figures 18.80 and 18.81 for details.




Figure 18.79 ×250




Figure 18.80 ×250




Figure 18.81 ×256




Figure 18.82 ×12.5



Figure 18.83 ×125

- 1. Albumen
- 2. Ciliated cell
- 3. Epithelium, regenerating gland
- 4. Epithelium, resting gland
- 5. Epithelium, secretory gland6. Isthmus, primary fold
- 7. Magnum
- 8. Muscularis
- 9. Pseudostratified epithelium
- 10. Secretory cell
- 11. Tubular gland

Figure 18.79. Magnum, Oviduct, Hen. Detail of the epithelium. Ciliated columnar and secretory (goblet) cells comprise the epithelium of the magnum. The nuclei of the secretory cells are round and are located close to the base of the cell, whereas the nuclei of the ciliated cells are oval and occupy the central to apical region of the cell. Accordingly, the epithelium is pseudostratified columnar.

Figure 18.80. Magnum, Oviduct, Hen. Detail of regenerating tubular glands. These glands have clearly defined lumens. The secretory cells are cuboidal.

Figure 18.81. Magnum, Oviduct, Hen. Detail of secretory tubular glands and resting tubular glands. The cells of the secretory stage are characterized by pyknotic, basal nuclei. The entire cytoplasm is filled with strongly acidophilic granules. The glandular lumens may be dilated by secreted albumen. The cytoplasm of the cells in the resting stage has a frothy appearance, and the lumens of the glands are obscure.

Figure 18.82. Isthmus, x.s., Oviduct, Hen. The primary folds of the isthmus are not as broad as those of the magnum. Compare with Figure 18.77. They are somewhat angular in appearance. A portion of an adjacent region of the magnum is present in this micrograph.

Figure 18.83. Isthmus, Oviduct, Hen. A portion of the epithelium and underlying tubular glands. The epithelium is ciliated pseudostratified columnar. A tubular gland can be seen opening to the surface. The glandular cells do not undergo obvious cyclic changes as in the magnum.




Figure 18.84 $\times 12.5$




Figure 18.85 ×250




Figure 18.86 ×625




Figure 18.87 ×25




Figure 18.88 ×125

- 1. Apical cell
- 2. Basal cell, nucleus
- 3. Blood vessel 4. Concave surface
- 5. Duct
- 6. Erythrocyte in capillary
- 7. Glandular epithelium
- 8. Muscularis
- 9. Primary fold
- 10. Pseudostratified epithelium
 11. Secondary fold
- 12. Sperm-host gland
- 13. Vacuole

Figure 18.84. Uterus (Shell Gland), Oviduct, Hen. The folds of the uterus are not as broad as those of the magnum, and there is less glandular tissue. This section was taken from a uterus that had been fixed while containing an egg. Accordingly, the luminal surface is somewhat concave.

Figure 18.85. Uterus (Shell Gland), Oviduct, Hen. Ducts of complex, branched, tubular glands pierce the ciliated pseudostratified columnar epithelium at intervals. Ducts are formed from polygonal gland cells.

Figure 18.86. Uterus (Shell Gland), Oviduct, Hen. Basal cells (nuclei close to the basement membrane) of the pseudostratified epithelium may contain vacuoles above and below their nuclei. Apical cells (nuclei centrally located) contain numerous granules before releasing their secretion.

Figure 18.87. Vagina, x.s., Oviduct, Hen. The mucosa of the vagina is characterized by long, slender, primary folds bearing numerous small secondary folds. The muscularis is highly developed.

Figure 18.88. Vagina, Proximal, Oviduct, Hen. Sperm-host glands are tubular glands, lined by tall columnar cells, and are located within the mucosa of the vagina near the uterovaginal junction. Sperm are stored in these glands, remaining functional for up to 21 days.

THE EYE

MAMMALS

he eye is a sensory organ designed for vision. Basically, it is composed of a lens and a wall that is divided into three layers: an outer fibrous tunic (corneoscleral layer), a middle vascular tunic (uvea), and an inner retina. The fibrous tunic is divided into the posterior, opaque sclera and the anterior, transparent cornea. The vascular tunic includes the choroid, ciliary body, and iris. The retinal tunic consists of a 10-layered, photosensitive retina and a bilayered, nonphotosensitive portion that covers the ciliary body and the posterior surface of the iris.

Fibrous Tunic

Sclera

The sclera consists of densely interwoven bundles of collagenous fibers arranged parallel to the surface of the wall of the eye. There are also fibroblasts, some fine elastic fibers, and scattered melanocytes, especially in the innermost region of the sclera.

Cornea

The cornea is avascular. Its anterior (outer) surface is covered by the nonkeratinized, stratified squamous anterior epithelium. Below this layer is Bowman's membrane, which is not distinct in domestic mammals. The underlying stroma (substantia propria) is composed of thin lamellae of collagenous fibers oriented parallel to the corneal surface. Fibroblasts occur between the layers of fibers. Descemet's membrane is a relatively thick membrane that separates the stroma from the posterior epithelium. The latter consists

of a single layer of squamous to low cuboidal cells that covers the posterior surface of the cornea.

Limbus

The corneoscleral junction is called the limbus. Here, the regular collagenous lamellae of the corneal stroma merge with the interwoven fibers of the sclera. The appearance of the stratified squamous epithelium of the cornea differs from that of the bulbar conjunctiva, which overlies the sclera near the limbus. The deepest cells in the epithelium of the bulbar conjunctiva are smaller and more closely packed than those of the anterior epithelium of the cornea. In addition, the basal border of the conjunctival epithelium is uneven with the presence of an underlying, papillated layer of loose connective tissue. The boundary between the corneal epithelium and its underlying stroma, however, is smooth.

Vascular Tunic

Choroid

The **choroid** is the portion of the vascular tunic of the eye that lies between the sclera and the photosensitive retina. It contains numerous melanocytes. The fine network of connective tissue of the **suprachoroid layer** joins the sclera to the **vascular layer** of the choroid. The latter is composed of a profusion of blood vessels surrounded by loose connective tissue. The **choriocapillary layer** contains a thin network of capillaries that are distributed in a single plane. **Bruch's membrane**, a refractile membrane that lies between the choriocapillary layer and the pigment epithelium of the retina, is difficult to resolve.

A reflective tapetum lucidum (responsible for eyeshine) is located between the choriocapillary and vascular layers of the choroid in the dorsal portion of the eye. It is present in all domestic mammals except the pig. The horse and ruminants have a fibrous tapetum lucidum composed of layers of collagenous fibers and fibroblasts. The cat and dog have a cellular tapetum lucidum formed by flattened, pentagonal or hexagonal cells that appear bricklike in profile. The tapetal cells are filled with numerous rodshaped granules. The flat surfaces of the cells and the long axes of their rod-shaped granules lie parallel to the surface of the retina.

Ciliary Body

The ciliary body is an anterior continuation of the choroid that extends to the base of the iris. The loose connective tissue of the stroma contains smooth muscle, the ciliary muscle, which lies peripheral to an inner, vascular region. The epithelium of the ciliary body, which is formed by cells of the nonphotosensitive portion of the retina, is called the pars ciliaris retinae. It is a bilayer of cells consisting of a basal layer of pigmented cells and a surface layer of nonpigmented columnar cells. Short folds of the posterior surface of the ciliary body become longer toward the iris and form ciliary processes that project toward the lens. Zonular fibers extend from the processes to the lens capsule near the equator of the lens.

Iris

The iris is the most anterior part of the uveal tract. It forms a thin, contractile diaphragm with a central aperature, the **pupil.** The base of the iris is attached to the anterior portion of the ciliary body. The stroma of connective tissue of the iris contains many melanocytes and blood vessels. The stroma contains circumferentially arranged bundles of smooth muscle that form the sphincter (constrictor) muscle. The anterior surface of the iris is not covered by an epithelium, but rather by a discontinuous layer of stromal cells (fibroblasts and melanocytes). The posterior surface is covered by a bilayer of epithelial cells, the pars iridica retinae, which represents the most anterior continuation of the nonphotosensitive portion of the retina. It consists of a superficial layer of pigmented columnar cells and a basal layer of partially pigmented myoepithelial cells. The latter are elongated, radially arranged, contractile cells that form the dilator "muscle" of the iris. They have an apical, pigmented portion containing the nucleus and a nonpigmented basal portion. The nonpigmented regions of these cells border the stroma and appear as an acidophilic band. The pigmented portion of each myoepithelial cell lies just below the layer of pigmented columnar cells. In the horse, pig, and ruminant, a number of corpora nigra (iris granules) project from the pupillary margin of the iris. They are highly vascularized proliferations of the stroma and the pigmented epithelial cells of the iris.

Filtration Angle

At the peripheral margin of the anterior chamber is the filtration angle, the area between the limbus, the base of the iris, and the ciliary body. This triangular region is spanned by a latticework of trabeculae and intertrabecular, fluid-filled spaces. The trabeculae are composed of connective tissue and pigment cells and are covered by a single layer of squamous cells. They form the pectinate ligament, the uveal trabecular meshwork, and the corneoscleral trabecular meshwork. At the peripheral margin of the anterior chamber, excess aqueous humor passes through the openings within the pectinate ligament into the spaces of Fontana within the uveal trabecular meshwork. These spaces communicate with those of the corneoscleral trabecular meshwork, which drain into the scleral venous plexus. In the horse, the limbus does not overlap the pectinate ligament of the filtration angle, so that the pectinate ligament is apparent by direct examination of the eye. In the other domestic mammals the limbus covers the pectinate ligament, which is therefore obscured from view by the opaque sclera.

Retina

The retina is the innermost layer of the wall of the eye. The photosensitive portion lines the inner surface of the eye (adjacent to the cavity of the vitreous humor) from the ora ciliaris retinae to the optic disc. The latter is the point of transition from the photosensitive retina to the nonphotosensitive retina. From the ora ciliaris retinae, the nonphotosensitive portion continues anteriorly as a bilayer of

cells, forming the pars ciliaris retinae and the pars iridica retinae, which cover the ciliary body and the posterior surface of the iris, respectively.

From the choroid to the cavity of the vitreous humor, the 10 layers of the photosensitive retina are as follows:

Pigment epithelium
Layer of rods and cones
Outer limiting membrane (usually not apparent)
Outer nuclear layer
Outer plexiform layer
Inner nuclear layer
Inner plexiform layer
Ganglion cell layer
Nerve fiber layer
Inner limiting membrane

The cuboidal cells of the pigment epithelium contain few or no pigment granules in the part of the eye where the tapetum lucidum is located. In other parts of the eye, pigment granules are numerous in the cells of the pigment epithelium.

The nerve-fiber layer consists of axonal processes of the ganglion cells that converge at the **optic disc** and form the optic nerve. Because photoreceptor cells are not present here, this region is also referred to as the **blind spot**. Bundles of fibers of the optic nerve pass through perforations of the sclera. This sievelike part of the sclera is the **lamina cribrosa**.

Lens

The transparent, biconvex lens is avascular. It is composed entirely of epithelial cells enclosed within a homogeneous capsule. The cells on the anterior surface of the lens just below the capsule are simple cuboidal and form the lens epithelium. Toward the equator of the lens, the cells become long, prismatic, and arranged in meridional rows, forming lens fibers. As new lens fibers develop from the lens epithelium at the germinal zone of the equator, older lens fibers are displaced centrally and lose their nuclei. The lens is suspended by zonular fibers (suspensory ligaments) that extend from the lens capsule to the ciliary processes.

Chambers of the Eye

The eye contains three fluid-filled regions. The anterior chamber is bordered by the cornea, iris, and lens. The posterior chamber is located between the iris, lens, zonular fibers, and ciliary processes. Both of these chambers contain aqueous humor. The most posterior compartment, the cavity of the vitreous humor, lies behind the lens.

Conjunctiva

The **conjunctiva** is a thin, transparent, mucous membrane. The **bulbar conjunctiva** is continuous with the anterior surface of the cornea at the limbus and covers the sclera for a short distance. The **palpebral conjunctiva** lines the internal surface of the eyelids. The **fornix of the conjunctiva** is the point of reflexion of the bulbar and palpebral

conjunctiva. The epithelium of the conjunctiva varies from stratified squamous to stratified columnar and may even appear transitional. Goblet cells are often present. The underlying layer of loose connective tissue may contain diffuse or nodular lymphatic tissue.

Eyelids

The eyelids are covered internally by the palpebral conjunctiva and externally by thin skin. The skin contains hair follicles, sweat glands (glands of Moll), and sebaceous glands (glands of Zeiss). In the pig, the glands are particularly well developed. Between the dermis of the skin and the lamina propria of the palpebral conjunctiva is a plate of dense connective tissue, the tarsus (tarsal plate). Large, multilobular, sebaceous glands, called tarsal (Meibomian) glands, are embedded in the tarsus. Their central ducts open onto the palpebral surface near its junction with the skin.

Nictitating Membrane

The nictitating membrane (third eyelid) is a ventromedial fold of conjunctiva. It is supported by hyaline cartilage in the dog and ruminants and by elastic cartilage in the cat, horse, and pig. The superficial gland of the nictitating membrane surrounds the base of the cartilage. It is a serous gland in the horse and cat, mixed in the dog and ruminants, and mucous in the pig. The pig also has a Harderian gland (deep gland of the nictitating membrane) that produces a fatty secretion.

Lacrimal Gland

The lacrimal gland is a tubuloacinar gland, serous in the cat and mixed in the horse, ruminant, dog, and pig. It is predominantly a mucous gland in the pig and mostly serous in the horse and ruminant. There are also accessory lacrimal glands, such as Krause's gland, which may be serous or mixed.

CHICKEN

Lens

The eye of the chicken is quite different from that of mammals. Within the **capsule** the **lens** is divided into the **annular pad** and the **lens body**. The annular pad forms an outer ring around the equator of the lens body. It consists of radially arranged lens fibers with peripheral nuclei. In the lens body the lens fibers are oriented parallel to the optical axis of the eye, and some nuclei are present, primarily near the annular pad.

Fibrous Tunic

Sclera

A remarkable feature of the avian sclera is the presence of a ring of overlapping scleral ossicles, anteriorly, and a cup-shaped layer of hyaline cartilage, the scleral cartilage, posteriorly. The latter terminates internal to the scleral ossicles. Dense connective tissue encloses the scleral ossicles and extends posteriorly, peripheral to the cartilage layer.

Cornea

The layers of the **cornea** of the chicken are similar to those of mammals. Bowman's membrane, however, is thicker and therefore more apparent in histologic preparations. Descemet's membrane is relatively thin and less distinct.

Vascular Tunic

Choroid

The choroid is a thick, vascularized coat with numerous pigment cells. The suprachoroid abuts the thin perichondrium of the scleral cartilage. The vascular layer of the choroid contains blood vessels and large spaces embedded in loose connective tissue. The choriocapillary layer is separated from the pigment epithelium of the retina by an indistinct Bruch's membrane. No tapetum lucidum is present in the chicken.

Ciliary Body

The ciliary body is a thin layer of loose connective tissue with a thick outer region of numerous elastic fibers. It is covered by a bilayer of more or less cuboidal, pigmented basal cells and cuboidal to columnar, nonpigmented surface cells. Folds of the lining of the ciliary body form ciliary processes that fuse to the lens capsule in the region of the equator of the lens. More posteriorly, zonular fibers extend from the ciliary body to the lens capsule. The ciliary muscles (Crampton's and Brücke's) are skeletal muscles that lie across from the ciliary body, just inside the main portion of the sclera.

Iris

The iris is thickest just above its narrow base, then tapers toward the pupillary margin. The stroma contains a sphincter and dilator muscle. Both of these are formed from small skeletal muscle cells that contain lipid vacuoles. The dilator muscle is sparse and posterior to the thicker sphincter muscle. The anterior (corneal) surface of the iris is covered by a simple layer of nonpigmented, flattened epithelial cells. The posterior (lens) surface of the iris is covered by a stratified layer of pigmented epithelial cells, three to five cells thick.

Retina

The photosensitive retina of the chicken is composed of 10 layers, as in mammals, but unlike that in mammals is avascular. The cells of the pigment epithelium are considerably different in the chicken. They are tall and narrow rather than cuboidal. The nucleus occupies the smaller, basal region of each cell, which contains few or no pigment granules. The apical portion is filled with rod-shaped pigment granules that are oriented parallel to the long axis of the cell. The apical cytoplasm often appears to be separated into tufts or strands of pigment granules.

Pecten

The pecten is a thin, highly vascular, pleated membrane that protrudes into the cavity of the vitreous humor from the ventral surface of the eye. Its base is secured intermittently to the linear, optic disc. The apical surface is attached to a thickened mass of pectineal tissue called the bridge. The pecten is characterized by an extensive network of capillaries lined by thick endothelial cells with plump nuclei. Polymorphic pigment cells fill the spaces between the capillaries and larger vessels. The pecten is draped by a covering membrane, which is thought to be continuous with the inner limiting membrane of the retina.

Filtration Angle

The filtration angle of the chicken is somewhat different from that of mammals. It is filled by a trabecular meshwork formed by the pectinate ligament (uveal meshwork) and the scleral trabecular meshwork. The pectinate ligament is a loose network of elastic fibers covered by simple squamous cells. It spans the filtration angle from the scleral trabecular meshwork to the iris and the elastic tissue of the ciliary body. The trabecular meshwork of the pectinate ligament encloses the spaces of Fontana. The latter communicate with the spaces of the scleral trabecular meshwork, which is formed by collagenous and elastic fibers. These spaces communicate with the canal of Schlemm within the sclera.

Structures Associated With the Eye

The chicken has a thin, well-developed nictitating membrane. A supportive cartilage is absent. The inner surface of the eyelids is lined by the palpebral conjunctiva. The external surface is covered by thin skin with sparse feathers. No glands are present.

The **lacrimal gland** is a small, tubular gland that produces a mucous secretion. It lies medial to the caudal part of the lower eyelid. The **Harderian gland** is a larger gland that lies on the dorsal posterior surface of the eye. It is characterized by numerous plasma cells that surround the tubular secretory units.

| Word Ro | Word Roots | | |
|-----------------|------------------------------------|--|--|
| ROOT | MEANING | EXAMPLE SENTENCE | |
| Corpus Nigra | Body Black, dark | The <i>corpora nigra</i> that project from the papillary margin of the iris are highly pigmented. | |
| Crib | A sieve | Bundles of fibers of the optic nerve pass through the sievelike layer called the <i>lamina cribrosa</i> of the sclera. | |
| Lachrym | Tears | The <i>lacrimal</i> gland produces tears. | |
| Limbus | An edge, usually curved | The <i>limbus</i> is the circular junction of the cornea with the sclera. | |
| Palpebra | Eyelid | The palpebral conjunctiva lines the internal surface of the eyelid. | |
| Scler | Hard | The <i>sclera</i> is the tough outer connective tissue layer of the wall of the eye. | |
| Tape Lucid | A carpet Light, clear, shiny | The <i>tapetum lucidum</i> is the reflective layer of the choroid. | |
| Vitre Humor | Glassy A fluid | Vitreous humor is a transparent fluid that fills the posterior compartment of the eye. | |






Figure 19.1. Eye, Sagittal Section, Dog. The relationship between various morphologic components of the canine eye are depicted.










Figure 19.4 ×125





Figure 19.5 ×250



| KEY | | |
|-----------------------------|-----------------------------|--|
| 1. Anterior chamber | 14. Lens epithelium | |
| 2. Bulbar conjunctiva | 15. Lens fibers, new | |
| 3. Ciliary body | 16. Lens fibers, old | |
| 4. Ciliary muscle | 17. Limbus | |
| 5. Ciliary process | 18. Palpebral conjunctiva | |
| 6. Cornea | 19. Pars ciliaris retinae | |
| 7. Corneoscleral trabecular | 20. Pectinate ligament | |
| meshwork | 21. Posterior chamber | |
| 8. Elastic fiber | 22. Sclera | |
| 9. Filtration angle | 23. Scleral venous plexus | |
| 10. Germinal zone | 24. Spaces of Fontana | |
| 11. Iris | 25. Vascular layer, choroid | |
| 12. Lens | , , | |
| 13. Lens capsule | | |


Figure 19.2. Eye, Dog. Anterior, peripheral portion of the eye.


Figure 19.3. Lens, Equator, Horse. Newly formed and older lens fibers are visible in this section through the germinal zone. The latter is the marginal band of lens epithelium that lies around the equator. Its cells are capable of dividing throughout adult life.


Figure 19.4. Ciliary Body, Dog. The epithelium of the ciliary body is called the pars ciliaris retinae. This portion of the non-photosensitive retina consists of an inner (closer to the cavity of the vitreous humor), nonpigmented layer and an outer, heavily pigmented layer of cells.

Figure 19.5. Ciliary Body, Cat (Orcein). In addition to smooth muscle (see Figure 19.4), the ciliary body contains an abundance of elastic fibers.

Figure 19.6. Eye, Horse. In a section through a horse's eye, the limbus does not overlap the pectinate ligament. Compare with Figure 19.7. The filtration angle is indicated by the triangular-shaped area marked by the dashed line.










Figure 19.10 \times 62.5




Figure 19.11 ×250

- 1. Bulbar conjunctiva
- 2. Ciliary process
- 3. Cornea
- 4. Corneoscleral trabecular meshwork
- 5. Descemet's membrane
- 6. Filtration angle 7. Iris
- 8. Limbus

- 9. Melanocyte
 10. Myoepithelial cell, cytoplasm
 11. Myoepithelial cell, nucleus
- 12. Myoepithelial cells

- 13. Nonpigmented cells
- 14. Palpebral conjunctiva
- 15. Pectinate ligament
- 16. Pigmented cells
- 17. Pigmented surface cells 18. Sclera
- 19. Spaces of Fontana
- 20. Sphincter muscle 21. Stroma
- 22. Uveal trabecular meshwork
- 23. Zonular fibers

Figure 19.7. Eye, Pig. In a section through the eye of a domestic mammal other than the horse, the limbus overlaps the pectinate ligament. The area of the filtration angle is outlined by dots.

Figure 19.8. Filtration Angle, Pig. Collagenous fibers, elastic fibers, fibroblasts, and pigment cells form the pectinate ligament and the uveal trabecular meshwork. The corneoscleral trabecular meshwork is a three-dimensional latticework of fine fibers of connective tissue and fibroblasts lying adjacent to the sclera.

Figure 19.9. Ciliary Processes, Pig. The nonpigmented cells of the epithelium of the ciliary processes cover the pigmented cells. Together these two layers of cells comprise the pars ciliaris retinae, which extends from the ora ciliaris retinae to the iris. The nonpigmented cells give rise to the zonular fibers.

Figure 19.10. Iris, Horse. The back surface of the iris is covered by a continuation of the pars ciliaris retinae and is called the pars iridica retinae. The surface layer of the pars iridica retinae consists of heavily pigmented cells, and its inner layer is formed by the contractile, pigmented myoepithelial cells that dilate the iris. Anteriorly, the iris is covered by a discontinuous layer of stromal

Figure 19.11. Iris, Dog. The heavily pigmented cells of the pars iridica retinae cover the posterior surface of the iris. The myoepithelial cells of the pars iridica retinae are partially pigmented in the region of their nuclei.




Figure 19.12 ×625



Figure 19.13 ×12.5




Figure 19.14 ×62.5




Figure 19.15 ×125




Figure 19.16 ×62.5

KEY 1. Anterior epithelium 10. Melanocyte 2. Bulbar conjunctiva, epithelium 11. Ora ciliaris retinae 3. Bulbar conjunctiva, lamina 12. Pars ciliaris retinae propria 13. Posterior epithelium 4. Choroid 14. Retina, photosensitive 5. Cornea 15. Sclera 16. Space artifacts 6. Corneal stroma 7. Corpus nigrum 17. Stroma 8. Descemet's membrane 9. Iris


Figure 19.12. Iris, Cat. In the cat, the melanocytes of the iris are binucleate and contain rod-shaped melanosomes.


Figure 19.13. Corpus Nigrum, Goat. In ungulates, the pupillary border of the iris is differentiated into corpora nigra, which are vascularized outgrowths of the stroma and pigmented epithelium of the iris.

Figure 19.14. Cornea, Dog. The anterior surface is covered by a nonkeratinized stratified squamous epithelium. The posterior surface is covered by squamous or low cuboidal cells.

Figure 19.15. Junction of Cornea and Bulbar Conjunctiva, Pig. Both the cornea and bulbar conjunctiva are covered by a nonkeratinized stratified squamous epithelium at their junction. The basal border of the conjunctiva is irregular, and the cells of its deepest layers are smaller than those of the anterior epithelium of the cornea.

Figure 19.16. Ora Ciliaris Retinae, Dog. The transition zone between the photosensitive and the nonphotosensitive (pars ciliaris retinae) regions of the retina is named the ora ciliaris retinae.










Figure 19.20

- 1. Blood vessel
- 2. Choriocapillary layer
- 3. Choroid
- 4. Cone
- 5. Ganglion cell layer
- 6. Inner limiting membrane
- 7. Inner nuclear layer
 8. Inner plexiform layer
- 9. Lamina cribrosa
- 10. Layer of rods and cones
- 11. Melanocytes12. Nerve fiber layer

- 13. Nerve fibers, bundle of
- 14. Optic disc
- 15. Optic nerve
- 16. Outer nuclear layer
- 17. Outer plexiform layer
- 18. Pigment epithelium
- 19. Retina 20. Rod
- 21. Sclera
- 22. Tapetum lucidum
- 23. Vascular layer, choroid

Figure 19.17. Optic Nerve, Sagittal Section, Dog. Nerve fibers of the retina converge to form the optic nerve at the optic disc (blind spot).

Figure 19.18. Lamina Cribrosa, x.s., Dog. At the lamina cribrosa the connective tissue of the sclera forms a sievelike framework, which subdivides the optic nerve into bundles of fibers.

Figure 19.19. Retina, Choroid, and Part of the Sclera, Pig. Note that the dendrites (cones) of cone cells of the pig are particularly plump and easily recognized.

Figure 19.20. Fibrous Tapetum Lucidum, Sheep. The tapetum lucidum of ruminants and horses is a compact membrane of connective tissue sandwiched between the choriocapillary and vascular layers of the choroid. The cells of the pigment epithelium of the retina contain few or no pigment granules where a tapetum lucidum is present. Compare with Figure 19.21.

Figure 19.19




Figure 19.21 ×250




Figure 19.22 ×250




Figure 19.23 ×250




Figure 19.24 ×625




Figure 19.25 ×12.5

| KEY | |
|--------------------------------------|---------------------------------------|
| 1. Anterior chamber | 13. Melanocytes |
| Bulbar conjunctiva | 14. Nerve fiber layer |
| 3. Capillary, x.s. | 15. Nictitating membrane, bulbar |
| 4. Choriocapillary layer | surface |
| 5. Cornea | Outer nuclear layer |
| 6. Cytoplasmic process | 17. Outer plexiform layer |
| 7. Descemet's membrane | 18. Pectinate ligament |
| 8. Elastic cartilage | 19. Pigment epithelium |
| 9. Ganglion cell layer | 20. Sclera |
| 10. Inner nuclear layer | 21. Tapetal cell |
| 11. Inner plexiform layer | 22. Tapetum lucidum |
| 12. Layer of rods and cones | 23. Vascular layer, choroid |


Figure 19.21. Retina, Choroid, and Part of the Sclera, Dog. Where the tapetum lucidum is lacking from the choroid layer, the cells of the pigment epithelium of the retina contain numerous melanosomes.

Figure 19.22. Cellular Tapetum Lucidum, Dog. In profile, the cells of the tapetum lucidum of carnivores are bricklike. Note that in this section the cells of the pigment epithelium of the retina are indistinct and lack pigment. Compare with Figure 19.21.

Figure 19.23. Melanocytes of Choroid Layer, Dog. Melanocytes of the choroid layer are flat polygonal cells with cytoplasmic processes. Their polygonal shape is evident in this tangential cut through the choroid layer.

Figure 19.24. Tapetum Lucidum, Dog. Tapetal cells are flattened and have a pentagonal or hexagonal outline, which is apparent in this tangential section through the choroid layer. The cells are filled with numerous small rods, whose long axes parallel the flat surfaces of the cells.

Figure 19.25. Nictitating Membrane and Cornea, Horse. A portion of the bulbar surface of the nictitating membrane and its supportive cartilage are shown. The nictitating membrane is a fold of the ventromedial portion of the conjunctiva. It contains elastic cartilage in the horse, pig, and cat, and hyaline cartilage in the dog and ruminants.








Figure 19.27 ×125




Figure 19.28 ×62.5




Figure 19.29 ×12.5



Figure 19.30 ×125

| KEY | | |
|---|--|--|
| Diffuse lymphatic tissue Duct Epidermis Goblet cell Hair follicle Hyaline cartilage Mucous acinus Palpebral conjunctiva | 9. Serous acinus 10. Stratified columnar epithelium 11. Stratified squamous epithelium 12. Superficial gland 13. Sweat gland 14. Tarsal gland 15. Tarsus | |


Figure 19.26. Superficial Gland of the Nictitating Membrane, Dog. The base of the cartilage of the nictitating membrane is surrounded by the superficial gland.

Figure 19.27. Superficial Gland of the Nictitating Membrane, Dog. This gland is mixed in the dog and ruminants. It is serous in the horse and cat and mucous in the pig.

Figure 19.28. Palpebral Conjunctiva, Pig. The palpebral conjunctiva is a mucous membrane that lines the inner surface of the eyelid. Its stratified epithelium varies from squamous through columnar and may even appear transitional. Goblet cells may be present.

Figure 19.29. Eyelid, Lower, Horse. The outer surface of the eyelid is covered by thin skin, while the inner surface is lined by the palpebral conjunctiva. The tarsal gland is a multilobulated gland whose duct opens onto the palpebral surface near the margin of the eyelid. The tarsal gland is surrounded by a condensed layer of connective tissue, the tarsus.

Figure 19.30. Eyelid, Upper, Pig. Numerous tubular sweat glands (and sebaceous glands, not shown) occur in the skin surface of the eyelid of the pig.







Figure 19.32 ×62.5



×250




Figure 19.35 ×125

- 1. Acinus
- 2. Epidermis
- 3. Fornix of conjunctiva
 4. Hair follicle
- 5. Interlobular connective tissue
- 6. Intralobular duct
- 7. Krause's gland 8. Palpebral conjunctiva

- 9. Sclera 10. Sebaceous gland 11. Skeletal muscle tissue
- 12. Smooth muscle tissue
- 13. Sweat gland
- 14. Tarsal gland
- 15. Tarsus

Figure 19.31. Eyelid, Lower, Goat. The skin surface of the eyelid contains hair follicles, sweat glands, and sebaceous glands.

Figure 19.32. Eyelid, Lower, Goat. Bundles of smooth and skeletal muscle fibers are scattered in the connective tissue between the tarsus and the skin surface of the eyelid.

Figure 19.33. Krause's Gland, Pig. Krause's gland is a small, accessory lacrimal gland (serous in this preparation) located near the fornix of the conjunctiva.

Figure 19.34. Harderian Gland, Pig. Among domestic mammals, this gland is present only in the pig. It secretes a fatty product.

Figure 19.35. Lacrimal Gland, Cow. A compound tubular acinar gland. The lacrimal gland is predominantly a serous gland in ruminants and horses.







Figure 19.36







| KEY | | |
|---|--|--|
| 15. Lens body | | |
| 16. Lens fibers, annular pad | | |
| 17. Lens fibers, lens body | | |
| 18. Nonpigmented epithelium, iris | | |
| 19. Pectinate ligament | | |
| 20. Pigmented epithelium, iris | | |
| 21. Posterior chamber | | |
| 22. Sclera | | |
| 23. Scleral cartilage | | |
| 24. Scleral ossicle | | |
| Scleral trabecular meshwork | | |
| | | |

12. Crampton's muscle13. Dilator muscle

14. Iris

Figure 19.36. Eye, Drawing of an Anterolateral Segment, Chicken. This drawing illustrates the relative positions of various structures seen in several micrographs of the chicken's eye in this chapter.

26. Spaces of Fontana


27. Sphincter muscle 28. Zonular fibers


Figure 19.37. Lens, Chicken. A portion of the annular pad and lens body.

Figure 19.38. Filtration Angle, Chicken. The filtration angle is bordered by the cornea, iris, ciliary body, and sclera in the chicken. It is bridged by a trabecular meshwork of the pectinate ligament, which encloses the spaces of Fontana.


Figure 19.39. Ciliary Process, Chicken (Masson's). Ciliary processes occur below the base of the iris and fuse with the lens capsule of the annular pad. The ciliary epithelium also attaches to the capsule by zonular fibers.

279









12

Figure 19.44 ×62.5

| KEY | |
|---|---|
| 1. Bowman's membrane 2. Ciliary body 3. Descemet's membrane 4. Dilator muscle cell 5. Elastic fiber 6. Lipid vacuole 7. Nonpigmented epithelium, iris | 10. Posterior epithelium 11. Scleral cartilage 12. Scleral ossicle 13. Space artifact 14. Sphincter muscle cell 15. Squamous cell, nucleus 16. Stratified squamous epithelium |
| 8. Pectinate ligament 9. Pigmented epithelium, iris | 17. Stroma 18. Zonular fibers |

Figure 19.40. Iris, Chicken. The iridial musculature of the chicken is composed of skeletal muscle cells, which are characterized by the presence of numerous lipid vacuoles. Unlike mammals, the anterior (corneal) surface of the iris is covered by a layer of flattened, nonpigmented epithelial cells. The posterior (lens) surface of the iris (see Figure 19.39) is covered by a stratified pigmented epithelium that is three to five cells thick.

Figure 19.41. Junction of Ciliary Body and Iris, Chicken (Orcein). The elastic fibers of the pectinate ligament insert into an elastic meshwork of the ciliary body.

Figure 19.42. Pectinate Ligament, Chicken. The elastic fibers of the pectinate ligament are covered by a simple squamous epithelium.

Figure 19.43. Cornea, Chicken. A well-developed Bowman's membrane separates the anterior, stratified squamous epithelium of the cornea from the underlying stroma.

Figure 19.44. Sclera, Chicken. The sclera is strengthened anteriorly by overlapping bony plates (scleral ossicles). Posteriorly, it consists of a thin layer of cartilage. See Figure 19.36 for the location of ossicles and cartilage. The region of overlap of these skeletal elements is shown here. In this section the ciliary body has separated from the sclera, creating a space artifact.




Figure 19.45 ×62.5




Figure 19.46 ×250




Figure 19.47 ×12.5




Figure 19.48 ×250




Figure 19.49 ×125

| 1. Blood vessels 2. Bridge 3. Capillary 4. Choriocapillary layer 5. Choroid 6. Covering membrane 7. Ganglion cell layer 8. Inner nuclear layer 9. Inner nuclear layer | | | |
|---|--|--|--|
| 2. Bridge 13. Nerve fiber layer 3. Capillary 14. Optic nerve 4. Choriocapillary layer 15. Outer nuclear layer 5. Choroid 16. Outer plexiform layer 6. Covering membrane 17. Pecten 7. Ganglion cell layer 18. Pigment epithelium 8. Inner nuclear layer 19. Retina | KEY | | |
| 10. Layer of rods and cones 11. Loose connective tissue | 2. Bridge 3. Capillary 4. Choriocapillary layer 5. Choroid 6. Covering membrane 7. Ganglion cell layer 8. Inner nuclear layer 9. Inner plexiform layer 10. Layer of rods and cones | 13. Nerve fiber layer 14. Optic nerve 15. Outer nuclear layer 16. Outer plexiform layer 17. Pecten 18. Pigment epithelium 19. Retina 20. Scleral cartilage | |

Figure 19.45. Retina and Choroid, Chicken. The bulk of the choroid is composed of blood vessels and large spaces supported by a loose connective tissue. The layers of the retina are comparable with those of mammals.

Figure 19.46. Retina and Choroid, Chicken. Cells of the pigment epithelium of the retina are tall and contain rod-shaped pigment granules. The basal region of each cell contains the nucleus and a few pigment granules.

Figure 19.47. Pecten, Chicken. The pecten is a thin, folded, and heavily pigmented membrane that projects into the vitreous humor from the posteroventral surface of the eye.

Figure 19.48. Pecten, Chicken. Numerous, polymorphic melanocytes are interspersed through this highly vascularized, nutritive membrane. The large capillaries are lined by thick endothelial cells with plump nuclei.

Figure 19.49. Bridge of Pecten, Chicken. This thickened, highly pigmented mass of pectineal tissue is located along the free edge of the pecten.





Figure 19.50 ×62.5



×25 **Figure 19.51**





Figure 19.52 $\times 25$



×250 **Figure 19.53**

- 1. Choroid

- 2. Collecting duct
 3. Dermis
 4. Epidermis
 5. Extrinsic muscle
- 6. Feather follicle 7. Harderian gland
- 8. Herbst corpuscle
- 9. Interlobular connective tissue
- 10. Plasma cells
- 11. Retina
- 12. Scleral cartilage
- 13. Secretory tubule

Figure 19.50. Eyelid, Chicken. A thin epidermis covers the eyelid. Herbst corpuscles are associated with a feather follicle.

Figure 19.51. Lacrimal Gland, Chicken. This compound tubular gland produces a mucoid secretion and is organized into lobules.

Figure 19.52. Harderian Gland, Chicken. This accessory immunologic gland contains a multitude of plasma cells. It is located on the dorsal posterior surface of the eye.

Figure 19.53. Harderian Gland, Chicken. Detail of the Harderian gland showing numerous plasma cells surrounding the vacuolated cells of the tubular secretory units.

THE EAR

ensations of sound and balance are received by separate and specialized areas of the ear before being transmitted to the brain, where they are interpreted. Based on anatomy, location, and function, the ear is divisible into external, middle, and internal components. The external ear collects sound waves, which it channels to the tympanic membrane. Vibrations produced in the latter are transmitted by the ossicles of the middle ear to fluids of the inner (internal) ear, where they generate movements of the delicate basilar membrane. Such movements stimulate sensory hair cells from which impulses are relayed by sensory nerves to the brain, where the sound is identified.

MAMMALS

External Ear

The external ear of domestic mammals is comprised of the pinna (auricle), which collects sound, and an external auditory meatus, which conveys the sound waves to the tympanic membrane.

The external auditory meatus is lined by a continuation of the surface skin. Hair, sebaceous glands, and tubular ceruminous glands are present. The combined secretions of these glands, plus sloughed epithelial cells, form cerumen (ear wax). The outer portion of the meatus is supported by cartilage, the remainder by bone.

Middle Ear

Middle-ear ossicles (malleus, incus, and stapes) are located in the tympanic cavity. They bridge the cavity from the tympanic membrane to the oval window located within the petrous part of the temporal bone.

The tympanic cavity is surrounded by bone. The tympanic membrane forms the lateral wall of the cavity. It is composed of a thin, outer layer of epithelium that is continuous with the skin of the external auditory meatus, a thin layer of connective tissue, and an inner layer of simple squamous or cuboidal epithelium. The remainder of the cavity is lined by ciliated columnar and simple squamous cells. The latter cover the ossicles, as well as portions of the wall of the cavity.

Inner Ear

The membranous labvrinth of the inner ear consists of the cochlear duct, sacculus, utriculus, and semicircular ducts. Cavities within the petrous segment of the temporal bone, lined by periosteum and containing perilymph (a fluid similar to cerebrospinal fluid), comprise the bony labyrinth and house the membranous labyrinth. The cavities containing the semicircular ducts are called the semicircular canals; the one containing the saccule (sacculus) and utricle (utriculus) is called the vestibule; and the one containing the cochlear duct (membranous cochlea, scala media) is named the cochlear canal (cochlea; bony cochlea). The cochlear canal spirals like a snail shell around a central pillar of bone, the modiolus. A thin shelf of bone, the osseous spiral lamina, travels up the modiolus like the thread of a screw. The number of turns in the cochlear canal varies. There are two and one-half in the horse, three in the cat, and four in the pig, for example.

Each semicircular duct is lined by a mesothelium, is filled with endolymph, and bears an expansion, the ampulla. A sensory structure, the crista ampullaris, is located in each ampulla. The sensory hair cells and supporting cells of each crista are covered by a gelatinous cupula. When the latter is deflected during rotational movements of the head, the sensory cells are stimulated and impulses are sent to the brain, where the signals are interpreted.

Both the saccule and utricle are filled with endolymph and are lined, in part, by maculae, which are patchlike collections of sensory hair cells and supporting cells. The remainder of these structures is lined by mesothelium. Embedded in the outer surface of the gelatinous otolithic membrane covering the maculae are numerous crystalline particles of calcium carbonate called otoliths (otoconia, statoconia). As the membrane shifts in response to gravity acting on the otoliths, sensory cells of the maculae are stimulated. Impulses sent to the brain in response to the stimulus make the animal aware of the position of its head in space. Also, because of the effect of inertia on the otolithic membranes when the body suddenly begins to move or slow down, hair cells are stimulated and the sensations of acceleration and deceleration are experienced.

The spirally organized cochlear duct is filled with endolymph and is roughly triangular in cross section. One side of the duct is attached to the spiral ligament, a thickening of the periosteal lining of the cochlear canal. This side consists of a stratified cuboidal epithelium, the stria vascularis. Capillaries occur among the superficial cuboidal cells of the stria. The side of the duct opposite the stria

is pointed. The floor of the duct is formed from the fibrous basilar membrane, which extends from the spiral ligament to the osseous spiral lamina. The roof is formed from the vestibular (Reissner's) membrane, which consists of two adjacent layers of simple squamous epithelium. Above the roof is a large chamber, the scala vestibuli, which is filled with perilymph. Below the floor of the cochlear duct is another large chamber filled with perilymph, the scala tympani. All three scalas follow a spiral path to the top of the cochlear canal. At the apex the scala vestibuli communicates with the scala tympani through a tiny opening called the helicotrema.

The upper surface of the basilar membrane supports the acoustically sensitive organ of Corti, which is bathed by the endolymph within the cochlear duct. The lower surface of the basilar membrane is lined by a simple squamous epithelium that faces the scala tympani. The organ of Corti is comprised of sensory hair cells and various different supporting cells. Overlying the organ of Corti and extending from the spiral limbus (an elevation of connective tissue above the osseous spiral lamina) is the proteinaceous tectorial membrane. Stereocilia of the sensory cells of the organ of Corti contact the tectorial membrane. The stereocilia are displaced when the basilar membrane vibrates in response to sound waves passing through the fluid-filled scalas. The sensory cells respond to this perturbation by initiating impulses in the cochlear nerve, which are transmitted to the brain for interpretation. The stimulatory sound waves are dissipated through the secondary tympanic membrane of the round window located in the lower part of the medial wall of the tympanic cavity.

CHICKEN

The ear of the chicken consists of the same basic components as that of the mammal, but there are some differences.

External and Middle Ear

Although an external auditory meatus is present in the chicken, it is relatively short, and there is no pinna.

The middle ear is lined by a cuboidal epithelium that also covers the columella, a single partially ossified rod that extends from the tympanic membrane to the oval window. The **columella** transmits vibrations from the tympanic membrane to the internal ear, taking the place of the malleus, incus, and stapes of mammals.

Inner Ear

Unlike mammals, the saccule (sacculus) of the inner ear contains two maculae. The cochlear duct is a short, narrow, slightly curved tube. It possesses a terminal expansion, the lagena, a structure peculiar to birds. The lagena contains a macula that is similar in structure and function to other maculae (see under Mammals). The cochlear duct is separated from the overlying scala vestibuli by the tegmentum vasculosum. The latter is composed of a thin membrane

of connective tissue integrated with a highly folded epithelium containing numerous blood vessels. The epithelium faces the cavity of the cochlear duct. The tegmentum occupies the same position as the vestibular membrane in mammals. The common wall separating the cochlear duct from the scala tympani below is formed from the basilar membrane, a platform that supports the organ of Corti (papilla acustica or basilaris). As in mammals, the organ of Corti is composed of sensory and supporting cells and is overlain by a tectorial membrane that is in contact with the sensory hairs (stereocilia) of the sensory cells.

| Word Roots | | |
|----------------------------|--------------------------------|--|
| ROOT | MEANING | EXAMPLE SENTENCE |
| Audi Meatus | Hearing A passageway | The external <i>auditory meatus</i> conveys sound waves to the tympanic membrane. |
| Helix Trema | A coil A hole | The <i>helicotrema</i> is the tiny opening at the apex of the bony cochlea through which the scala vestibuli and scala tympani communicate. |
| Labyrinth | A maze | The bony labyrinth and membranous labyrinth of the inner ear are complex, mazelike passageways. |
| Malleus Incus Stapes | A hammer Anvil A stirrup | The three ossicles of the middle ear, the <i>malleus</i> , <i>incus</i> , and <i>stapes</i> , resemble a hammer, anvil, and stirrup, respectively. |
| Scala | Ladder or stairs | The passageways, namely the scala vestibuli, scala media, and scala tympani, follow a spiral path to the top of the cochlear canal. |
| Tecto | A roof or cover | The tectorial membrane covers the stereocilia of the sensory cells of the organ of Corti. |
| Tympani | A drum | The <i>tympanic membrane</i> is nicknamed the eardrum. |




Figure 20.1 $\times 25$





Figure 20.2 ×25



×25 Figure 20.3




Figure 20.4




Figure 20.5 ×62.5

| KEY | |
|--|---|
| Adipose tissue Bone Ceruminous gland Duct Elastic cartilage Epidermis | 7. External auditory meatus 8. Hair follicle 9. Pigment granules 10. Sebaceous gland 11. Tympanic cavity 12. Tympanic membrane |


Figure 20.1. External Auditory Meatus, Puppy. The outer portion of the meatus is supported by elastic cartilage. The thin epidermis is underlain by numerous sebaceous glands and a few ceruminous glands. Small hair follicles are present.

Figure 20.2. External Auditory Meatus, Puppy. The external auditory meatus, near the tympanic membrane, contains large sebaceous glands.

Figure 20.3. External Auditory Meatus, Goat. The outer portion of the external auditory meatus with numerous ceruminous glands is shown. Hair follicles and portions of sebaceous glands are also

Figure 20.4. Ceruminous Gland, Goat. The secretory epithelium of these apocrine glands varies from cuboidal to columnar. The cells contain tiny, brown pigment granules.

Figure 20.5. Tympanic Membrane, Periphery, Puppy. The tympanic membrane has a core of collagenous fibers. Its outer (external auditory meatus) surface is covered by a stratified squamous epithelium; its inner (tympanic cavity) surface is covered by a simple squamous or cuboidal epithelium.








Figure 20.7 ×250




Figure 20.8 ×12.5



Figure 20.9 ×25




Figure 20.10 ×125

KEY 1. Annular ligament 2. Articular cartilage 3. External auditory meatus 4. Facial nerve 5. Footplate, stapes 6. Incus 7. Joint capsule with elastic fibers 8. Lenticular process, articular 10. Malleus 11. Saccule, macula of 12. Stapes, articular cartilage 13. Synovial cavity 14. Temporal bone, petrous part 15. Tympanic cavity 16. Tympanic membrane 17. Utricle, macula of

cartilage

9. Ligament

Figure 20.6. Malleus and Tympanic Membrane, Puppy. The handle (manubrium) of the malleus is attached to the tympanic membrane.

18. Vestibule

Figure 20.7. Tympanic Membrane, Puppy. Where the manubrium of the malleus is embedded in the tympanic membrane, the connective tissue of the tympanic membrane is thicker than elsewhere, and blood vessels are abundant.

Figure 20.8. Portion of Stapes and Incus, Puppy. The footplate of the stapes is attached to the oval window by an annular ligament (broken on one side in this section). The stapes articulates with the lenticular process of the incus.

Figure 20.9. Joint, Malleus and Incus, Puppy. The head of the malleus articulates with the body of the incus in this synovial joint.

Figure 20.10. Junction of Lenticular Process and Stapes, Puppy. The stapes articulates with the lenticular process of the incus. See Figure 20.8 for orientation.

287




Figure 20.11 ×125




Figure 20.12




Figure 20.13




Figure 20.14 $\times 62.5$

- 1. Ampulla, cavity
- 2. Annular ligament
- 3. Connective tissue
- 4. Cupula, portion of 5. Mixed glands
- 6. Otolith
- 7. Otolithic membrane
- 8. Pseudostratified epithelium
- 9. Sensory cell, nucleus
- 10. Stapes, articular cartilage
- 11. Supporting cell, nucleus
 12. Temporal bone, petrous part
- 13. Type 1 cell

Figure 20.11. Annular Ligament, Stapes, Puppy. The stapes is fastened to the circular cartilage of the oval window by the fibroelastic, annular ligament. See Figure 20.8 for orientation.

Figure 20.12. Macula of Saccule, Puppy. Otoliths are embedded in a gelatinous otolithic membrane, which lies on an epithelium consisting of sensory and supporting cells. Chalicelike, Type I sensory cells and the basal nuclei of supporting cells are evident in this micrograph.

Figure 20.13. Crista Ampullaris, Puppy. This ridge of sensory epithelium, supported by connective tissue, protrudes into the ampulla of a semicircular duct and is oriented at right angles to the long axis of the duct. The epithelium consists of sensory and supporting cells similar to those found in the maculae. A mass of gelatinous material, the cupula, covers the surface epithelium.

Figure 20.14. Guttural Pouch, Horse. This diverticulum of the Eustachian tube is lined by a ciliated, pseudostratified columnar epithelium with goblet cells. Mixed glands occur in the lamina propria.



Figure 20.15 ×18






Figure 20.16 ×90


| KEY | |
|--|---|
| Basilar membrane Cochlear duct Cochlear nerve Inner tunnel Modiolus Organ of Corti Osseous spiral lamina Scala tympani Scala vestibuli | 10. Spiral ganglion11. Spiral ligament12. Spiral limbus13. Spiral tunnel14. Stria vascularis15. Tectorial membrane16. Temporal bone, petrous part17. Vestibular membrane |


Figure 20.15. Cochlea, Puppy. Cross section through a portion of the spiral cochlea. See Figure 20.16 for details of cochlear duct region.

Figure 20.16. Cochlear Duct in Cochlea, Puppy. Detail of the region of the cochlear duct (scala media, membranous cochlea) and surrounding temporal bone that forms the cochlea.









1. Basilar membrane

2. Blood vessel of connective

4. Cartilaginous frame, caudal

5. Cartilaginous frame, rostral

9. Endolymphatic space

11. Homogeneous cells 12. Light cell

tissue base 3. Capillary

> Cochlea Cochlear nerve

8. Dark cell

10. Hair ćells

- 13. Organ of Corti
- 14. Perilymphatic space
- 15. Raphe

KEY

- 16. Scala media (cochlear duct)
- 17. Scala tympani
- 18. Scala vestibuli
- 19. Semicircular canal, wall
- 20. Semicircular duct, wall
- 21. Supporting cells
- 22. Tectorial membrane
- 23. Tegmentum vasculosum

Figure 20.17. Semicircular Canal, x.s., Chicken. The semicircular canal is a part of the bony labyrinth. It contains the semicircular duct, a part of the membranous labyrinth. The duct is lined by a simple squamous epithelium except at the raphe, where cuboidal cells form the lining.

Figure 20.18. Cochlear Duct in Cochlea, x.s., Chicken (Masson's). Only a small portion of the bony cochlea that encloses the scala media is visible. The three passageways of the cochlea (the scala vestibuli, scala media, and scala tympani) are evident. The tegmentum vasculosum (see Figure 20.20) separates the scala vestibuli and the scala media. A portion of the organ of Corti is magnified in Figure 20.19.

Figure 20.19. Organ of Corti, Cochlea, x.s., Chicken (Masson's). The organ of Corti is composed of sensory cells and supporting cells with an overlying tectorial membrane.

Figure 20.20. Tegmentum Vasculosum, Cochlea, Chicken (Masson's). This thick membrane rests on a thin base of connective tissue and possesses a highly folded epithelial surface with numerous vascular loops. The epithelium consists of light and dark cells. Dark cells have a constricted neck region that extends to the surface of the epithelium. Their basal portion contains the nucleus and is irregularly shaped. The cytoplasm is very dense. Light cells have a pale cytoplasm and surround the dark cells.

GLOSSARY

A band: The part of a sarcomere, of both skeletal and cardiac muscle cells, containing actin and myosin myofilaments and located centrally between the I bands of the sarcomere.

Abomasum: The glandular stomach of a ruminant.

Accessory gland: The name applied to any of the following glands of the male reproductive system: bulbourethral, prostate, seminal vesicle.

Acid stain (anionic stain): A dye, used for histologic or cytologic staining procedures, that carries a negative charge. Such dyes attach electrostatically to positively charged components of cells or other tissue elements.

Acidophil (alpha cell): An acidophilic chromophil found within the pars distalis of the pituitary.

Acidophilic: Literally, possessing an affinity for acid (anionic) dyes which carry a negative charge. For example, positively charged (cationic) amino groups of proteins in cells and tissues become stained with the acid dye, eosin, which carries a net negative charge.

Acinus: A small, grape-shaped secretory unit of an acinar gland or tubuloacinar gland.

Adenohypophysis: The glandular portion of the pituitary. It is derived during development from Rathke's pouch. It includes the pars distalis, pars tuberalis, and pars intermedia.

Adipocyte: Synonym for fat cell.

Adipose tissue: Aggregates of adipocytes within loose connective tissue.

Adrenal cortex: That part of the adrenal gland surrounding the medulla. It is divisible into four zones of cells that produce steroid hormones.

Adrenal medulla: The region interior to the cortex of the adrenal gland. The chromaffin cells of the medulla secrete the hormones epinephrine (adrenalin) and norepinephrine.

Adventitia: The external layer of connective tissue covering a structure.

Afferent arteriole: Smallest branch of the renal artery that delivers blood directly to the glomerulus of a renal corpuscle.

Agranulocyte: A leukocyte that lacks specific granules, e.g., a lymphocyte or monocyte.

Air capillary: In birds, a tiny respiratory tubule forming part of a network connecting parabronchi. Respiratory exchange occurs between the blood in surrounding blood capillaries and the gas within air capillaries.

Air sac: One of several rather large, thin-walled, and airfilled sacs found within various parts of the body cavity of a bird. Many hollow bones of the chicken contain extensions of air sacs. The designation air sac is also used as a synonym for alveolus.

Alveolar bone: The bone lining the alveolus (root socket) of a tooth. Collagenous fibers attach the root of the tooth to the alveolar bone.

Alveolar duct: The portion of the bronchial tree of a mammal lying between a respiratory bronchiole and an alveolar sac.

Alveolar sac: A terminal, baglike expansion of the bronchial tree whose wall is formed entirely from alveoli.

Alveolar septum: All of the tissue separating the cavities of adjacent alveoli of the lungs formed by type I alveolar cells and pulmonary capillaries.

Alveolus: One of millions of thin-walled, tiny sacs forming the terminal limits of the bronchial tree. It is sometimes called an air sac. The term is also used as a synonym for the acinus of an exocrine gland.

Ameloblasts: Cells that form the enamel organ of a tooth and produce the enamel.

- Amorphous ground substance: A viscous and colorless material that fills the spaces between the fibers, cells, and vessels of connective tissue.
- Ampulla: A dilated portion of a tubular structure such as the ampulla of a semicircular duct of the inner ear, or the ampulla of the vas deferens or oviduct.
- Anal gland: A tubuloacinar gland found within the submucosa and muscularis of the anal canal of carnivores and pigs.
- Anal sac: One of a pair of pouchlike glands found within the tissue adjacent to the anus.
- Anastomosis: A connection between two structures.
- Anestrus: In nonprimate mammals, the period of sexual inactivity.
- Anisocytosis: A condition exemplified by the existence of considerable variation in the size and shape of an animal's erythrocytes. It is often associated with certain pathological conditions, but may occasionally be seen in a healthy individual.
- Annular ligament: A fibroelastic ligament that attaches the stapes of the middle ear to the cartilage of the oval window.
- **Annular pad:** The portion of the lens of the chicken's eye located around the equator of the lens.
- Annular sinus: A large blood sinus lying between the inner and outer layers of the connective tissue sheath of a sinus hair follicle.
- Anterior chamber: The space, filled with aqueous humor, that is bounded by the cornea, iris, and lens of the eye.
- Anterior epithelium: The stratified squamous epithelium of the outer surface of the cornea.
- Antrum: A cavity, such as the antrum of an ovarian follicle.
- Aortic body: An encapsulated chemical receptor located between the pulmonary artery and aorta. It detects changes in oxygen, carbon dioxide, and pH levels and functions in neural reflexes that adjust cardiac output and respiratory rate.
- Apocrine gland: A gland whose secretory cells release their product by pinching off blebs of cytoplasm containing secretory product from their free surface.
- **Aponeurosis:** A white fibrous membrane that connects a muscle to a bone or fascia.
- **Arachnoid layer:** The middle meninx enveloping the spinal cord and brain.
- **Argyrophilic:** A substance having an affinity for silver.
- Arrector pili muscle: The smooth muscle originating from the dermis of the skin and inserting on the connective tissue sheath of a hair follicle. Elevates the hair.
- Arteriole: The smallest of the arteries. Consists of an endothelium and one or two layers of smooth muscle.
- Arteriovenous anastomosis (arteriovenous shunt): A vessel that enables blood to flow directly from an artery to a vein without passing through a capillary bed.
- **Artery:** A blood vessel that carries blood away from the heart to a capillary bed.
- **Artifact:** Any alteration in appearance or structure that has been caused by artificial means.
- **Arytenoid cartilage:** One of two small cartilages located at the back of the larynx.

- **Astrocyte:** A neuroglial cell having long processes and a starlike appearance.
- Atresia: The degenerative regression of an ovarian follicle.
- Atretic follicle: An ovarian follicle that has undergone atresia.
- **Atrium:** A cavity or chamber such as an atrium of the heart or an air vesicle of a chicken's parabronchus.
- Auerbach's plexus: A collection of nerve cells and fibers lying between the circular and longitudinal layers of the muscularis externa of the digestive tract.
- Axon: The extension of a neuron that transmits impulses away from the cell body.
- Axon hillock: The place of origin of an axon from a nerve cell body. The hillock lacks Nissl granules.
- Azurophilic granule: Nonspecific granules found in the cytoplasm of some leukocytes.
- Band cell: A granulocyte in the process of development and characterized by the presence of a straplike unsegmented nucleus.
- Barb: A subdivision, bearing barbules, of a vane of a contour feather. The barbules of adjacent barbs interlock with one another by means of their hooklets.
- Barb stem: The stalk of the barb of a feather.
- Barbule: A projection, bearing hooklets, from a barb.
- Basal cell: A pale cell found in the fundus of the gastric glands of the ventriculus of the chicken. The term basal cell is also applied to a cell attached to the basement membrane of a stratified epithelium.
- Basement membrane: A fibrous membrane separating the cells of an epithelium from the underlying connective tissue
- Basic stain (cationic stain): A dye used for histologic or cytologic staining procedures that carries a positive charge. Such stains attach electrostatically to negatively charged elements within cells and tissues.
- Basilar membrane: A membrane of the inner ear that supports the organ of Corti.
- **Basophil:** A granulocyte characterized by large, basophilic, specific granules. Also, a cell of the pars distalis of the pituitary gland.
- Basophilic: Literally, having an affinity for basic (cationic) dyes that carry a net positive charge. For example, negatively charged (anionic) groups in cells and tissues, such as phosphate groups of amino acids, sulfate groups of glycosaminoglycans, and carboxyl groups of proteins, become colored by the basic dyes toluidine blue and methylene blue.
- Basophilic erythroblast: A developing cell of the erythroid line characterized by an intensely basophilic cytoplasm and a large, round, and deeply stained nucleus.
- Bile canaliculus: A tiny channel, formed from the cell membranes of adjacent hepatocytes, that receives bile from the hepatocyte and delivers it to a bile ductule.
- Bile ductule: A subdivision of a bile duct found within a portal area (tract) and lined by a simple cuboidal epithelium.
- Bistratified epithelium: A stratified epithelium consisting of two layers of cells.

- Blastocyst: The vesicular embryonic stage of a mammal that consists of an inner cell mass and outer cell mass (trophoblast) surrounding a fluid-filled cavity (blastocele, blastocyst cavity).
- Blind spot: The optic disc of the eye that lacks photoreceptor cells.
- Bone marrow (active): The site of the formation of blood cells (hematopoiesis) within the marrow cavity of a bone.
- Bone matrix: The hard, calcified substance of a bone, consisting of hydroxyapatites and collagenous fibers.
- Bony labyrinth: The three interconnected bony spaces, the cochlea, vestibule, and semicircular canals, of the inner ear located within the temporal bone of the skull.
- Bowman's capsule: The hollow, bulbus, initial portion of a renal corpuscle. Its outer wall is the capsular epithelium (simple squamous), and its inner wall or glomerular epithelium is composed of podocytes.
- Bowman's gland: A mucoserous gland located in the lamina propria below the respiratory epithelium. Its duct opens onto the respiratory epithelium.
- Bowman's membrane: A membrane of connective tissue that lies below the anterior epithelium of the cornea.
- Brain sand: A granular, calcified material that may be found in various parts of the brain.
- Bronchiole: A subdivision of the bronchial tree that begins at the termination of the smallest bronchus. Bronchioles vary in diameter, amount of smooth muscle present, and whether a ciliated epithelium is present.
- Bronchus: Any subdivision of the bronchial tree from the end of the trachea to a bronchiole. All bronchi are supported by cartilage and are lined by an epithelium that varies from ciliated pseudostratified columnar in the larger bronchi to ciliated simple columnar in the smaller ones.
- Bruch's membrane: A thin, refractile membrane of the retina separating the choriocapillary layer of the choroid from the pigment epithelium of the retina.
- Brücke's muscle: A ciliary muscle (skeletal) of the chicken eye.
- Brunner's gland (duodenal or submucosal gland): A compound tubuloacinar mucous gland found within the submucosa, and to some degree the lamina propria, of the duodenum.
- **Brush border:** A border formed from numerous microvilli of variable length on the apical surface of certain epithelial cells, e.g., those of the proximal convoluted tubules of the kidney.
- **Buffy coat:** The thin layer of leukocytes between the plasma and packed red blood cells in a sample of blood that has been centrifuged.
- Bulbar conjunctiva: That portion of the conjunctiva located on the surface of the eyeball.
- Bulbourethral gland (Cowper's gland): An accessory male reproductive gland that empties into the urethra. Its secretion forms part of the seminal fluid.
- Bursa of Fabricius: A saclike lymphatic diverticulum from the dorsal surface of the proctodeum of birds.
- Calamus: The hollow quill of a feather.

- Calcified cartilage: Cartilage matrix that has become impregnated with calcium salts, as in the zone of calcification of the epiphyseal disc.
- Canal of Hering: A tiny channel, lined by simple cuboidal epithelium, that connects a bile canaliculus to a bile ductule within a portal tract (area).
- Canal of Schlemm: A channel within the sclera that drains excess aqueous humor from the eye to the circulatory system.
- Canaliculi: In bone, tiny channels that contain the processes of osteocytes.
- Capillary: A blood vessel ranging from 4 to 12 micrometers in diameter that is networked with other capillaries and located between an arteriole and venule.
- Capsular epithelium: The outer simple squamous epithelium of Bowman's capsule.
- Capsule of Glisson: A thin layer of connective tissue enveloping the liver.
- Cardiac gland: A mucous gland located in the cardiac gland region of the glandular stomach.
- Cardiac gland region: The proximal part of the glandular stomach preceding the fundic gland region.
- Cardiac muscle: The striated, involuntary muscle of the
- Cardiac skeleton: A supportive framework of connective tissue found at various locations within the wall of the heart.
- Carotid body: A chemical receptor located at the bifurcation of the carotid arteries. It detects changes in oxygen, carbon dioxide, and pH levels and functions in neural reflexes that adjust cardiac output and respiratory rate.
- Caruncle: In ruminants, any nonglandular region of the mucosa of the uterus.
- Cavernous space: A thin-walled vein that forms part of the erectile tissue of the urethra.
- Cavity of the vitreous humor: A large space behind the lens of the eye that contains a transparent and colorless gelatinous mass, the vitreous body (humor).
- Cecum: In mammals, that portion of the large intestine that unites with the small intestine. In chickens, one of two diverticuli that arise from the region of junction of the ileum and large intestine.
- Cellular tapetum lucidum: A cellular reflective layer of the choroid of the eye of cats and dogs.
- Cementoid (precementum): Uncalcified cementum of a tooth, produced by cells called cementoblasts.
- Cementum: A layer of bonelike mineralized tissue covering the dentin of the root of a tooth.
- Central artery: A branch of the splenic artery that passes through the white pulp of the spleen.
- Central canal: The fluid-filled cavity within the center of the spinal cord.
- Central nervous system: That part of the nervous system consisting of the brain and spinal cord.
- Central pallor: The pale central area, evident in a surface view, of the biconcave erythrocyte.
- Centroacinar cell: An extension of an intercalated duct of the pancreas into the interior of a secretory acinus.
- Cerebellum: A part of the brain that lies above the medulla and pons and below the posterior part of the cerebrum.

- Cerebrospinal fluid: The clear fluid found within the ventricles of the brain and the central canal of the spinal cord.
- Cerumen: A soft, waxy substance formed from the combined secretions of the ceruminous and sebaceous glands of the external auditory meatus and containing sloughed epithelial cells.
- Ceruminous gland: A gland of the external auditory meatus whose secretions contribute to the composition of the cerumen.
- Cervical os: The opening of the cervix into the vagina. Chestnut: A small round or oval thickening of the epidermis located on the inner side of the legs of a horse.
- Chief cell (stomach): A cell found in the fundic gland region of the stomach that secretes pepsinogen.
- Chondrocyte: A cartilage cell surrounded by cartilage
- **Chondrogenic:** The capacity to produce cartilage. For example, the perichondrium is a chondrogenic layer.
- Chorioallantoic membrane: An extraembryonic membrane of amniotes formed from the fusion of the chorion and the allantois.
- Choriocapillary layer: A network of capillaries of the choroid, distributed in a single plane, below Bruch's membrane of the eye.
- Chorion laeve: A part of the chorionic sac, having a smooth surface, that is not involved in the formation of the placenta.
- Choroid: The portion of the uvea (vascular tunic) of the eye located between the sclera and the photosensitive retina.
- Choroid plexus: A highly vascularized portion of the roof of the fourth ventricle and of other ventricles of the brain whose villi are surrounded by cerebrospinal fluid.
- Chromaffin cells: Cells of the adrenal medulla that form clusters and anastomosing cords separated by sinusoids.
- Chromophil: A cell of the pars distalis of the pituitary that can be stained with various kinds of histologic dye substances.
- **Chromophobe:** A small cell of the pars distalis of the pituitary that stains poorly or not at all with dyes.
- Cilia: Relatively long, motile extensions of the free surface of the cell membrane of certain epithelial cells.
- Ciliary body: The portion of the uvea between the choroid and the iris.
- Ciliary muscle: The musculature of the ciliary body.
- Ciliary process: A projection from the surface of the pars ciliaris retinae of the ciliary body to which zonular fibers are attached.
- Circumanal gland: A partly sebaceous gland found in the subcutis near the anus of the dog.
- Circumferential lamellae: The bony lamellae that lie parallel to the outer and inner surfaces of long bones.
- Circumvallate papilla: A large, round elevation bearing numerous taste buds, found on the surface of the tongue.
- Classic liver lobule: A cylindrical subunit (about 1×2 mm) of the liver that is composed of plates of hepatocytes separated by an extensive network of sinusoids.

- Claw: A keratinized derivative of the skin found on the tips of the digits.
- Clear cell: A cell of the secretory units of the carpal gland of the pig.
- Clearing: In a tissue sample being processed, the replacement of alcohol with a substance miscible with paraffin prior to infiltrating the sample with paraffin.
- **Clitoris:** A structure of the female that is homologous to the glans penis of the male.
- Cloaca: A chamber, internal to the vent, of various vertebrates that receives digestive wastes, reproductive cells, and urinary products.
- Cochlea (cochlear canal): A spiral-shaped cavity within the temporal bone of the skull forming a subdivision of the bony labyrinth of the inner ear. It contains the scala tympani, scala vestibuli, and the cochlear duct, which is part of the membranous labyrinth.
- Cochlear duct (membranous cochlea; scala media): A spirally arranged part of the membranous labyrinth, whose walls are formed from the basilar membrane, stria vascularis, and vestibular membrane.
- Collagen: A form of fibrous protein of which several types exist.
- Collagenous fibers: Fibers formed from one of the types of the fibrous protein collagen.
- Collecting tubule: The part of a uriniferous tubule that receives urine filtrate from a nephron.
- Colloid: A gelatinous material found within follicles of the thyroid gland and the pars intermedia of the pituitary gland, among other places.
- **Columella:** An ossicle of the middle ear of birds and some other vertebrates.
- Comb: A highly vascularized derivative of the skin found atop the head of the chicken.
- Compact bone: A bone whose dense matrix contains few marrow spaces.
- Compound follicle: A hair follicle bearing several hair bulbs.
- Conjunctiva: An epithelial layer covering part of the bulbar surface of the eye and the inner surface of the eyelid.
- Connecting duct: Tubules of the epididymis of the rooster, which are also called excretory canals.
- Connective tissue proper: A category of connective tissue that includes loose, dense, reticular, elastic, and adipose tissues.
- Connective tissue sheath (of hair follicle): A layer of connective tissue surrounding a hair follicle.
- Contour feather: A flight feather of birds with a central shaft consisting of a rachis supporting the vanes and a hollow quill.
- **Coprodeum:** The portion of the cloaca that joins the large intestine.
- **Cornea:** The anterior, transparent portion of the fibrous tunic of the eye.
- Corneal stroma (substantia propria): Lamellae of collagenous fibers oriented parallel to the corneal surface and located between Bowman's and Descemet's membranes of the cornea.
- Corneoscleral trabecular meshwork: A meshwork of trabeculae of connective tissue, fibroblasts, and pigment

- cells found in the filtration angle of the eye immediately adjacent of the sclera.
- Corneous cells: Keratinized cells of a feather follicle or surface epidermis.
- **Corneous connection:** A strand of tissue extending from the corneous layer of a feather follicle to the calamus of a developing feather.
- Corona (of lymphatic nodule): The external layer, consisting of numerous small lymphocytes, of an activated lymphatic nodule.
- Corona radiata: The columnar cells abutting the zona pellucida and forming the innermost layer of the cumulus oophorus of an ovarian follicle.
- Coronary region: The proximal, peripheral part of the horse's hoof located above the laminae.
- Corpora amylacea: Concretions of casein and cellular debris found within the cavities of the secretory units of the mammary gland.
- Corpora nigra: Highly vascularized proliferations of the pupillary margin of the iris.
- Corpus albicans: Scar tissue remaining following regression of the corpus luteum of the ovary.
- Corpus cavernosum: A mass of erectile tissue located within the body of the penis.
- Corpus cavernosum clitoridis: Erectile tissue of the clitoris.
- Corpus luteum: An endocrine gland derived from granulosa cells and theca interna cells of a postovulatory ovarian follicle; its cells secrete progesterone and estrogen.
- Corpus spongiosum (corpus cavernosum urethra): Erectile tissue within the glans of the penis and surrounding the penile urethra.
- Cortex (hair): That portion of a hair located between the cuticle and the medulla.
- Cortical nephron (chicken): A nephron with a relatively small renal corpuscle located in the cortex of the kidney.
- Cotyledonary placenta: The type of placenta found in ruminants where numerous, bean-shaped portions of the chorioallantoic membrane form the functional connections with the endometrium.
- Crampton's muscle: One of the ciliary muscles (skeletal) of the chicken's eye.
- Crenated: Having a corrugated or scalloped border, e.g., the cell membrane of an erythrocyte following the loss of water.
- Crista ampullaris: A sensory structure located within the ampulla of a semicircular duct of the inner ear.
- **Crop:** A caudal diverticulum of the esophagus of the chicken used for the temporary storage of food.
- Crypt of Lieberkühn (intestinal gland): A simple tubular gland located within the lamina propria of the small and large intestine.
- Cumulus oophorus: The mound of granulosa cells surrounding an oocyte of the ovarian follicle of a mammal.
- Cupula: The gelatinous covering of the sensory hair cells of the crista ampullaris.
- Cuticle (hair): The layer of flat cells, arranged in a shinglelike manner, on the surface of a hair.

- Cytoplasmic bridge: The region of adhesion (at the desmosomes) between adjacent epithelial cells of the stratum spinosum of the epidermis.
- Cytotrophoblast: The cellular layer of the trophoblast (covering layer of the blastocyst) that contributes to the formation of the placenta.
- Dark zone (glandular stomach): The distal and longest part of the fundic gland region of the stomach of carnivores; characterized by a thick mucosa and relatively shallow gastric pits.
- Deciduate placenta: A type of placenta wherein the chorioallantoic membrane has fused with the endometrium.
- Dendrite: The neuronal process that receives stimuli.
- Dense irregular connective tissue: A form of connective tissue proper that consists of relatively thick collagenous fibers arranged as a feltwork with scant space between them. The predominant cell is the fibroblast.
- Dense regular connective tissue: A form of connective tissue proper that consists of relatively thick collagenous fibers, arranged in parallel, with scant space between them. The fibroblast is the only cell type present.
- Dental lamina: A band of ectodermal cells from the embryonic jaw that grow into the underlying mesenchyme. Buds from the dental lamina give rise to the primordia of the enamel organs of the teeth.
- Dental papilla: A projection of mesenchyme of the embryonic jaw into the developing enamel organ.
- Dental pulp: The soft tissue of the pulp cavity of a tooth.
- **Dental sac:** The mesenchymal precursor of the connective tissue surrounding a developing tooth, or the connective tissue surrounding a developing tooth.
- **Dentin:** The ivory that forms the mass of the tooth.
- **Dermal papilla:** A projection of the dermis into the overlying epidermis.
- **Dermis:** The connective tissue of the skin located beneath the epidermis.
- Descemet's membrane: An acellular layer separating the stroma from the posterior epithelium of the cornea. Diaphysis: The shaft of a long bone.
- Diencephalon: A subdivision of the brain composed of the thalamus, subthalamus, and hypothalamus.
- Diestrus: That part of the estrous cycle, between metestrus and anestrus, when development and secretory activity of the endometrial glands peak.
- Diffuse lymphatic tissue: A type of lymphatic tissue characterized by a moderate amount of scattered lymphocytes.
- Diffuse placenta: The name given to a placenta when the chorioallantoic membrane makes a major structural contribution.
- Digital cushion: The loose connective tissue (subcutis) below the dermis of the digital pad.
- Digital pad: The soft, shock-absorbing tissue formed by the epidermis, dermis, and subcutis beneath the digits of many mammals.
- **Diplokaryocyte:** A binucleate giant cell of the epithelium of a chorionic villus of the placenta of the cow.

Distal convoluted tubule: That portion of a nephron located between the loop of Henle and a collecting tubule.

Dorsal plate (claw): The keratinized, upper part of a claw. Dorsal root ganglion: The ganglion of the dorsal root of a spinal nerve.

Down feather: A soft, fluffy feather that lacks barbules with hooklets.

Duct of the epididymis: The highly convoluted duct forming a major part of the epididymis.

Dura mater: The outer meninx of dense connective tissue surrounding the brain and spinal cord.

Eccrine (merocrine) gland: The cells of these glands release their product by exocytosis.

Efferent arteriole: The arteriole carrying blood away from the glomerulus of a renal corpuscle.

Efferent ductule: One of several small tubes connecting the rete testis to the duct of the epididymis.

Elastic cartilage: A type of cartilage whose matrix contains numerous elastic fibers.

Elastic fiber: A component of certain connective tissues that forms from the protein elastin.

Ellipsoid: A fusiform structure, composed of macrophages, that surrounds a portion of each of the capillaries of the penicillus of the spleen.

Embedding: When an infiltrated tissue is transferred to a fresh solution of embedding medium (e.g., melted paraffin) and the paraffin is then allowed to harden.

Embryonal connective tissue: A form of connective tissue exemplified by mesenchyme and mucous connective tissue.

Enamel: The hard, glistening material covering the exposed surface of a tooth.

Endocardium: The tissue layer lining the atria and ventricles of the heart.

Endochondral bone: Bone that has been formed by replacing a cartilaginous model.

Endocrine gland: A ductless gland.

Endolymph: A clear fluid contained within the semicircular ducts, sacculus, utriculus, and cochlear duct of the inner ear.

Endometrial gland: A simple tubular gland of the endometrium of the uterus.

Endometrium: The mucosa of the uterus.

Endomysium: The connective tissue immediately surrounding individual muscle cells of a fascicle of a muscle.

Endoneurium: The connective tissue immediately surrounding the processes of nerve cells within a fascicle of a nerve.

Endosteum: The layer of squamous osteogenic cells lining the entire marrow cavity and extending into the Haversian canals.

Endotheliochorial placenta: A type of placenta, found in carnivores, where the maternal and fetal blood are separated by four layers of tissue.

Endothelium: The simple squamous epithelium lining the cardiovascular system and lymphatic vessels.

Enterochromaffin cell: A form of APUD cell (amineprecursor uptake decarboxylase cell) found in the gastrointestinal tract, among other places; it can be stained with bichromate solutions and produces either a hormone or a paracrine (a substance that acts locally by diffusing to a target organ).

Eosinophil: A granulocyte characterized by the presence of eosinophilic specific granules.

Ependymal cell: A neuroglial cell that lines the ventricles of the brain and the central canal of the spinal cord.

Epicardium: The outermost layer of the heart.

Epidermal collar: A thick ring of epidermal cells situated at the base of the follicle of a feather.

Epidermal laminae: The platelike epidermal structures of the wall of the horse's hoof.

Epidermal peg: A downward extension of the epidermis that alternates with upward-extending dermal papillae.

Epidermis: The ectodermally derived stratified squamous epithelium forming the surface layer of the skin.

Epididymis: A highly coiled tube of the male reproductive system located between the efferent ductules and the vas deferens.

Epimysium: The outer sheath of connective tissue of a muscle.

Epineurium: The outer sheath of connective tissue of a nerve.

Epiphyseal disc: A plate of hyaline cartilage between the epiphysis and diaphysis of a developing (growing) long bone.

Epiphysis: The end (proximal or distal) of a long bone.

Epithelial tuft: A grouping of simple columnar cells at the apex of a follicle within the bursa of Fabricius of a chicken.

Epitheliochorial placenta: A placenta with six tissue layers between the maternal and fetal blood streams.

Epithelium: A cellular tissue that covers external surface

Epithelium: A cellular tissue that covers external surfaces or lines cavities.

Erectile tissue: A highly vascular tissue that allows for an increase in turgidity of the penis or clitoris.

Ergot: An epidermal thickening found on the posterior surface of a horse's foot at the level of the distal end of the metacarpal bone.

Erythrocyte: A red blood cell.

Estrous cycle: A cyclic event of a female nonprimate mammal involving changes in anatomic structure of reproductive organs, changes in physiologic condition, and changes in sexual behavior.

Estrus: That segment of the estrous cycle during which the female is sexually receptive of the male.

Euchromatic: The relatively pale appearance of the nucleus of a fixed and stained cell when the chromatin is not highly coiled.

Exocrine gland: A gland with a duct system that carries a secretion to a body surface or cavity.

Exocytosis: A process that occurs when a cytoplasmic vesicle fuses with the cell membrane and releases its contents into the extracellular space.

External auditory meatus: The canal of the external ear. External ear: That portion of the ear external to the tympanic membrane, including the external auditory meatus and the pinna.

- External elastic membrane: The elastic membrane external to the tunica media of some arteries.
- External root sheath: The layer of cells in the wall of a hair follicle between the inner root sheath and the connective tissue sheath.
- Eyelid: The movable, muscular fold covering the eye.
- Eyeshine: The reflected glow in the eye of some animals at night due to the presence of a tapetum lucidum in the choroid.
- Fascia: A sheet of connective tissue surrounding, investing, or binding together parts of the body.
- **Fascicle:** A bundle of structures such as a bundle of axons or a bundle of muscle cells.
- **Feather follicle:** The epidermal structure from which a feather grows.
- **Feather pulp:** The vascularized, mesenchyme-like tissue in the center of a developing feather.
- Feather sheath: The thin, epidermal surface layer of the calamus of a growing feather. It eventually disintegrates, except for a collarlike remnant around the calamus.
- Fibroblasts: Ubiquitous cells of the connective tissue, responsible for producing the precursors of collagen and elastin and for producing the amorphous ground substance.
- Fibrocartilage (fibrous cartilage): Dense connective tissue containing isolated groups of chondrocytes surrounded by small amounts of cartilage matrix.
- Fibroelastic tissue: Connective tissue proper containing a mixture of collagenous and elastic fibers.
- **Fibrous tapetum lucidum:** A reflective layer of the choroid of the eyes of horses and ruminants that is composed of collagenous fibers and fibroblasts.
- **Fibrous tunic:** An external sheath comprised of fibers of connective tissue.
- Filiform papilla: An outgrowth on the surface of the tongue bearing threadlike projections or spines.
- Filoplume: A small, hairlike feather.
- **Filtration angle:** In the eye, the angle formed between the limbus, ciliary body, and the base of the iris.
- **Fimbria of the infundibulum:** The fimbria is the fringed border of the infundibulum of the oviduct.
- **Fixative:** A chemical or mixture of chemicals used to preserve the structural characteristics of fresh tissue.
- Foliate papilla: One of several foldlike elevations of the surface of the tongue.
- Follicular tonsil: A tonsil with deep invaginations (crypts) of its surface epithelium.
- Forestomach: The nonglandular portion of the stomach of the horse, ruminant, or pig.
- Formed elements: Collectively, the erythrocytes, leukocytes, and platelets of the blood.
- Fornix of conjunctiva: The point of reflection of the bulbar and palpebral conjunctiva.
- Fossa: A cavity or a pit.
- Fourth ventricle: The fluid-filled cavity of the medulla oblongata of the brain.
- Frog: A caudal wedge-shaped part of the horse's hoof that lies between the bars.
- Fundic gland region: A portion of the glandular stomach lying between the cardiac and pyloric gland regions.

- Fungiform papilla: A mushroom-shaped elevation of the surface of the tongue.
- Gallbladder: A saclike diverticulum of the common bile duct; functions as a storage depot for bile.
- Ganglion cell layer (retina): Layer of cell bodies of neurons between the inner, plexiform layer and the nerve fiber layer.
- Gastric furrow: A depression in the mucosa of the glandular stomach lined by columnar surface mucous cells.
- Gastric pit (foveola): An invagination of the mucosa of the glandular stomach lined by columnar, surface mucous cells.
- Germinal center: The central region of an activated lymphatic nodule, consisting mainly of lymphocytes of medium to large size.
- Germinal epithelium: The epithelial layer covering the cortex of the ovary; composed of cuboidal or flattened cells.
- Germinal vesicle: The nucleus of an oocyte.
- Germinal zone (lens of eye): A band of epithelial cells located around the equator of the lens, capable of dividing throughout adult life.
- Gingiva: The gum surrounding a tooth.
- Glands of Moll: Sweat glands of the skin of the eyelids. Glands of the anal sac: Tubular glands in the wall of an anal sac
- Glands of Zeiss: Sebaceous glands of the skin of the eyelids.
- Glandular stomach: The portion of the stomach characterized by the presence of various types of tubular glands. It includes cardiac, fundic, body, and pyloric regions.
- Globular (globule) leukocyte: A leukocyte-like cell with large, round, eosinophilic granules, found in the mucosa of the intestine and stomach and reported to be derived from mast cell precursor cells in ruminants and rats.
- Glomerular epithelium: The layer of podocytes in intimate contact with the glomerular capillary loops of a renal corpuscle.
- Glomerulus: The tuft of capillary loops in the center of a renal corpuscle positioned between the afferent and efferent arteriole.
- Glycogen body: A structure, centrally located in the spinal cord of birds, whose cells contain a central mass of glycogen.
- Glycosaminoglycans (GAGS): Long-chained, very hydrophilic, sulfated polysaccharides. Most are directly bound to proteins and are collectively known as proteoglycans. Their hydrophilic property enables the diffusion of water-soluble material throughout the ground substance of loose connective tissue and cartilage.
- Goblet cell: A mucus-secreting cell having the shape of a goblet.
- Golgi apparatus: Membranous cellular organelle functioning to modify, sort, and package proteins secreted by a cell.
- Gonocyte: A primordial germ cell.
- Granulocyte: A leukocyte having specific granules, e.g., eosinophil, neutrophil, basophil, and heterophil.

- Granulosa lutein cell: The principal cell type of the corpus luteum; derived from a membrana granulosa cell of a postovulatory follicle.
- Gray matter: That portion of the brain and spinal cord containing the cell bodies of neurons and mainly unmyelinated neuronal fibers.
- Ground substance: The substance, consisting mainly of glycoproteins and glycosaminoglycans, that fills the spaces between the cellular and fibrous elements of connective tissue.
- Guttural pouch: A diverticulum of the Eustachian tubes of the horse.
- H&E: Hematoxylin and eosin.
- H band: A pale zone, devoid of actin filaments, located at the center of an A band of a sarcomere.
- Hair bulb: The expanded base of a hair follicle.
- Hair follicle: A derivative of the epidermis from which one or more hairs grow.
- Hair matrix: That portion of a hair bulb where cell division occurs, giving rise to a hair shaft
- Hard keratin: A tough form of keratin found in hair and nails, among other places; contains more disulfide bonds and cystine than soft keratin.
- Harderian gland: A large, tubular gland that lies on the dorsal posterior surface of the eye.
- Hassall's corpuscle: Acidophilic, concentric arrangements of reticular cells found in the medulla of the thymus of mammals.
- Haversian canal: The canal in the center of a Haversian system; contains blood vessels and nerves.
- Haversian system (osteon): Collectively, concentrically arranged bony lamellae, osteocytes, and a Haversian canal and its contents.
- **Head of the epididymis:** The initial portion of the epididymis.
- Helicine artery: A tortuous vessel that supplies blood to the cavernous spaces of the penis.
- Helicotrema: A tiny opening at the apex of the cochlear canal of the inner ear connecting the cavity of the scala vestibuli with the cavity of the scala tympani.
- Hemal node: A nodular structure found along blood vessels in ruminants; contains blood-filled sinuses between cellular cords.
- Hematoma: Blood that has escaped from a blood vessel(s) within a tissue or organ.
- Hemolymph node: A nodular structure, containing lymphatic vessels, whose sinuses receive a mixture of blood and lymph.
- Henle's loop: The U-shaped portion of a nephron located between its proximal and distal convoluted tubules.
- Hepatocyte: A liver cell.
- **Hepatoid gland:** The nonsebaceous portion of the circumanal glands of the dog. Its cells resemble hepatocytes, hence the name hepatoid.
- **Herbst corpuscle:** A tactile, encapsulated nerve ending of the skin of birds resembling the pacinian corpuscle but smaller.
- Herring body: A neurosecretion found within the axons of unmyelinated neurosecretory cells of the hypothal-amohypophyseal tract.

- Heterochromatic: The appearance of the nucleus of a fixed and stained cell whose chromatin is tightly coiled. Such chromatin stains readily, resulting in a grainy nucleus.
- **Heterophil:** The most abundant of the granulocytes of the chicken. Its specific granules are spindle-shaped and sometimes possess a distinct, ruby-red, spheric granule centrally.
- Hilus: An invaginated region of the margin of an organ where blood vessels enter and leave.
- Hilus cell: A group of epithelioid cells that may be found close to the rete ovarii near the hilus of the ovary in some mammals.
- Histiocyte: A synonym for macrophage.
- Holocrine secretion: A mode of secretion where disintegrated secretory cells and their product are released from a gland.
- Hoof: The keratinized, epidermal, digital outgrowth of an ungulate.
- Horn: A keratinized, epidermal outgrowth, with a bony core, of the head of some mammals.
- Horn tubule (tubular horn): A tapering, keratinized, epidermal structure of the equine hoof that extends toward the surface from a dermal papilla.
- Howship's lacuna: An eroded area of bone surface produced by the activity of an osteoclast.
- **Hyaline cartilage:** The most common form of cartilage whose matrix has the appearance of ground glass in the fresh state.
- **Hydroxyapatites:** Crystals of calcium phosphate that mineralize bone matrix, making it hard.
- Hypothalamohypophyseal tract: Collectively, the axons of neurosecretory cells within the infundibular stalk and infundibular process of the pituitary gland.
- I band: That portion of adjacent sarcomeres lying to either side of the Z-line of an uncontracted myofibril. The I band contains actin myofilaments exclusively.
- Immature bone: A highly cellular form of bone that is replaced by mature bone during development.
- **Incus:** The ossicle of the middle ear located between the malleus and stapes.
- Indeciduate placenta: A placenta whose endometrium and chorioallantoic membrane are in contact but do not fuse, seen in the mare, ruminants, and the sow
- **Infundibular cavity:** An extension of the third ventricle into the infundibular stalk of the pituitary, and in some animals, into the infundibular process.
- **Infundibular stalk:** That portion of the neurohypophysis between the median eminence and the infundibular process.
- **Infundibulum:** A ventral outpocketing of the diencephalon that develops into the neurohypophysis of the pituitary gland. Also, that part of the oviduct having a funnel shape and lying closest to the ovary.
- **Inner limiting membrane:** The part of the photosensitive retina of the eye abutting the vitreous body.
- **Inner nuclear layer:** The part of the photosensitive retina of the eye lying between the outer and inner plexiform layers.

Inner plexiform layer: The part of the photosensitive retina of the eye lying between the inner nuclear and ganglion cell layer.

Inner root sheath: That portion of a hair follicle lying between the hair and the external root sheath.

Inner tunnel: A large cavity lying within the organ of Corti of the inner ear.

Intercalated disc: A cell-surface modification found at both ends of cardiac muscle cells that enables cell-tocell adhesion and physiologic exchanges.

Intercalated duct: That part of the duct system of many glands that connects directly to the secretory unit.

Intermediate cell: An epithelial cell of the vagina with round corners, but larger than a parabasal cell, that occurs during the estrous cycle.

Internal ear: That part of the ear comprised of the semicircular ducts in semicircular canals, saccule, and utricle in the vestibule, and the cochlea.

Internal elastic membrane: A sheetlike elastic membrane forming the outer boundary of the tunica intima of many arteries.

Interstitial cell (Leydig cell): A cell that produces testosterone and is found in the connective tissue between seminiferous tubules of the testes.

Interstitial gland cells: Epithelioid cells, arranged as cords, in the stroma of the ovaries of bitches and queens but not ordinarily found in other domestic mammals.

Interstitial system (interstitial lamellae): A remnant of a preexisting Haversian system (osteon) of bone.

Interterritorial matrix: The matrix of hyaline cartilage surrounding the territorial matrix and possessing lesser concentrations of glycosaminoglycans than territorial matrix.

Intertubular horn: The keratinized epidermis of the hoof that surrounds the tubular horn.

Intralobular: That which is located within a lobule of a gland.

Intralobular duct: A duct located within the lobule of a gland.

Intramembranous bone (membrane bone): Bone that develops directly within or under a membrane of connective tissue.

Iris: The pigmented portion of the uvea surrounding the pupil of the eye.

Islet of Langerhans: A multicellular, endocrine structure embedded within the exocrine pancreas that produces insulin, glucagon, pancreatic polypeptide hormone, and somatostatin.

Isogenous group: A small group of cells derived by cell division from a single cell, as exemplified by isogenous groups of chondrocytes often seen in cartilage matrix.

Isthmus (of the oviduct): The section of the oviduct attached to the uterus.

Isthmus (of the uropygial gland): The part of the drainage system of the uropygial gland located between the primary duct and the nipple.

Juxtaglomerular apparatus: A trinity of cellular structures associated with a nephron and consisting of the macula densa, juxtaglomerular cells, and mesangial cells.

Juxtaglomerular cell: A modified cell (smooth muscle) of an afferent arteriole of the kidney.

Keratinized: A structure whose cells have become filled with the protein keratin.

Keratinocyte: A skin cell whose cytoplasm becomes filled with keratin, a scleroprotein, or albuminoid substance over time. Found in the dead cells of the outer layer of the epidermis, hairs, horns, feathers, hooves, beaks, etc.

Keratinoid: A tough proteinaceous substance lining the gizzard of the chicken.

Keratohyalin granules: Vesicles found within the cells of the stratum granulosum of the epidermis, whose contents will form the amorphous portion of keratin.

Köhler illumination: Bright, even illumination required for optimal light microscopy and achieved by adjusting the light source in a prescribed manner.

Krause's gland: An accessory lacrimal gland that may be either serous or mixed.

Kupffer cell: A macrophage located within the blood vascular system of the liver.

l.s.: Longitudinal section.

Labia: Lips.

Labial gland: Mixed glands found within the lips of the mouth.

Labyrinth: A complicated structural arrangement.

Lacrimal gland: The tear gland.

Lacuna: A small cavity or space.

Lagena: A terminal expansion of the cochlear duct of the inner ear of birds.

Lamella: A layer of material, e.g., of bone.

Lamina cribrosa: The sievelike part of the sclera that partitions groups of axons of the optic nerve.

Lamina propria: The loose connective tissue beneath the epithelium of a mucous membrane.

Lamina subglandularis: A thick sheet of collagenous fibers (stratum compactum) and the accompanying layer of fibroblasts (stratum granulosum) between the base of the glands and the muscularis mucosae of the stomach of the cat and sometimes the dog. The lamina may also occur in the small intestine of carnivores.

Laminar region: The part of the wall of the hoof that is composed of numerous plates (laminae) that function to suspend the third phalanx from the hoof.

Layer of rods and cones: In the retina of the eye, the layer comprised of the dendrites of the photoreceptor cells.

Lens: A biconvex, transparent structure comprised of lens fibers, positioned between the iris and the vitreous body.

Lens body: The part of the eye of the chicken, exclusive of the annular pad, whose lens fibers are oriented parallel to the optical axis of the eye.

Lens epithelium: The layer of simple cuboidal epithelium on the anterior surface (iris side) of the lens.

Lens fibers: Transparent, elongated, prismatic cells forming the bulk of the substance of the lens of the eye.

Leukocyte: A white blood cell, either granulocyte or agranulocyte.

Leydig cell: Synonym for the interstitial cell of the testis. Leydig cells produce testosterone.

- Ligamentum nuchae: A thick band of elastic tissue in the dorsal neck that is especially well developed in grazing animals.
- Light cell: Any cell, among contrasting dark cells, with pale cytoplasm; found in such places as the epithelium of the gallbladder or the tegmentum vasculosum of the inner ear of the chicken, among other places.
- Light zone (of the glandular stomach): The initial portion of the fundic gland region of the stomach of a carnivore; compared with the dark zone of the fundic gland region, it is shorter and its mucosa is thinner.
- Limbus: The boundary line between the cornea and sclera.
- **Lobule:** The subdivision of a lobe of an organ such as a gland or the lung.
- Loop of Henle: The U-shaped portion of a nephron joining a proximal and distal convoluted tubule.
- Loose (areolar) connective tissue: A type of connective tissue proper whose fibers are in the form of a non-compacted, open meshwork.
- Luteal cells (interstitial cells): The lutein cells of the corpus luteum of the mammalian ovary, or the pale interstitial cells within the theca externa surrounding the oocyte of an ovarian follicle of the chicken.
- Lymph node: A lymphatic organ having both afferent and efferent lymphatic vessels.
- Lymphatic nodule: A temporary, spherical, or oval structure, consisting of numerous lymphocytes, found within various lymphatic organs; may also be found anywhere within the loose connective tissue of the body.
- Lymphocyte: An agranulocyte with scant cytoplasm and, typically, a large, round or broadly oval nucleus.
- Macrophage: A phagocytic cell, derived from a monocyte, and widely distributed in tissues and organs throughout the body.
- Macula (of the ear): A patch of sensory and supporting cells found within the sacculus (saccule) and utriculus (utricle) of the inner ear.
- Macula densa: A part of the juxtaglomerular apparatus of a nephron and formed from closely packed epithelial cells of a portion of the wall of the distal convoluted tubule.
- Magnum: The part of the chicken's oviduct whose gland cells produce the albumin of the egg.
- **Malleus:** The middle ear ossicle in contact with the tympanic membrane.
- Mammary gland: The milk-producing organ of the mammalian female.
- Marginal zone: Splenic tissue located between the white and red pulp.
- Mast cell: A large, granular, ubiquitous cell of the connective tissue that produces histamine.
- Mature bone: Bone with an acidophilic matrix and fewer osteocytes than the immature bone it replaces during development.
- Median eminence: The region of the floor of the diencephalon of the brain from which the infundibular stalk of the pituitary arises.
- Medulla: The inner region of an organ such as the medulla of the kidney, ovary, or lymph node.

- Medullary cone: The cone-shaped medullary portion of the chicken's kidney that contains segments of uriniferous tubules.
- Medullary cord: One of many interconnected segments of diffuse connective tissue, surrounded by medullary sinuses, within the medulla of a lymph node.
- Medullary nephron: The larger of two forms of nephrons of the chicken's kidney having the characteristics of the mammalian nephron.
- Medullary sinuses: The lymph-filled spaces surrounding the medullary cords of a lymph node.
- Megakaryocyte: An extraordinarily large cell of the bone marrow that produces blood platelets by a budding process and releases them directly into the sinusoids of the marrow.
- Meissner's plexus: As seen in histologic section, a spindleshaped collection of parasympathetic neurons and their processes found within the submucosa of the digestive tract.
- Melanocyte: A large, branched cell that produces melanosomes (tiny vesicles containing pigment).
- Membrana granulosa: The cells that line the antrum of a mammalian ovarian follicle.
- Membrane bone: Bone that is formed within or beneath a membrane by intramembranous ossification, e.g., within embryonic mesenchyme or at the osteogenic surface of the periosteum or endosteum.
- Membranous labyrinth: A group of structures located within the bony labyrinth of the inner ear. The membranous labyrinth includes the semicircular ducts, the utricle and saccule, and the cochlear duct (membranous cochlea; scala media).
- Meninges: The three membranes enveloping both the brain and spinal cord: pia mater, arachnoid mater, and dura mater.
- Merkel's cell: A cell of the epidermis of the skin that has contact with tactile nerve endings.
- Merocrine gland (eccrine gland): An exocrine gland whose secretory cells release droplets of secretion by exocytosis.
- Mesangial cells: Phagocytic cells found within the renal glomerulus.
- Mesenchyme: Embryonal connective tissue that consists of stellate mesenchyme cells and ground substance.
- Mesobronchus: The intrapulmonary, primary bronchus of the chicken's lung.
- Mesometrium: The mesentery supporting the uterus from the abdominal wall.
- Mesosalpinx: The mesentery supporting the oviduct from the abdominal wall.
- Mesothelium: The mesodermally derived, simple squamous epithelium covering the surface of mesenteries and organs that protrude into coelomic cavities of the body.
- Metachromsia: The circumstance in which a cell or tissue component acquires a color different from the dye solution with which it is stained.
- Metamyelocyte: A developing granulocyte that possesses an indented nucleus and specific granules.

- Metestrus: That part of the estrous cycle between estrus and diestrus. The development of the corpus luteum occurs during metestrus.
- Microplacentome: A placental structure of the mare consisting of a small tuft of chorionic villi and a crypt of the endometrium into which it is inserted.
- Microvillus: A short, nonmotile, fingerlike projection of the free surface of certain epithelial cells. Usually present in large numbers, thereby greatly increasing the cell's surface area.
- Middle ear: A subdivision of the ear comprised of three small ossicles (malleus, incus, and stapes) or of a columella.
- Mixed gland: An exocrine gland whose secretory units consist of either mucous or serous cells or a combination of these cells. Or, a gland with both endocrine and exocrine components.
- Modiolus: The pillarlike bone in the center of the cochlea. Monocyte: A large, agranular leukocyte with an oval, indented or horseshoe-shaped nucleus and pale, bluegray, often vacuolated, cytoplasm.
- Mucosa: In the digestive tract, the mucous membrane comprised of an epithelium, lamina propria, and muscularis mucosae (latter is lacking in the mouth, pharynx, and portions of the esophagus). In other organ systems, the epithelium lining the organ together with the underlying lamina propria constitutes the mucosa.
- Mucous acinus: The bulblike secretory unit of a gland whose cells secrete mucus.
- Mucous connective tissue: A form of embryonal connective tissue consisting of amorphous ground substance, loosely arranged collagenous fibers, and fibroblasts.
- Mucous membrane: A synonym for mucosa.
- Mucous neck cell: A cell of the neck region of a gastric gland that produces mucus.
- Mucus: A viscous, slimy mixture of mucin, water, electrolytes, and cells. It is secreted by gland cells lining nasal, esophageal and other body cavities and serves to protect and lubricate surfaces.
- Multilaminar primary follicle: A preantral ovarian follicle whose oocyte is surrounded by several layers of follicle cells
- Multilocular adipocyte: An adipocyte (fat cell) containing numerous small lipid-filled vacuoles.
- Multinucleate giant cell: A large, phagocytic cell with many nuclei; formed by the coalescence of macrophages.
- Multipolar neuron: A nerve cell having numerous dendrites and a single axon.
- Muscularis externa: The outermost layers of muscle in the wall of the digestive tract; may be smooth muscle, skeletal muscle, or both.
- Muscularis mucosae: The layer(s) of smooth muscle below the lamina propria of the mucosa of the digestive tract.
- Myelin sheath: A derivative of the cell membrane of a Schwann cell or of an oligodendrocyte; the sheath is arranged in concentric layers around axons.
- Myeloblast: An early stage in the development of a granulocyte of the bone marrow.

- Myelocyte: The developmental stage of a granulocyte following the promyelocyte stage.
- Myocardium: The middle, muscular layer of the heart wall. Myoepithelial cell: An epithelial cell with contractile properties, as found on the surface of many glandular secretory units; also, one of the many contractile cells forming the iridial dilator of the eye.
- Myofibril: One of many contractile units, formed from linearly joined sarcomeres, as in skeletal and cardiac muscle cells.
- **Myofilament:** A linear subunit of a sarcomere made of the protein actin or myosin. Numerous myofilaments of actin and myosin become assembled to form the sarcomeres of myofibrils.
- Myoid cell: A contractile cell, such as found at the surface of a seminiferous tubule.
- Myometrium: The layers of smooth muscle external to the endometrium of the uterus.
- Nasal cavity: One of a pair of bilaterally arranged chambers located between the external nares and nasopharynx.
- Nasolabial gland: An exocrine gland of the subcutis of the planum nasolabiale of ruminants.
- Nasopharynx: That portion of the pharynx between the internal nares and the oropharynx.
- Necrotic: Pertaining to dead cells or tissue.
- Nephron: The portion of a uriniferous tubule before the collecting tubule; includes the renal corpuscle, proximal convoluted tubule, loop of Henle, and distal convoluted tubule.
- Nerve fiber layer (of the retina): The portion of the retina between the ganglion cell layer and the inner limiting membrane; consists of axons of ganglion cells.
- Neurilemma (neurolemma): In the peripheral nervous system, the sheath formed by Schwann cells that surround a neuron's axon. The neurilemma may or may not include a myelin component.
- Neuroglia: Supportive cells of the central nervous system; considered to be about 10 times more numerous than neurons. They include oligodendrocytes, astrocytes, microglia, and ependymal cells.
- Neurohypophysis: The portion of the pituitary gland that is derived from the infundibulum; consists of the median eminence, infundibular stalk, and infundibular process (pars nervosa).
- Neuromuscular spindle: An encapsulated, fusiform (spindle-shaped) stretch receptor found within skeletal muscle. It contains sensory and motor nerve endings and intrafusal fibers (modified muscle cells). Neuromuscular spindles regulate muscle tone.
- **Neutrophil:** A polymorphonuclear granulocyte with fine specific granules.
- Nictitating membrane: The third eyelid found in some animals.
- Nissl granule: The rough endoplasmic reticulum of a neuron.
- **Nonglandular stomach:** The part of the stomach lacking glandular elements.
- Nonsinusal spleen: A type of spleen having poorly developed sinuses or no sinuses, as found in the cat, horse, pig, and ruminants.

- **Nuclear bag fiber:** An intrafusal fiber (modified skeletal muscle cell) of a neuromuscular spindle, characterized by the presence of many closely packed nuclei.
- **Nucleolus:** A small round or oval structure, within the nucleus of a cell, where ribonucleoprotein is synthesized.
- Odontoblasts: Cells on the surface of dental papillae that produce uncalcified dentin (predentin).
- Olfactory epithelium: The pseudostratified columnar epithelium of the nasal cavity that is comprised of sensory cells, supporting cells, and basal cells.
- Oligodendrocyte: A highly branched neuroglial cell that forms the myelin sheath of axons within the central nervous system. By means of its many processes, a single oligodendrocyte can produce the myelin sheaths of many separate axons.
- Omasum: The third subdivision of the ruminant forestomach.
- Optic disc: The portion of the eye where the axons of the nerve fiber layer of the retina converge to form the optic nerve. There are no light sensitive cells present in the disc, hence the name blind spot.
- Optic nerve: The second cranial nerve.
- Ora ciliaris retinae: The point of transition from the photosensitive to the nonphotosensitive part of the retina.
- Organ of Corti: The part of the inner ear that is sensitive to sound.
- Oropharynx: The portion of the pharynx that is located behind the mouth.
- Orthochromatophilic erythroblast: In the erythroid line, the smallest nucleated cell.
- Os penis: A bone within the glans of the penis of carnivores.
- Osseous spiral lamina: A spiral shelf of bone around the modiolus of the cochlea.
- Osteoblast: A cell that synthesizes and secretes bone matrix.
- Osteoclast: A multinucleate giant cell that resorbs bone matrix.
- Osteocyte: A mature bone cell.
- Osteoid: Uncalcified bone matrix.
- Otolith: A tiny structure with a prismatic shape found embedded in the gelatinous covering (otolithic membrane) of a macula of the utricle and saccule of the inner ear.
- Otolithic membrane: The gelatinous covering of a macula of the utricle and saccule of the inner ear.
- Outer enamel epithelium: A layer of cells abutting (externally) the stellate reticulum of a developing fetal tooth.
- Outer limiting membrane: The retinal layer formed from the plasma membranes of Müller cells and located between the layer of rods and cones and the outer nuclear layer.
- Outer nuclear layer: The retinal layer consisting of the nuclei of the rod and cone cells.
- Outer plexiform layer: The retinal layer composed of neuronal fibers and located between the outer nuclear layer and the inner plexiform layer.

- Outer root sheath (external root sheath): The layer of cells in the wall of a hair follicle between the inner root sheath and the connective tissue sheath.
- Ovarian follicle: The cellular unit surrounding an oocyte in the ovary.
- Oviduct: The tubular organ that receives an oocyte from the ovary and conveys it, after fertilization, to the exterior (birds) or to the uterus (mammals, except prototherians) for implantation.
- Oxyphil: A parenchymal cell of the parathyroid gland. Pacinian corpuscle: An encapsulated nerve ending that
- responds to heavy pressure as opposed to light touch. Palpebral conjunctiva: The mucous membrane lining the
- Palpebral conjunctiva: The mucous membrane lining the eyelid.
- Paneth cell: A secretory cell whose acidophilic granules contain lysozyme; found in the intestinal glands of some mammals.
- Papillary duct: A large urinary duct that opens to the renal pelvis from the tip of a renal papilla of the kidney.
- Papillary (superficial) layer of the dermis: The upper layer of loose connective tissue of the dermis with fingerlike extensions called dermal papillae that interdigitate with the epidermis.
- Parabasal cell: A sloughed, small, round vaginal epithelial cell, found in vaginal smears taken from an animal in anestrus.
- Parabronchus (tertiary bronchus): An intrapulmonary branch of a secondary bronchus of the chicken lung.
- Parafollicular cell (C cell): A large, pale cell found between epithelial cells of thyroid follicles and also between follicles; produces the hormone calcitonin, whose action lowers blood calcium level.
- Parenchyma: The specific tissue cells of an organ as distinguished from the organ's supportive connective tissue, e.g., muscle cells as opposed to the connective tissue supporting them.
- Parietal cell: A large acidophilic cell of the fundic and pyloric gland regions of the stomach that produces hydrochloric acid.
- Pars ciliaris retinae: The bilayered, nonphotosensitive portion of the retina associated with the ciliary body and located between the ora ciliaris retinae and the pars iridica retinae.
- Pars convoluta (cortical labyrinth): That part of the renal cortex that contains renal corpuscles and convoluted tubules, and is located between medullary rays.
- Pars disseminata: The scattered portions of the prostate gland.
- Pars distalis: Derived from Rathke's pouch and the largest component of the pituitary gland. Alone or with the pars tuberalis, called the anterior lobe of the pituitary.
- Pars intermedia: That part of the pituitary gland located between the pars distalis and pars nervosa. A derivative of Rathke's pouch.
- Pars iridica retinae: That part of the nonphotosensitive retina located on the side of the iris facing the lens.
- Pars nervosa: The major part of the neurohypophysis and a derivative of the hypothalamus; with the pars intermedia, forms the posterior lobe of the pituitary.

Pars radiata (medullary or cortical rays): The part of the renal cortex alternating with the pars convoluta and consisting of collecting tubules and the straight portions of nephrons.

Pars tuberalis: The part of the pituitary gland that forms a collar around the infundibular stalk; derived from Rathke's pouch.

Pecten: A thin, vascular, pleated membrane that protrudes from the ventral surface of the chicken's eye into the cavity of the vitreous body.

Pectinate ligament (uveal meshwork): A loose network of elastic fibers, covered by squamous cells, that spans the filtration angle of the eye.

Penicillus: Term applied to the pulp arteries of the spleen and their branches because, collectively, they resemble the bristles of an artist's brush.

Penis, body of: Shaft of the intromittent organ of the male.

Penis, glans of: The expanded terminal end of the penis.

Periarterial lymphatic sheath: The white pulp of the spleen.

Pericardium: The visceral and parietal serosa of the pericardial cavity.

Perichondrium: The chondrogenic, dense irregular connective tissue covering of hyaline or elastic cartilage.

Perilymph: The fluid found in the bony labyrinth of the inner ear surrounding the membranous labyrinth.

Perimetrium: The serosa of the uterus.

Perimysium: The connective tissue surrounding a fascicle of muscle cells.

Perineurium: The connective tissue surrounding a bundle of nerve cell fibers (axons, dendrites, or both).

Perinuclear halo: A lightly stained area of cytoplasm separating the nuclear surface from the remainder of the cytoplasm in some small lymphocytes.

Periople: The proximal border of the horse's hoof.

Periosteum: The osteogenic, dense irregular connective tissue that covers portions of many bones.

Peritoneum: The serosa lining a coelomic cavity.

Perivitelline membrane: The membrane abutting the cell membrane of the oocyte of a chicken's ovarian follicle

Pessulus: A small bone supporting the syrinx of a chicken. Peyer's patch: An aggregation of lymphatic tissue (nodular and diffuse) in the lamina propria and submucosa of the small intestine, especially the ileum.

Photosensitive retina: The portion of the retina containing light-sensitive rod and cone cells.

Pia mater: The delicate, well vascularized meninx in contact with the surface of the brain.

Pigment epithelium: The pigmented layer of cells forming the outermost boundary of the retina.

Pineal gland: A dorsal evagination from the roof of the diencephalon of the brain.

Pinealocytes: The epithelioid, acidophilic, parenchymal cells of the pineal gland.

Pinna (auricle): The sound-collecting auricular appendage of the head.

Pituicytes: Neuroglial cells located among the neuronal fibers of the pars nervosa of the pituitary gland.

Placenta: A nutritive organ, derived in part from both the endometrium and the chorion.

Placentome: A structure formed from a cotyledon (clump of chorionic villi) and caruncle (elevation of the endometrium) of a cotyledonary placenta.

Planum: The flat surface of skin located between the external nares.

Plasma: The acellular fluid portion of circulating blood.

Plasma cell: A derivative of the B cell that synthesizes immunoglobulins.

Platelet: A fragment of membrane-bound cytoplasm, derived by budding from a megakaryocyte, with an important role in blood clotting.

Plexus: A localized network of any of the following: neurons and their processes, blood vessels, or lymphatic vessels.

Plica: A fold.

Podocyte: A highly branched cell whose processes interdigitate with those of other podocytes to form the glomerular epithelium of the Bowman's capsule of the kidney.

Polychromatophilic erythroblast: An erythroblast of medium size with cytoplasm exhibiting both basophilic and acidophilic areas.

Polymorphonuclear leukocyte: A granulocyte with a segmented nucleus.

Portal tract (area): An aggregation of blood vessels (and sometimes lymphatic vessels) and a bile ductule within the interlobular connective tissue of the liver.

Postcapillary venule: A venule, of the deep cortex of a lymph node, whose endothelial cells are cuboidal.

Posterior chamber (of the eye): The cavity, containing aqueous humor, that is located between the iris and the lens.

Posterior epithelium (of the cornea): The simple cuboidal or squamous epithelium covering the side of the cornea in contact with the aqueous humor.

Preantral follicle: A growing ovarian follicle that has not yet formed an antrum.

Predentin: Uncalcified dentin.

Prepuce: The foreskin of the penis.

Primary bronchus: A large bronchus branching directly from the trachea.

Primary follicle: An ovarian follicle whose oocyte is surrounded by a single layer of cuboidal cells. The term is also used, by some authors, for a multilaminar follicle.

Primary hair: A large hair shaft produced by a compound follicle.

Primary spermatocyte: A diploid cell formed by differentiation from a spermatogonium.

Primordial follicle: The earliest, smallest, and most numerous of ovarian follicles; consists of an oocyte surrounded by a layer of flat follicle cells.

Principal cell (chief cell): A parenchymal cell of the parathyroid gland. The name is also used for the small, basophilic cell of the glandular stomach that secretes pepsinogen.

Proctodeum: That part of the chicken's cloaca that joins the large intestine.

- Proerythrocyte (rubriblast): A large, round cell of the erythroid line with basophilic cytoplasm and a large round nucleus.
- **Proestrus:** The first stage of the estrous cycle; characterized by growth of the endometrium.
- **Promyelocyte:** An early granulocyte, recognized by a large nucleus with nucleoli and azurophilic cytoplasmic granules.
- **Proprioceptor:** A sensory receptor, mainly in skeletal muscles, tendons, and joints, that responds to stimuli arising within the body.
- Prostate gland: An accessory male reproductive gland whose secretion contributes to the seminal fluid at ejaculation.
- **Proventriculus:** The glandular portion of the stomach of the chicken.
- Proximal convoluted tubule: The long, highly convoluted tubule of a nephron that arises from a renal corpuscle and whose cells have a distinctive brush border.
- Pseudostratified epithelium: An epithelium that appears to be stratified but is not. All of its cells are in contact with the basement membrane. Its stratified appearance is the result of its cells being of different heights and their nuclei being located at different levels.
- Pulp artery: An artery within the red pulp of the spleen that arises from the central artery of the periarterial lymphatic sheath.
- Pupil: The opening in the center of the iris.
- Purkinje cell: A large, modified cardiac muscle cell that forms a part of the heart's conduction system. The word also identifies the large multipolar nerve cells present in the cerebellum at the junction of the granular and molecular layers.
- Pyknotic cell: A cell with a shrunken, basophilic nucleus or a cell that has become reduced in size.
- Pyloric gland region: The terminal glandular region of the stomach, characterized by deep gastric pits and mucous glands with some parietal cells.
- Pyramidal cell: A nerve cell having the shape of a pyramid, as found in the cerebral cortex.
- Ouill: The hard basal portion of a feather.
- Rachis: The part of the central shaft of a contour feather bearing the vanes.
- Rathke's pouch: An ectodermal diverticulum from the roof of the oral cavity of an embryo.
- **Red pulp:** The portion of the parenchyma of the spleen other than the white pulp; characterized by the abundance of erythrocytes.
- Reissner's membrane (vestibular membrane): The upper wall of the scala media of the inner ear separating it from the scala vestibuli.
- Renal corpuscle: A component of the nephron consisting of Bowman's capsule and the glomerulus.
- **Renal cortex:** The outer part of the kidney, identified by the presence of numerous renal corpuscles.
- Renal medulla: The region of the kidney internal to the cortex and dominated by loops of Henle, collecting tubules, and vasa rectae.
- Renal papilla: The tip of a renal pyramid.
- Renal pelvis: The expanded end of the ureter located within the hilus of the kidney.

- **Respiratory bronchiole:** A bronchiole with scattered alveoli within its walls and positioned between a terminal bronchiole and an alveolar duct.
- Rete ovarii: Channels, lined by cuboidal cells, located within the medulla of the ovaries of carnivores and ruminants.
- Rete testis: A network of channels located within the loose connective tissue of the mediastinum testis.
- Reticular fiber: A thin, argyrophilic, collagenous fiber.
- Reticular (deep) layer of the dermis: The layer of the dermis, consisting of dense irregular connective tissue, located below the papillary layer.
- Reticular structure: A diffuse form of Hassall's corpuscle in the thymus of the chicken, consisting of an irregular mass of reticular cells, including degenerating ones, in the medulla of lobules.
- Reticular tissue: A special form of connective tissue proper consisting of a feltwork of reticular fibers functioning as a supportive framework for cells of the parenchyma. Among other locations, it is well represented in the liver, spleen, and bone marrow.
- Reticulocyte: A newly produced erythrocyte.
- Retina (photosensitive): The part of the retinal tunic of the eye containing light-sensitive rod and cone cells.
- Retinal tunic: The innermost layer of the wall of the eye, consisting of the photosensitive retina, pars ciliaris retinae, and pars iridica retinae.
- Romanovsky stain: A compound dye substance used for staining the various different cells of the blood and bone marrow.
- Rouleau: An arrangement of erythrocytes, in a smear preparation, resembling a stack of discs or coins.
- Rumen: The largest subdivision of the forestomach of ruminants.
- Saccule (sacculus): A part of the membranous labyrinth within the vestibule of the inner ear; contains a macula whose sensory cells, when stimulated, make an animal aware of the position of its head in space and the sensations of linear acceleration and deceleration.
- Sarcolemma: The plasmalemma (cell membrane) of a muscle cell.
- **Sarcomere:** The unit of contraction of a skeletal or cardiac muscle myofibril; every sarcomere of a myofibril is located between adjacent Z lines.
- Sarcoplasm: The cytoplasm of a muscle cell.
- Satellite cell: A neuroglia cell in close proximity to the cell body of a neuron. Also, a cell with mesenchymal properties found intimately associated with skeletal muscle cells.
- Scala tympani: A chamber of the inner ear, filled with perilymph, lying below the floor of the cochlear duct.
- Scala vestibuli: A chamber of the inner ear, filled with perilymph, lying above the roof (vestibular membrane) of the cochlear duct.
- Schiff's reagent: A colorless fuchsin sulfurous acid (leucofuchsin) solution. Leucofuchsin is used in the periodic acid-Schiff (PAS) reaction for demonstrating the presence of complex carbohydrates, e.g., glycogen, starch, cellulose, and proteoglycans, among others.

- Schwann cell: A cell of the peripheral nervous system closely associated with neuronal processes and responsible for forming the neurilemma.
- Sclera: A part of the outermost tunic of the eye, consisting of dense irregular connective tissue.
- Scleral cartilage: A cup-shaped layer of cartilage within the sclera of the eye of the chicken.
- Scleral ossicle: A plate of bone, located anterior to the scleral cartilage, within the sclera of the eye of the chicken.
- **Scleral trabecular meshwork:** A webwork of elastic and collagenous fibers, within the filtration angle of the eye of the chicken, whose spaces are continuous with the spaces of Fontana of the uveal meshwork (pectinate ligament).
- Scleral venous plexus: A system of channels that drain excess aqueous humor away from the eye.
- Scute: A large scale covered by keratin.
- Sebaceous gland: A holocrine gland of the skin that produces an oily secretion.
- **Sebaceous zone:** The sebaceous region of the uropygial gland of the chicken.
- Secondary follicle: An ovarian follicle of the mammal with a C-shaped antrum
- Secondary spermatocyte: The stage of spermatogenesis between primary spermatocyte and early spermatid.
- **Secretory duct:** See striated duct.
- Semicircular canals: Tubular cavities within the temporal bone of the skull forming a subdivision of the bony labyrinth of the inner ear. They contain the semicircular ducts of the membranous labyrinth.
- Seminal vesicle: An accessory reproductive gland that contributes to the seminal fluid at ejaculation.
- Seminiferous tubule: A highly convoluted tubule within the testis where spermatogenesis occurs.
- Sensory cell: A neuron that conveys sensory information to the central nervous system.
- Sensory hair cells (ear): Cells, with stereocilia on their apical surfaces, that will initiate an impulse to the brain for interpretation when stimulated; found in the organ of Corti, sacculus, utricle, and crista ampullaris.
- **Serosa:** The lining membrane of the coelom in all its subdivisions; consists of a mesothelium and layer of underlying connective tissue.
- Serous demilune: A crescent-shaped cap, as seen in histologic section, of serous cells on the surface of a mucous acinus.
- Serous membrane: A synonym for serosa.
- Sertoli cell: A large, multifunctional cell, forming part of the epithelium of the wall of a seminiferous tubule.
- Serum: The acellular fluid part of blood obtained following coagulation.
- Sex cord: A convoluted cord of cells, within the testis of an embryo or young animal, whose cells will differentiate into stem cells.
- **Sharpey's fibers:** Collagenous fibers that anchor the periosteum to bone or a tendon to bone.
- **Sheathed artery:** A vessel of the white pulp of the spleen of the chicken that is surrounded by a ring of reticular cells.

- Shell gland: That portion of the oviduct of the chicken whose secretion produces the egg shell; the shell gland is also called the uterus.
- Simple epithelium: An epithelium consisting of a single layer of cells, all of which are attached to the basement membrane.
- Sinus hair: A tactile hair, limited to the facial region, that originates from a highly innervated follicle containing a large blood-filled sinus.
- Sinus pad: In the sinus hair follicle of the dog, the sinus pad is a thickening of the inner connective tissue sheath of the follicle that protrudes into the upper portion of the blood-filled sinus.
- Sinusal spleen: A spleen whose red pulp is characterized by numerous, blood-filled sinuses.
- Sinusoid: A thin-walled blood vessel with the characteristics of a capillary but having a larger and more irregular diameter.
- Small artery: Arbitrarily, an artery with up to eight or nine layers of smooth muscle cells in the tunica media. The smallest of the small arteries is called an arteriole.
- Smegma: The soft, cheesy deposit of desquamated epithelial cells and glandular secretions found on the glans and prepuce of the penis; it is also found in the urethral pouch of the stallion.
- Smudged cell: A ruptured, or otherwise distorted, leukocyte found in a blood or bone marrow smear.
- **Soft keratin:** A form of keratin, found in such places as the stratum corneum, that has fewer disulfide bonds and less cystine than hard keratin.
- **Spaces of Fontana:** Cavities filled with aqueous humor within the trabecular meshworks of the eve.
- Specific granules: Granules within the cytoplasm of a granulocyte that characterize the cell, e.g., eosinophilic granules of an eosinophil.
- Sperm-host gland: Tubular glands of the vagina of the oviduct of the chicken that serve as storage depots for deposited sperm.
- **Spermatid:** The haploid cell following the secondary spermatocyte during spermatogenesis.
- **Spermatogenic cell:** Any precursor of a spermatozoon found within the seminiferous epithelium.
- **Spermatogonium:** The most immature spermatogenic cell of the seminiferous epithelium
- **Sphincter** (iris): The circumferentially arranged smooth muscle of the iris that enables pupillary constriction.
- **Spicule:** A small, frequently irregularly shaped piece of bone.
- Spiral ganglion: The auditory ganglion associated with the modiolus of the cochlea.
- **Spiral ligament:** A thickening of the periosteal lining of the cochlear canal.
- Spiral limbus: An elevation of connective tissue resting on the osseous spiral lamina of the cochlea.
- **Spiral tunnel:** The space below the tectorial membrane of the inner ear.
- Spleen: A major lymphatic and blood-filtering organ.
- **Spongy bone** (cancellous bone): An architectural form of bone consisting of a three-dimensional meshwork of bony trabeculae containing numerous marrow spaces.

- Spur: A sharp, horny process on each leg of various birds.
 Stapes: The middle ear ossicle attached to the oval window in the petrous portion of the temporal bone.
- Stellate reticulum: A portion of the enamel organ of a developing tooth characterized by star-shaped cells.
- Stereocilia: Very long, nonmotile, flexible cell surface projections on the apices of certain cells, e.g., on the epithelial cells lining portions of the duct of the epididymis and on the sensory hair cells of the inner ear; referred to as long microvilli.
- Straight tubule: A tubule that connects a seminiferous tubule with the rete testis.
- Stratified epithelium: An epithelium consisting of two or more layers of cells with only the basal layer being in contact with the basement membrane.
- Stratum basale: The layer of cells in contact with the basement membrane of the epidermis.
- Stratum cavernosum: The erectile tissue of the pelvic urethra.
- **Stratum compactum:** A thick layer of collagenous fibers located between the base of the glands and the muscularis mucosae of the glandular stomach of the cat and some dogs.
- Stratum corneum: The outermost, keratinized layer of the epidermis.
- Stratum germinativum: The layers of the epidermis below the stratum corneum of the chicken, i.e., the basal, intermediate, and transitional layers. In mammals, stratum germinativum is a term sometimes employed to include both the stratum basale and the stratum spinosum.
- Stratum granulosum: The layer of cells below the stratum corneum whose cytoplasm contains keratohyalin granules.
- **Stratum intermedium:** The layer of cells located between ameloblasts and the stellate reticulum of a developing tooth.
- **Stratum lucidum:** The layer of pale epidermal cells located between the stratum corneum and stratum granulosum of thick skin.
- **Stratum medium:** The major portion of the wall of the horse's hoof, consisting of tubular and intertubular horn that extends from the coronary region to the surface of the ground.
- **Stratum spinosum:** The layer of the epidermis directly above the stratum basale. It is characterized by cells whose membranes have shrunken in all places except where desmosomes occur, giving the cells the appearance of having spines.
- Stratum vasculare: The richly vascularized and well innervated layer wedged between the inner circular and outer longitudinal layers of the myometrium of the bicornuate uterus of domestic mammals.
- Stria vascularis: The stratified cuboidal epithelium of the side of the cochlear duct that is attached to the spiral ligament of the cochlear canal. Capillaries occur among the superficial cuboidal cells of the stria.
- Striated border: The border of apical microvilli of intestinal epithelial cells as seen in profile view.

- Striated duct: An intralobular duct of a salivary gland whose epithelial cells have vertical stripes (invaginations of the plasma membrane) along their basal ends.
- Stroma (corneal): The predominant layer of the cornea; also called the substantia propria. It consists of lamellae of collagenous fibers oriented parallel to the corneal surface.
- Subarachnoid space: A space, filled with cerebrospinal fluid, located between the arachnoid layer and pia mater of the brain and spinal cord.
- Subcapsular sinus: The space, filled with lymph, beneath the capsule of a lymph node.
- Subcutis (subcutaneous connective tissue): The layer of loose connective tissue deep to the skin.
- Submucosa: The layer of connective tissue beneath a mucous membrane.
- Sulcus (chicken proventriculus): A depression between folds of the mucosa of the proventriculus.
- Superficial cell: Similar in size and shape to a superficial intermediate cell of a vaginal smear, but with a pyknotic nucleus, faded nucleus, or no nucleus.
- Superficial gland of the nictitating membrane: Depending on the type of animal, a serous, mucous, or mixed gland surrounding the base of the cartilage supporting the membrane.
- Superficial intermediate cell: A large, vaginal, epithelial cell with angular edges and a round nucleus found in vaginal smears.
- Surface mucous cells: Cells whose apical ends contain mucigen and form the epithelium of the glandular stomach of mammals. The mucus they secrete protects the epithelial surface from abrasion and the acidity of the stomach content.
- Sweat gland: A tubular or saclike gland of the skin; may be either apocrine or merocrine.
- Syndesmochorial placenta: The type of placenta of ewes and nanny goats where five layers of tissue separate maternal from fetal blood.
- Synovial fluid: The clear, lubricating fluid within the synovial cavity of a synovial joint.
- **Syntrophoblast:** The syncytial outer layer of the trophoblast.
- Syrinx: The voice box of the chicken, located where the trachea bifurcates into two bronchi.
- Taenia ceci: Flat bands of smooth muscle and elastic fibers within the cecum of horses and pigs.
- Taenia coli: Flat bands of smooth muscle and elastic fibers within the colon of horses and pigs.
- Tail of the epididymis: The end of the epididymis that joins with the vas deferens.
- **Tapetum lucidum:** A fibrous or cellular reflective layer of the choroid coat of the eye.
- Tarsal glands: Large multilobular sebaceous glands located within the tarsus (plate of dense connective tissue) of the palpebral conjunctiva.
- **Tarsus:** A plate of dense connective tissue within the eyelid. It is located between the dermis of the skin and the lamina propria of the palpebral conjunctiva.
- Taste bud: A multicellular, barrel-shaped structure, consisting of sensory and supportive cells and located within various parts of the epithelium of the tongue.

- Taste pore: A tiny opening at the tip of a taste bud.
- Teat canal: A channel, lined by stratified squamous epithelium, that opens onto the tip of a teat.
- Teat sinus: A channel, lined by a bistratified epithelium, that opens into a teat canal.
- **Tectorial membrane:** A proteinaceous membrane that overlies and contacts the stereocilia of the sensory cells of the organ of Corti of the inner ear.
- Tegmentum vasculosum: A vascularized membrane separating the cochlear duct from the overlying scala vestibuli of the inner ear of the chicken.
- **Tendon:** A bundle or band of dense regular connective tissue connecting a muscle to a bone.
- **Tendon sheath:** A layer of cells and fibers of connective tissue on the surface of a tendon.
- **Territorial matrix:** The matrix, rich in sulfated glycosaminoglycans, immediately surrounding a chondrocyte(s) of hyaline cartilage.
- Tertiary follicle: The large ovarian follicle just prior to ovulation; also called a Graafian follicle.
- Theca externa: The outer (connective tissue) layer of the wall of a mammalian ovarian follicle.
- Theca folliculi: A sheath of stromal cells, surrounding a growing, mammalian ovarian follicle, that will differentiate into a theca externa and theca interna.
- Theca interna: The inner cellular and well vascularized layer of the wall of a mammalian ovarian follicle.
- Theca lutein cell: A small lutein cell of a corpus luteum derived from a cell of the theca interna.
- Thick skin: Hairless skin with an epidermis that is many cells thick, e.g., skin of a digital pad or of the planum nasolabiale.
- **Thin skin:** Skin with an epidermis that is only a few cells thick, e.g., skin of the trunk or of the legs among other places.
- **Thrombocyte:** A blood cell of the chicken with a role in blood clot formation. Also, incorrectly used as a synonym for a platelet of mammals.
- Thyroid follicle: A vesicle of the thyroid gland, formed of a simple epithelium, that contains the storage form of thyroxin called thyroglobulin.
- Tomial edge of beak: The cutting edge of a bird's beak. Tonsil: A lymphatic organ found below the epithelium of the pharynx. It is composed of lymphatic nodules and diffuse lymphatic tissue.
- **Trabecula:** A part of the framework of connective tissue of an organ or structure, e.g., a bundle of fibers of the splenic stroma or any of the irregularly shaped pieces of bone that form part of the three-dimensional latticework of spongy bone.
- Transitional epithelium: An epithelium, limited to the urinary system, whose appearance depends on the amount of fluid pressure applied against it.
- **Trophoblast:** The mesectodermal layer covering the blastocyst.
- Tubuloacinar gland: A gland whose secretory units consist of tubules and acini.
- Tunica adventitia: The outermost layer of connective tissue of a blood vessel.
- Tunica albuginia: A layer of dense connective tissue surrounding a structure, e.g., the layer beneath the germi-

- nal epithelium of the ovary or the layer surrounding the testis.
- Tunica intima: The innermost tunic of the wall of a blood vessel.
- Tunica media: The middle, muscle layer of the wall of a blood vessel.
- Tunica vaginalis: The serosa of the testis and the epididymis.
- Tympanic cavity: The cavity containing the middle ear ossicles; also called the cavity of the middle ear.
- Tympanic membrane (eardrum): The membrane between the external auditory meatus and the tympanic cavity (cavity of the middle ear).
- Type I alveolar cells: Simple squamous epithelial cells that line the alveoli of the lungs and form part of the alveolar septa.
- Type II alveolar cell: A cell present in the alveolar lining of the mammalian lung that produces surfactant. Surfactant reduces surface tension.
- Type I cell (of the macula): A chalice-shaped sensory cell of the epithelium of the macula of a sacculus.
- **Unilocular adipocyte:** An adipocyte (fat cell) containing a single, very large, lipid-filled vacuole.
- Unipolar neuron: A nerve cell with two processes that arise from a single site on the surface of the nerve cell.
- **Urachus:** The portion of the reduced allantoic stalk between the apex of the bladder and the umbilicus (belly button).
- **Ureter:** The tube extending from the renal pelvis to the urinary bladder.
- **Urethra:** The tube extending from the urinary bladder to the exterior.
- **Urethral process:** An extension of the urethra beyond the penis as in the stallion and ruminants.
- Urinary space: The cavity between the capsular epithelium and the glomerular epithelium of a renal corpuscle.
- **Urodeum:** The portion of the cloaca of a chicken into which urinary wastes are deposited.
- **Uropygial gland** (preen gland): A holocrine gland that produces an oily secretion and is located dorsally within the base of the tail of the chicken.
- Uterine gland: A simple, tubular gland within the endometrium of the uterus.
- Utricle (utriculus): Function same as for saccule.
- **Uveal trabecular meshwork:** One of three meshworks of connective tissue, within the filtration angle of the eye, whose cavities are filled with aqueous humor.
- Vacuolar cell: A cell of the cortex of the chicken ovary containing numerous fat vacuoles and a pyknotic nucleus. Collections of these cells may represent the remnants of a postovulatory follicle.
- Vagina (of the chicken oviduct): The segment of the oviduct that opens into the urodeum of the cloaca.
- **Valve:** A leaflet of connective tissue, covered by endothelial cells, that assures fluid flow in one direction, as in the heart, veins, and lymphatic vessels.
- Vane (of a feather): The most prominent feature of a contour feather. It is formed of barbs and interlocking barbules positioned on opposite sides of the rachis.

- Vas deferens: The sperm duct that extends from the testis to the urethra.
- Vasa vasorum: Blood vessels within the wall of a blood vessel.
- Vasa recta: Straight, thin-walled, large-diameter blood vessels located within the medulla of the kidney.
- Vascular tunic (uvea): The middle layer of the wall of the eye.
- Vascular layer (of choroid): The portion of the choroid layer of the eye that contains numerous blood vessels.
- Ventral root: The motor portion of a spinal nerve that arises from the ventrolateral part of the spinal cord.
- Venule: A small, thin-walled vein.
- Vestibular membrane: A thin, epithelial membrane separating the cochlear duct from the scala vestibuli of the inner ear.
- Vestibule: An entrance chamber, e.g., of the nose or vulva. Vestibule of the inner ear: A central cavity within the temporal bone of the skull forming a subdivision of the bony labyrinth of the inner ear. It contains the saccule and utricle of the membranous labyrinth.
- Villus: A finger-shaped process, e.g., intestinal villus.
- Visceral pleura: The serous membrane covering the surface of the lung.
- **Vocal ligament:** A band of elastic fibers enclosed in a fold of a mucous membrane.
- Volkmann canal: In the diaphysis of a long bone, any transverse channel connecting two Haversian canals or piercing the shaft of the bone.
- Vulva: The external genitalia of the female mammal; it includes the vestibule, labia, and clitoris.
- Wall (of hoof): The part of the hoof that is visible when the digit is on the ground.
- Wattle: A fleshy appendage of the skin as in the throat region of the neck of a chicken, goat, or pig.
- White line: The junction, at the surface of the ground, of the wall and sole of the horse's hoof.
- White matter: The part of the brain or spinal cord containing numerous myelinated neuronal processes.

- White pulp: The scattered but numerous concentrations of diffuse and nodular lymphatic tissue found throughout the red pulp of the spleen.
- x.s.: Cross section.
- **Z** band (line): The boundary between adjacent sarcomeres of a myofibril; structurally, it represents the point at which the actin filaments of adjacent sarcomeres are in contact with the Z filament.
- Zona fasciculata: The thickest portion of the adrenal cortex; located between the zona glomerulosa or zona intermedia and the zona reticularis.
- Zona glomerulosa (zona multiformis): The outermost portion of the adrenal cortex.
- Zona intermedia: The portion of the adrenal cortex located between the zona glomerulosa and the zona fasciculata.
- Zona pellucida: An acidophilic membrane separating an oocyte from the cumulus oophorus.
- Zona reticularis: The innermost portion of the adrenal cortex located between the zona fasciculata and the adrenal medulla.
- **Zonary placenta:** A placenta, found in carnivores, that is wrapped around the chorionic sac in the manner of a cummerbund.
- **Zone of calcification:** The portion of the epiphyseal disc where the cartilage matrix becomes infiltrated by calcium salts.
- **Zone of hypertrophy:** The portion of an epiphyseal disc where the chondrocytes become enlarged.
- Zone of multiplication (proliferation): The portion of an epiphyseal disc where chondrocytes are duplicated.
- **Zone of ossification:** The portion of an epiphyseal disc where cartilage is being replaced by bone.
- **Zone of reserve cartilage:** The portion of an epiphyseal disc that is attached to the bone of the epiphysis.
- Zonular fibers (suspensory ligaments): Collagenous suspensory fibers that extend from the capsule of the lens to the ciliary processes.

BIBLIOGRAPHY

- Adam, W.S., Calhoun, M.L., Smith, E.M., and Stinson, A.W.: Microscopic Anatomy of the Dog: A Photographic Atlas. Springfield, IL, Charles C. Thomas, 1970.
- Amann, R.P., Johnson, L., and Pickett, B.W.: Connection between the seminiferous tubules and the efferent ducts in the stallion. Am. J. Vet. Res., 38:1571,
- Amoroso, E.C.: Histology of the placenta. Br. Med. Bull., 17:81, 1961.
- Arey, L.B.: Human Histology. 4th Ed. Philadelphia, W.B. Saunders, 1974.
- Banks, W.J.: Applied Veterinary Histology. 2nd Ed. Baltimore, Williams & Wilkins, 1986.
- Björkman, N.H.: Fine structure of the fetal-maternal area of exchange in the epitheliochorial and endotheliochorial type of placentation. Acta Anat., 86(Suppl. 1):1, 1973.
- Bloom, W., and Fawcett, D.W.: A Textbook of Histology. 10th Ed. Philadelphia, W.B. Saunders, 1975.
- Borror, D.J.: Dictionary of Word Roots and Combining Forms. Mountain View, California, Mayfield Publishing Co., 1960.
- Calhoun, M.L.: The microscopic anatomy of the digestive tract of Gallus domesticus, Iowa State Coll. I. Sci., 7:261, 1933.
- Cole, H.H., and Cupps, P.T. (eds.): Reproduction in Domestic Animals. Vol. 1. New York, Academic Press, 1959.
- Cole, H.H., and Cupps, P.T. (eds.): Reproduction in Domestic Animals. New York, Academic Press, 1969.
- Czarnecki, C.M., and Hammer, R.F.: Auto-tutorial Laboratory Guide. Microscopic Anatomy of Domestic Animals. St. Paul, MN, University of Minnesota, 1978.
- Dellmann, H.: Textbook of Veterinary Histology. 4th Ed. Philadelphia, Lea & Febiger, 1993.
- Delly, J.G.: Photography Through the Microscope. 7th Ed. Rochester, NY, Eastman Kodak, 1980.
- Department of Anatomy: Histology Laboratory Manual (Microscopic Anatomy, VAN. 306). Ames, IA, Iowa State University, 1987
- Di Fiore, M.S.H.: Atlas of Human Histology. 5th Ed. Philadelphia, Lea & Febiger,
- Elias, H.: Comparison of duodenal glands in domestic animals. Am. J. Vet. Res., 8:311, 1947
- Evans, H.E., and Christensen, G.C.: Miller's Anatomy of the Dog. 2nd Ed. Philadelphia, W.B. Saunders, 1979.
- Gartner, L.P., and Hiatt, J.L.: Atlas of Histology. Baltimore, Williams & Wilkins,
- Geneser, F.: Color Atlas of Histology. Philadelphia, Lea & Febiger, 1989.
- Gentle, M.J.: The lingual taste buds of Gallus Domesticus L. Br. Poult. Sci., 12:245, 1971.
- Getty, R.: Sisson and Grossman's The Anatomy of the Domestic Animals. Vol. 1. 5th Ed. Philadelphia, W.B. Saunders, 1975
- Ham, A.W., and Cormack, D.H.: Histology. 8th Ed. Philadelphia, J.B. Lippincott,

- Hammersen, F.: Sabotta/Hammersen. Histology. A Color Atlas of Microscopic Anatomy. 3rd Ed. Baltimore, Urban & Schwarzenberg, 1985.
- Hamre, C.J.: Origin and differentiation of heterophil, eosinophil and basophil leukocytes of chickens. Anat. Rec., 112:339, 1952.
- Hodges, R.D.: The Histology of the Fowl. New York, Academic Press, 1974.
- Humason, G.L.: Animal Tissue Technique. 4th Ed. San Francisco, W.H. Freeman and Co, 1979.
- Huntley, J.F.: Mast cells and basophils: A review of their heterogeniety and function. J. Comp. Path., 107:349, 1992.
- Jain, N.C.: Schalm's Veterinary Hematology. 4th Ed. Philadelphia, Lea & Febiger, 1986.
- Junqueira, L.C., and Carneiro, J.: Basic Histology. 2nd Ed. Los Altos, CA, Lange Medical Publications, 1980.
- Kelly, D.E., Wood, R.L., and Enders, A.C.: Bailey's Textbook of Microscopic Anatomy. 18th Ed. Baltimore, Williams & Wilkins, 1984.
- Lawson, D.D., Nixon, G.S., Noble, H.W., and Weipers, W.L.: Dental anatomy and histology of the dog. Res. Vet. Sci., 1:201, 1960.
- Leeson, T.S., and Leeson, C.R.: A Brief Atlas of Histology. Philadelphia, W.B. Saunders, 1979.
- Lewis, H.B., and Rebar, A.H.: Bone Marrow Evaluation in Veterinary Practice. St. Louis, Ralston Purina, 1979.
- Lindenmaier, P., and Kare, M.R.: The taste end-organs of the chicken. Poult. Sci., 38:545, 1959
- Lucas, A.M., and Jamroz, C.: Atlas of Avian Hematology. Washington, D.C., United States Department of Agriculture, 1961.
- Lucas, A.M., and Stettenheim, P.R.: Avian Anatomy. Integument. Part I. Washington, D.C., United States Department of Agriculture, 1972.
- Lucas, A.M., and Stettenheim, P.R.: Avian Anatomy. Integument. Part II. Washington, D.C., United States Department of Agriculture, 1972.

 Melfi, R.C.: Permar's Oral Embryology and Microscopic Anatomy. 7th Ed.
- Philadelphia, Lea & Febiger, 1982.
- Mescher, A.L.: Junqueira's Basic Histology Text and Atlas. 12th Ed. New York. McGraw-Hill 2010.
- Murphy, C.J.: Raptor ophthalmology. Compend. Contin. Educ. Pract. Vet., 9:241, 1987.
- Natt, M.P., and Herrick, C.A.: Variation in the shape of the rodlike granules of the chicken heterophil leukocyte and its possible significance. Poult. Sci.,
- Nickel, R., Schummer, A., and Seiferle, E. (eds.): Anatomy of the Domestic Birds. Translated by W.G. Siller and P.A.L. Wight. New York, Springer-Verlag, 1977
- Olson, P.N., Behrendt, M.D., and Weiss, D.E.: Reproductive problems in the bitch: Finding answers through vaginal cytology. Vet. Med., 82:344, 1987.
- Prince, J.H., Diesem, C.D., Eglitis, I., and Ruskell, G.L.: Anatomy and Histology of the Eye and Orbit in Domestic Animals. Springfield, IL, Charles C. Thomas,
- Rebar, A.H.: Handbook of Veterinary Cytology. St. Louis, Ralston Purina,

- Reith, E.J., and Ross, M.H.: Atlas of Descriptive Histology. 3rd Ed. New York, Harper & Row, 1977.
- Rich, L.J.: The Morphology of Canine and Feline Blood Cells, Including Equine References. St. Louis, Ralston Purina, 1976.
- Ross, M.H., and Romrell, L.J.: Histology. 2nd Ed. Baltimore, Williams & Wilkins, 1989.
- Ross, M.H., Romrell, L.J., and Kaye, G.I.: Histology. A Text and Atlas. 3rd Ed. Baltimore, Williams & Wilkins, 1995.
- Samuel, C.A., Allen, W.R., and Steven, D.H.: Studies on the equine placenta. I. Development of the microcotyledons. J. Reprod. Fertil., 41:441, 1974.
 Seely, J.C.: The harderian gland. Lab. Anim., 16:33, 1987.
- Shackleford, J.M., and Wilborn, W.H.: Ultrastructure of bovine parotid glands. J. Morphol., 127:453, 1969.
- Sloss, M.W.: The microscopic anatomy of the digestive tract of Sus scrofa domestica. Am. J. Vet. Res., 1:578, 1954.
- Smith, R.F.: Microscopy and Photomicrography. A Practical Guide. New York, Appleton-Century-Crofts, 1982.
- Stedman, T.L.: Illustrated Stedman's Medical Dictionary. 24th Ed. Baltimore, Williams & Wilkins, 1983.
- Strickland, J.H., and Calhoun, M.L.: The integumentary system of the cat. Am. J. Vet. Res., 24:1018, 1963.

- Stump, J.E.: Anatomy of the normal equine foot, including microscopic features of the laminar region. J. Am. Vet. Med. Assoc., 151:1588, 1967.
- Sturkie, P.D. (ed.): Avian Physiology. 3rd Ed. New York, Springer-Verlag, 1976.
- Telford, I.R., and Bridgman, C.F.: Introduction to Functional Histology. 2nd Ed. New York, Harper and Collins, 1995.
- Trautmann, A., and Fiebiger, J.: Fundamentals of the Histology of Domestic Animals. Translated and revised by R.E. Habel and E.L. Biberstein. Ithaca, NY, Comstock, 1952.
- Trautmann, A., and Fiebiger, J.: Fundamentals of the Histology of Domestic Animals. Translated and revised by R.E. Habel and E.L. Biberstein. Ithaca, NY, Comstock, 1957.
- Webb, A.J., and Calhoun, M.L.: The microscopic anatomy of the skin of mongrel dogs. Am. J. Vet. Res., 15:274, 1954.
- Weiss, L.: Cell and Tissue Biology, A Textbook of Histology. 6th Ed. Baltimore, Munich, Urban & Schwarzenberg, 1988.
- Weiss, L.: The red pulp of the spleen: Structural basis of blood flow. Clin. Haematol., 12:375, 1983.
- Weiss, L., and Greep, R.O.: Histology. 4th Ed. New York, McGraw-Hill, 1977. Wheater, P.R., Burkitt, H.G., and Daniels, V.G.: Functional Histology. A Text and

Colour Atlas. New York, Churchill Livingstone, 1987.

310

INDEX

Page numbers in italics denote figures.

```
A band, 62
A cell, 176, 177, 182
Abomasum, 160, 162
Accessory glands, 226
Acidic (anionic) stains, 5
Acidophil, 212, 216, 217, 222
Acinar cell, 176
Acinus, 176, 177, 182, 278
Adenohypophysis, 211, 213
Adipocyte (adipose cell), 20, 24, 25, 165
Adipose tissue
  cardiovascular system, 81, 82, 84, 85, 86, 87
  connective tissue, 20, 21, 25
  digestive system, 145, 148, 151, 153, 154, 160, 165,
    169, 170
  ear, 286
  endocrine system, 218, 223
  female reproductive system, 250
  integument, 117, 118, 135, 136, 138
  lymphatic system, 98, 103
  male reproductive system, 238
  nervous system, 72
  respiratory system, 199, 200, 201
  urinary system, 189
Adrenal cortex, 212
Adrenal gland, 212, 213, 220, 221, 223
Adrenal medulla, 213
Adventitia
  digestive system, 139, 153, 154, 175
  female reproductive system, 259
  male reproductive system, 234
  tunica adventitia, 77, 80, 81, 82, 83, 85, 252
  urinary system, 191
Afferent arteriole, 188, 189
Agranulocyte, 42, 43
```

```
Air capillary, 197, 209
Air sac, 197
Air space, 133
Air vesicle (atrium), 197, 208, 209
Albumen, 264
Allantoic blood vessel, 254, 255, 256, 257, 258
Allantoic epithelium, 254
Alpha cell. See Acidophil
Alveolar bone, 143, 144
Alveolar (type II) cell, 204
Alveolar duct, 203
Alveolar sac, 203
Alveolar septum, 204
Alveolus, 195, 202, 203, 204, 205
Ameloblast, 144
Amorphous ground substance, 19, 22, 27
Ampulla
  ear, 284, 288
  female reproductive system, 244, 250
  male reproductive system, 226, 234
Anal canal, 140, 170, 172
Anal gland, 140, 170, 171
Anal sac, 141, 170
Anastomotic artery, 83, 176
Anestrus, 244, 245, 259, 260
Anionic (acidic) stains, 5
Annular ligament, 287, 288
Annular pad, 269, 279
Annular sinus, 106, 119
Anterior chamber, 269, 271, 272, 276, 279
Anterior epithelium, 267, 274
Antrum, 244, 247, 248
Aorta, 83, 84, 85
Aortic body, 86, 87
Apical cell, 265
```

| Apocrine tubular gland, 106, 171 | Basophilic band cell, 56 |
|--|--|
| Arachnoid layer, 68 | Basophilic erythroblast, 54, 55, 56 |
| Areolar (loose) connective tissue | Basophilic myelocyte, 55, 56 |
| defined, 20, 22, 23, 24 | Basophilic substances, 5 |
| digestive system, 145 | Beta cell (basophil) |
| eye, 281 | blood, 42, 43, 44, 45, 46, 47, 52 |
| male reproductive system, 232, 233 | endocrine system, 212, 217, 222 |
| Argyrophilic fiber, 20 | Bifurcation, 63 |
| Arrector pili muscle, 106, 114 | Bile canaliculus, 141, 173 |
| Arteriole | Bile duct, 173, 174 |
| cardiovascular system, 77, 79, 80, 86, 87 | Binocular microscope, 8 |
| connective tissue, 24 | Binucleate hepatocyte, 174 |
| digestive system, 155, 162 | Bipolar neurons, 65 |
| lymphatic system, 90 | Bistratified epithelium, 13, 17, 18 |
| muscle, 60 | bistratified columnar epithelium, 124, 125 |
| Arteriovenous anastomoses, 78, 83 | bistratified cuboidal epithelium, 124, 125 |
| Arteriovenous shunt, 75 | Bitch. See Dog |
| Artery, 39, 81, 87, 252 | Blind spot, 269 |
| Articular cartilage, 37, 287 | Blood, 41–52 |
| Articulating surface, 40 | agranulocyte, 42, 43 |
| Artifact (imperfections) | azurophilic granule, 42, 49 |
| crackling artifiact, 10 | basophil (beta cell), 42, 43, 44, 45, 46, 47, 52 |
| defined, 9, 10 | buffy coat, 45, 50 |
| digestive system, 144, 148, 164 | central pallor, 41 |
| endocrine system, 219 | chicken, 43, 51–52 |
| eye, 274, 280 | crenated erythrocyte, 42 |
| female reproductive system, 257 | eosinophil, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 |
| nervous system, 73, 76 | erythrocyte (red blood cell), 41, 43, 44, 45, 46, 47, |
| space artifact, 73, 76, 148, 219, 274, 280 | 48, 49, 50, 51 |
| Arytenoid cartilage, 199, 200 | formed elements, 41 |
| Astrocyte, 69 | granulocyte, 42, 43, 51 |
| Atresia, 244 | heterophil, 43, 51, 52 |
| Atretic follicle, 246, 262 | leukocyte (white blood cell), 41, 42, 43, 44 |
| Atrium, myocardium, 85 | lymphocyte, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51 |
| Atrium (air vesicle), 197, 208, 209 | mammals, 41–43 |
| Attachment epithelium, 144 | monocyte, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 |
| Auerbach's plexus, 72, 167 | neutrophil, 42, 44, 45, 46, 47, 48, 49, 50 |
| Auricle (pinna), 29, 188, 283 | plasma, 41, 44 |
| Axial blood vessel, 132 | platelet, 41, 43, 44, 45, 46, 47, 49, 50 |
| Axon, 65, 73, 74, 76 | polymorphonuclear leukocyte, 43 |
| Axon hillock, 71 | rouleaux, 42, 45, 47 |
| Azurophilic granule, 42, 49, 55 | serum, 41 |
| | smudged cell, 44, 45, 47, 52 |
| B cell, 176, 177, 182 | thrombocyte, 43, 51 |
| Band cell, 54, 55, 56 | vacuole, 43, 51 |
| Bar, hoof, 128, 129 | word roots, 43 |
| Barb, 107, 132 | Blood vessel |
| Barbule, 107, 133 | allantoic blood vessel, 254, 255, 256, 257, 258 |
| Basal cell | axial blood vessel, 132 |
| digestive system, 179 | cardiovascular system, 103 |
| female reproductive system, 265 | digestive system, 144, 151, 152 |
| integument, 106, 131, 134, 138 | ear, 290 |
| male reproductive system, 233, 234, 235 | endocrine, 216 |
| respiratory system, 198, 205, 206 | eye, 275, 281 |
| Basal striations, 149 | female reproductive system, 253, 265 |
| Basement (glassy) membrane, 11, 16, 106, 121 | integument, 118, 124, 125, 131 |
| Basic (cationic) stain, 5 | lymphatic system, 103 |
| Basilar membrane, 284, 289, 290 | maternal blood vessel, 255, 256, 257 |
| Basophil (beta cell) | mesenteric blood vessel, 75 |
| blood, 42, 43, 44, 45, 46, 47, 52 | nervous system, 67, 74 |
| endocrine system, 212, 217, 222 | Boar. See Pig |

| | 1 1 1 1 1 1 2 |
|--|--|
| Body of penis (corpus penis), 227, 238, 239 | basophilic band cell, 56 |
| Body of prostate gland, 226, 235 | basophilic erythroblast, 54, 55, 56 |
| Bone, 31–40 | basophilic myelocyte, 55, 56 |
| artery, 39 | chicken, 54 |
| articular cartilage, 37 | eosinophil, 56 |
| articulating surface, 40 | eosinophilic band cell, 55 |
| bone marrow, 35, 38 | eosinophilic myelocyte, 55 |
| bone matrix, 31, 32, 33, 34, 36, 37, 38, 40 | erythrocyte, 55, 56 |
| calcified cartilage, 34, 36, 37, 38 | erythroid, 53 |
| canaliculi, 32, 38 | granulocyte, 53 |
| cartilage, 40 | hematopoietic tissue, 53 |
| circumferential lamellae, 32 | heterophil, 56 |
| | |
| compact bone, 32, 38, 39 | mammals, 53 |
| concentric lamellae, 32 | megakaryocyte, 54, 55 |
| distal interphalangeal joint, 40 | metamyelocyte, 54 |
| ear, 286 | mitotic figure, 55 |
| endochondral bone, 34 | myeloblast, 54 |
| endochondral (intracartilaginous) ossification, 32, 38 | myelocyte, 54 |
| endosteum, 31, 39 | neutrophil, 56 |
| epiphyseal disc, 32, 37 | neutrophilic band cell, 55, 56 |
| extensor tendon, 40 | neutrophilic metamyelocyte, 56 |
| fibrous capsule, 40 | orthochromatophilic erythroblast, 54, 55, 56 |
| haversian canal, 32, 38, 39 | osteoblast, 54, 55 |
| haversian systems, 32, 38 | osteoclast, 54, 55 |
| hyaline cartilage, 34, 35 | plasma cell, 55 |
| immature bone, 32, 33 | pluripotent stem cell, 53 |
| integument, 130, 131, 135, 137 | polychromatophilic erythroblast, 54, 56 |
| interstitial system, 38 | proerythrocyte, 53 |
| intramembranous ossification, 32, 33 | promyelocyte, 54, 56 |
| joint cavity, 40 | reticulocyte, 54, 56 |
| lacuna, 32, 33, 38 | smudged cell, 55 |
| lamellae, 32 | vascular sinusoid, 53 |
| marrow cavity, 31, 34, 35, 36, 37, 39 | word roots, 54 |
| mature bone, 32, 33 | Bone matrix, 31, 32, 33, 34, 36, 37, 38, 40 |
| | |
| megakaryocyte, 36, 39 | Bony tracked sing 200, 207 |
| nerve, 39 | Bony tracheal ring, 206, 207 |
| nervous system, 67, 76 | Bowman's capsule, 184, 187, 188, 194 |
| osteoblast, 31, 33, 36, 37, 38 | Bowman's gland, 198, 206 |
| osteoclast, 32, 33, 36, 38 | Bowman's membrane, 267, 280 |
| osteocyte, 32, 33, 36, 37, 38, 39 | Brain sand, 69 |
| osteoid, 32, 33 | Bridge, 270, 281 |
| osteon, 32, 38 | Bronchi, 195, 202, 204 |
| periosteum, 31, 34, 35, 37 | Bronchial cartilage, 208 |
| periosteum of femur, 39 | Bronchial rings, 207 |
| respiratory system, 198, 209 | Bronchial tree, 195, 196 |
| Sharpey's fiber, 39, 40 | Bronchiole, 196, 202, 203, 204 |
| sinusoid, <i>36</i> , <i>38</i> | Bruch's membrane, 268 |
| spongy bone, 32, 33, 37 | Brücke's muscle, 279 |
| synovial fold, 40 | Brünner's gland, 140, 162, 164, 166 |
| tendon, 39 | Brush border, 183, 188 |
| vein, 39 | Buffy coat, 45, 50 |
| Volkmann's canal, 32, 38, 39 | Bulb, hoof, 128, 129 |
| word roots, 32 | Bulbar conjunctiva, 268, 269, 271, 272, 273, 274, 276 |
| zone of calcification, 36, 37, 38 | 279 |
| zone of hypertrophy, 35, 36, 37, 38 | Bulbourethral (Cowper's) gland, 226, 236 |
| zone of multiplication, 35, 37, 38 | Bursa of Fabricius, 91, 104 |
| zone of ossification, 35, 36, 37, 38 | Daton of 1 noticino, 7 1, 10 1 |
| zone of reverse cartilage, 35, 37 | C (parafollicular) cell, 212, 218 |
| Bone marrow, 35, 38, 53–56 | Calamus (quill), 107, 133, 134 |
| azurophilic granule, 55 | Calcified cartilage, 34, 36, 37, 38 |
| band cell, 54 | CAM (chorioallantoic membrane), 244, 254, 256, 257 |
| | 21111 (enotioniumore inclination), 2 11, 201, 200, 207 |

| Canal of Hering, 141 | internal elastic membrane, 77, 80, 81, 82 |
|--|---|
| Canal of Schlemm, 270, 279 | lymph node, 87 |
| Canaliculi, 32, 38 | lymphatic vessel, 78, 87 |
| Capillary | macrophage, 79 |
| air capillary, 197, 209 | mast cell, 79 |
| cardiovascular system, 77, 79 | medium artery, 77, 81, 82 |
| choriocapillary layer, 268, 275, 276, 281 | medium vein, 78, 82 |
| | mesenchyme-like tissue, 85 |
| digestive system, 155, 176, 182 ear, 290 | |
| | myocardium, 78, 84, 85, 86 |
| eye, 276, 281 | myofibrils, 86 |
| integument, 107, 118, 120, 134, 135 | neurons, 86 |
| lymphatic system, 90, 100 | pancreas, 81 |
| nervous system, 69 | parenchyma cell, 86 |
| postcapillary venule, 90, 97 | pericardial cavity, 86 |
| respiratory system, 197, 209 | pericardium, 86 |
| sheathed, 90 | plasma, 80, 87 |
| sinus capillary, 107, <i>134</i> , <i>135</i> | plasma cell, 79 |
| urinary system, 192 | portal vein, 82 |
| Capillary layer, 91, 104 | pulmonary artery, 83, 84, 85 |
| Capsule | pulmonic (semilunar) valve, 85 |
| bone, 40 | Purkinje cell, 86 |
| Bowman's, 184, 187, 188, 194 | renal artery, 82 |
| endocrine system, 220, 221, 222, 223 | right auricle, 84 |
| eye, 269, 279 | skeletal muscle, 79, 80 |
| lens, 272 | small artery, 77, 79, 80, 81 |
| lens capsule, 272 | small vein, 77, 79, 80 |
| lymphatic system, 90, 94, 95, 96, 98, 99, 100, 101 | smooth muscle, 79, 84 |
| male reproductive system, 235 | sweat gland, 80 |
| nervous system, 75 | tunica adventitia, 77, 80, 81, 82, 83, 85 |
| urinary system, 183, 186, 187 | tunica intima, 77, 81, 82, 83, 84, 85 |
| Capsule of Glisson, 141, 142, 174 | tunica media, 77, 80, 81, 82, 83, 84, 85 |
| Cardiac gland, 157, 158, 160 | Type I cell, 87 |
| Cardiac gland region, 140, 157, 158, 160 | Type II cell, 87 |
| Cardiac muscle, 57, 58, 64, 84 | umbilical artery, 83 |
| Cardiac skeleton, 78, 85, 86 | uterine gland, 79 |
| Cardiac sphincter, 158 | valves, 78, 80, 82, 85, 87 |
| Cardiovascular system, 77–87 | vasa vasorum, 78, 84 |
| adipose tissue, 81, 82, 84, 85, 86, 87 | vein, 81, 87 |
| anastomotic artery, 83 | vena cava, 84 |
| aorta, 83, 84, 85 | venule, 77, 79, 80, 87 |
| aortic body, 86, 87 | word roots, 78 |
| arteriole, 77, 79, 80, 86, 87 | Carpal gland, 121 |
| arteriovenous anastomoses, 78, 83 | Cartilage, 27–30 |
| artery, 81, 87 | amorphous ground substance, 27 |
| atrium, myocardium, 85 | chondroblast, 27 |
| | |
| capillary, 77, 79 | chondrocyte, 27, 29, 30 |
| cardiac muscle, 84 | collagenous fiber, 27, 28, 30 |
| cardiac skeleton, 78, 85, 86 | elastic cartilage, 27, 29, 30 |
| cecum, 87 | elastic fiber, 28, 29, 30 |
| chondrocyte, 86 | extracellular matrix, 27 |
| collagenous fiber, 84 | fibrocartilage, 27, 30 |
| elastic fiber, 81, 84 | hyaline cartilage, 27, 29 |
| endocardium, 78, 84 | integument, 130 |
| endothelial cell, 79, 80 | interterritorial matrix, 27, 29 |
| endothelium, 77, 87 | isogenous groups, 27, 29 |
| epicardium, 78 | lacuna, 27, 29, 30 |
| epithelioid cell, 83 | perichondrium, 27, 29 |
| erythrocyte, 79, 80, 81, 87 | plate, 95 |
| external elastic membrane, 81, 82 | respiratory system, 198 |
| glomus, 83 | territorial matrix, 27, 29 |
| heart, 78 | word roots, 28 |

| Cartilage plate, 95 | blood, 43, 51, 52 |
|--|---|
| Cartilaginous frame, 290 | bone, 38 |
| Cartilaginous tracheal ring, 206, 207 | bone marrow, 56 |
| Caruncle, 244, 245, 253 | cardiovascular system, 81, 87 |
| Cat | cartilage, 30 |
| blood, 44, 45 | connective tissue, 22 |
| bone, 37, 38, 39 | digestive system, 141–142, 177, 178, 179, 180, 181, |
| bone marrow, 55, 56 | 182 |
| cardiovascular system, 80, 81, 82, 86 | ear, 284–285, 290 |
| cartilage, 29 connective tissue, 22 | endocrine system, 213–214, 221, 222, 223 erythrocyte, average size, 9 |
| digestive system, 146, 148, 153, 159, 161, 162, 164, | eye, 269–270, 279, 280, 281, 282 |
| 165, 167, 168, 170, 171, 173 | female reproductive system, 246, 261, 262, 263, 264 |
| endocrine system, 215, 220 | 265 |
| epithelium, 14, 16, 17, 18 | integument, 132, 133, 134, 135, 136, 137, 138 |
| eye, 272, 274 | lymphatic system, 91, 103, 104 |
| female reproductive system, 247, 251, 259, 260 | male reproductive system, 227–228, 240, 241, 242 |
| integument, 108, 112, 114, 122, 123 | nervous system, 75, 76 |
| lymphatic system, 92, 93, 103 | respiratory system, 196–197, 205, 206, 207, 208, |
| male reproductive system, 237 | 209 |
| muscle, 59, 61, 63, 64 | urinary system, 184–185, 193, 194 |
| nervous system, 69, 74, 75 | Chief cell |
| respiratory system, 200, 202, 203, 204, 205 | digestive system, 140, 158, 159, 160, 161, 179 |
| urinary system, 187, 191 | endocrine system, 223 |
| Cationic (basic) stain, 5 | Chondrocyte, 27, 29, 30, 86 |
| Caudal larynx (syrinx), 197, 207, 208 | Chorioallantoic connective tissue, 255 |
| Caudal region, 213 | Chorioallantoic membrane (CAM), 244, 254, 256, 257 |
| Cavernous space | Chorioallantoic villus, 256, 257, 258 |
| female reproductive system, 259 | Choriocapillary layer, 268, 275, 276, 281 |
| male reproductive system, 227, 236, 237, 238, | Chorion laeve, 254, 256 |
| 239 Cavom ava voin 100 | Choroid, 268, 270, 271, 274, 275, 281, 282 |
| Cavernous vein, 198 | Charonaffa cell 221 |
| Cavity of air sac, 209 Cavity of lens, 279 | Chromaffin cell, 221 Chromophil, 212, 221 |
| Cavity of Rathke's pouch, 212, 215, 216 | Chromophobe, 212, 221 Chromophobe, 212, 216, 217, 221 |
| Cavity of renal pelvis, 189, 190 | Cilia |
| Cavity of vitreous humor, 269, 271, 279 | endocrine system, 219 |
| Cecal band, 169 | female reproductive system, 250, 253, 258 |
| Cecal tonsil, 103 | male reproductive system, 241, 242 |
| Cecum, 87, 140, 142, 168, 169, 180 | nervous system, 71 |
| Cellular tapetum lucidum, 268, 276 | stereocilia, 11, 233 |
| Cementum, 144 | Ciliary body, 268, 270, 272, 279, 280 |
| Centering slides, 5 | Ciliary muscle, 268, 270, 271, 272 |
| Central artery, 90, 99, 100 | Ciliary process, 268, 271, 272, 273, 279 |
| Central canal, 70, 71, 75 | Ciliated cell, 222, 264 |
| Central lacteal, 163, 167 | Ciliated epithelium, 263 |
| Central layer, 138 | Ciliated pseudostratified columnar epithelium, 16, 197 |
| Central pallor, 41 | Circular muscle, 251 |
| Central vein | Circumanal gland, 141, 170, 171, 172 |
| digestive system, 173, 174, 182 | Circumferential lamellae, 32 |
| urinary system, 184 | Circumvallate papilla, 139, 147 |
| Centroacinar cell, 176, 177 | Classic lobule, 141 |
| Cephalic region, 213 Cerebellum, 67, 69, 222 | Claw, 126, 137 Claw fold, 126 |
| Cerebral cortex, 67 | Clear cell, 121 |
| Cerebrospinal fluid, 68 | Clearing specimens, 3 |
| Ceruminous gland, 286 | Clitoris, 245 |
| Cervix, 244, 258 | Cloaca, 142, 181, 262 |
| Cheek, 143 | Cochlea, 289, 290 |
| Chestnut, 127 | Cochlear canal, 284 |
| Chicken | Cochlear duct, 284, 289 |

| Cochlear nerve, 289, 290 | lymphocyte, 20, 22, 23, 25 |
|---|---|
| Collagenous band, 124 | macrophage, 20, 24 |
| Collagenous fiber | male reproductive system, 237, 238 |
| cardiovascular system, 84 | mast cell, 20, 22, 23 |
| cartilage, 27, 28, 30 | mesenchyme tissue, 20, 22 |
| connective tissue, 20, 22, 23, 24 | mucous connective tissue, 20, 22 |
| integument, 117 | muscle, 19 |
| Collecting duct, 194, 282 | neutrophil, 20, 22 |
| Collecting tubule, 184, 185, 187, 188, 189, 190, 191, | plasma cell, 20, 22, 23 |
| 194 | reticular fiber, 20, 25 |
| Colloid, 218, 219, 221 | reticular tissue, 20, 25 |
| Colon, 140, 169 | Schiff's reagent, 20 |
| Columnar cell 250 260 | silver stains, 20 |
| Columnar cell, 259, 260 | tendon and tendon sheath, 24, 25 |
| Columnar epithelium bistratified, 124, 125 | urinary system, 187 |
| ciliated pseudostratified columnar epithelium, 16, 197 | Weigert's resorcin fuchsin, 20 word roots, 21 |
| digestive system, 157, 158, 160, 161, 167, 172, 181 | Connective tissue (band of), 155, 156 |
| eye, 277 | Connective tissue (coat of), 155, 156 Connective tissue (coat of), 151 |
| female reproductive system, 250, 251, 263 | Connective tissue (core of), 147, 148 |
| male reproductive system, 232, 237 | Connective tissue (lamina of), 209 |
| pseudostratified columnar epithelium, 12, 13, 16, | Connective tissue (papilla of), 143, 145, 146, 147, 154, |
| 196, 263 | 158 |
| respiratory system, 196, 197, 202, 208 | Connective tissue (partition of), 174 |
| simple columnar epithelium, 158, 160, 161, 172, 181 | Connective tissue (septum of), 176 |
| stratified columnar epithelium, 12, 13, 18, 208, 237, | Connective tissue sheath, 106, 116, 118, 119 |
| 277 | Connective-tissue (trabecula of), 238 |
| types, 13, 16 | Constrictor (sphincter) muscle, 268, 271, 273, 279, 280 |
| Comb, 107, 134, 135 | Contour feather, 107 |
| Common follicular opening, 106, 114 | Coprodeum, 181 |
| Compact bone, 32, 38, 39 | Cornea, 267, 269, 270, 271, 272, 273, 274, 276, 279, |
| Compound follicle, 106 | 280 |
| Concentric lamellae, 32 | Corneal stoma, 274 |
| Cones, 275, 276, 281 | Corneoscleral layer (fibrous tunic), 267, 269 |
| Conical papilla, 148, 156 | Corneoscleral trabecular meshwork, 268, 272, 273 |
| Conjunctiva, 269 | Corneous cell, 132 |
| Connecting duct (excretory canal), 228, 240, 242 | Corona, 89, 92 |
| Connective tissue, 19–25. See also specific tissues | Corona radiata, 244, 247 |
| adipocyte, 20, 24, 25 | Coronary dermis, 128 |
| adipose tissue, 20, 21, 25 | Coronary epidermis, 128 |
| amorphous ground substance, 19, 22 argyrophilic fiber, 20 | Cornary region of hoof, 128, 129, 130 |
| arteriole, 24 | Corpora amylacea, 122, 123 Corpora nigra (iris granule), 268 |
| collagenous fiber, 20, 22, 23, 24 | Corpus albicans, 244, 248 |
| connective tissue proper, 19, 20 | Corpus cavernosum, 227, 237, 238 |
| dense connective tissue, 20 | Corpus cavernosum clitoridis (erectile tissue), 245 |
| dense irregular connective tissue, 20, 24 | Corpus cavernosum urethra (corpus spongiosum), 227, |
| dense regular connective tissue, 20, 24 | 236, 237 |
| ear, 288 | Corpus luteum, 244, 247, 249 |
| elastic fiber, 20, 22, 25 | Corpus nigrum, 274 |
| elastic tissue, 20, 21, 25 | Corpus penis (body of penis), 227, 238, 239 |
| embryonal connective tissue, 19 | Corpus spongiosum (corpus cavernosum urethra), 227, |
| eosinophil, 20, 22, 23, 24 | 236, 237 |
| epithelium, 19, 22, 24 | Cortex |
| erythroctye in capillary, 22 | hair, 106, 116, 118 |
| extracellular matrix, 19, 20 | kidney, 183, 186, 188, 193, 194 |
| fibroblast, 19, 20, 22, 23, 24, 25 | lymph node, 90, 95, 96, 97, 101, 102, 104 |
| globular leukocyte, 20, 23 | Cortical cell, 223 |
| integument, 123 | Cortical labyrinth (pars convoluta), 183 |
| loose (areolar) connective tissue, 20, 22, 23, 24 | Cortical nephrons, 184 |
| lymphatic system, 103 | Cortical sinus, 94, 96 |

| Cortical stroma, 243 | Dental lamina, 143 |
|--|--|
| Cortical tissue, lymphatic system, 95 | Dental pad, 145 |
| Cortical tissue (interrenal), 213 | Dental papilla, 143, 144 |
| Cortical type, 184 | Dental pulp, 144 |
| Cotyledonary placenta, 244, 257 | Dental sac, 143, 144 |
| Covering membrane, 281 | Dentin, 144 |
| Coverslipping, 4 | Dentinoenamel junction, 144 |
| Cow | Dermal feather papilla, 107, 132, 135 |
| blood, 47 | Dermal lamina, 131 |
| cartilage, 29 | Dermal papilla, 106, 108, 110, 116, 127, 130, 132, |
| connective tissue, 22, 23, 24, 25 | 240 |
| digestive system, 145, 149, 155, 156, 161, 166, 169, | Dermis |
| 177 | coronary, 128 |
| endocrine system, 219, 220, 221 | eye, 282 |
| epithelium, 15, 16 | histology, 10 |
| eye, 278 | integument, 106, 108, 109, 110, 113, 114, 115, 116, |
| female reproductive system, 248, 249, 250, 253, 257, | 117, 118, 121, 125, 126, 127, 130, 132, 133, 134 |
| 258 | 135, 136, 137 |
| integument, 110, 111, 119, 122, 123, 124, 125, 127 | laminar, 128, 130, 131 |
| lymphatic system, 94, 95, 96, 97, 100 | male reproductive system, 237 |
| nervous system, 71, 75 | nervous system, 76 |
| respiratory system, 201, 202, 205 | perioplic, 128, 130 |
| urinary system, 190, 191, 192 | Descemet's membrane, 267, 273, 274, 276, 280 |
| Cowper's gland (bulbourethral), 226, 236 Crackling artifact, 10 | Diencephalon, 221 Diestrus, 244, 245, 252, 261 |
| Crampton's muscle, 279 | Diffuse lymphatic tissue |
| Crenated erythrocyte, 42 | diffuse placenta, 244, 256, 257 |
| Crista ampullaris, 284, 288 | eye, 277 |
| Crop, 141, 178 | lymphatic system, 92, 93, 94, 95, 96, 97, 98 |
| Crypt, 94, 256, 257, 258 | Digestive system, 139–182 |
| Crypt of Lieberkühn | A cell, 176, 177, 182 |
| digestive system, 140, 142, 162, 164, 165, 166, 167, | abomasum, 160, 162 |
| 168, 169, 170, 171, 172, 180, 181 | acinar cell, 176 |
| lymphatic system, 103 | acinus, 176, 177, 182 |
| Cryptal epithelium, 258 | adipose cell, 165 |
| Cuboidal epithelium | adipose tissue, 145, 148, 151, 153, 154, 160, 165, |
| bistratified, 124 | 169, 170 |
| defined, 11, 13, 15 | adventitia, 139, 153, 154, 175 |
| female reproductive system, 251, 253 | alveolar bone, 143, 144 |
| male reproductive system, 232 | ameloblast, 144 |
| respiratory system, 203 | anal canal, 140, <i>170</i> , <i>172</i> |
| Cumulus oophorus, 244, 247 | anal gland, 140, <i>170</i> , <i>171</i> |
| Cupula, 288 | anal sac, 141, <i>170</i> |
| Cuticle, 116 | anastomotic artery, 176 |
| Cyst, 219, 221 | apocrine tubular gland, 171 |
| Cystic duct, 176 | arteriole, 155, 162 |
| Cytoplasmic process, 276 | artifact, 164 |
| Cytotrophoblast, 255 | attachment epithelium, 144 |
| D 1 11 424 450 475 477 240 200 | Auerbach's plexus, 167 |
| Dark cell, 121, 159, 175, 176, 219, 290 | B cell, 176, 177, 182 |
| Dark principal cell, 212 | basal cell, 179 |
| Dark zone, 140 | basal striations, 149 |
| Dartos muscle, 117 | bile canaliculus, 141, 173 |
| Deciduate, 244 | bile duct, 173, 174 |
| Deep cortex, 94, 97 | binucleate hepatocyte, 174 |
| Deep glandular layer, 254 | blood vessel, 144, 151, 152 |
| Dehydration, 3 Dendrite, 65, 67, 71 | Brünner's gland, 140, 162, 164, 166 canal of Hering, 141 |
| Dense connective tissue, 20 | capillary, 155, 176, 182 |
| Dense irregular connective tissue, 20, 24 | capsule of Glisson, 141, 142, 174 |
| Dense regular connective tissue, 20, 24 | cardiac gland, 157, 158, 160 |
| | |

| Digastive exetem Continued | gallbladder, 141, 142, 175 |
|--|---|
| Digestive system, Continued | |
| cardiac gland region, 140, 157, 158, 160 | gastric furrow, 140, 162 |
| cardiac sphincter, 158 | gastric gland, 179 |
| cecal band, 169 | gastric pit (foveolae), 140, 158, 159, 160, 161, 162, 179 |
| cecum, 140, 142, 168, 169, 180 | gingiva, 144 |
| cementum, 144 | gland cell, 178, 179 |
| central lacteal, 163, 167 | glands of the anal sac, 170, 171 |
| central vein, 173, 174, 182 | glandular stomach, 140 |
| centroacinar cell, 176, 177 | globular leukocyte, 161, 165, 167 |
| cheek, 143 | goblet cell, 140, 141, 151, 163, 164, 167, 168, 169, |
| chicken, 141–142 | 170, 172, 174, 175, 176, 177 |
| chief cell, 140, 158, 159, 160, 161, 179 | hair follicle, 143, 172 |
| circumanal gland, 141, 170, 171, 172 | hard palate, 145 |
| circumvallate papilla, 139, 147 | hepatic artery, 173 |
| classic lobule, 141 | hepatocyte, 141, 173, 174, 182 |
| cloaca, 142, 181 | hepatoid gland, 141 |
| colon, 140, 169 | Herbst corpuscle, 181 |
| columnar epithelium, 157, 167 | horn tubule-like structure, 145 |
| conical papilla, 148, 156 | hyaline cartilage, 177 |
| connective tissue (band of), 155, 156 | ileum, 140, 141, 168, 180 |
| connective tissue (coat of), 153, 150 | intercalated duct, 141, 149, 150, 151, 176, 177 |
| connective tissue (core of), 137 connective tissue (core of), 147, 148 | intercellular space, 178 |
| connective tissue (papilla of), 143, 145, 146, 147, | interlobular connective tissue, 148, 149, 150, 151 |
| | |
| 154, 158 | interlobular duct, 141, 148, 151, 176 |
| connective tissue (partition of), 174 | intestinal absorptive cell, 163 |
| connective tissue (septum of), 176 | intestinal lumen, 167 |
| coprodeum, 181 | intestine, 140, 141–142 |
| crop, 141, 178 | intralobular duct, 143, 148 |
| crypt of Lieberkühn, 140, 142, 162, 164, 165, 166, | islet of Langerhans, 141, 176, 177 |
| 167, 168, 169, 170, 171, 172, 180, 181 | jejunum, 140, 141, 167, 168 |
| cystic duct, 176 | keratinized cell, 155 |
| dark cell, 159, 175, 176 | keratinized epidermis, 172 |
| dark zone, 140 | keratinized epithelium, 171 |
| dental lamina, 143 | keratinoid, 141, 179 |
| dental pad, 145 | Kupffer cell, 174 |
| dental papilla, 143, 144 | labial gland, 143 |
| dental pulp, 144 | lamina propria, 139, 144, 145, 152, 153, 154, 155, |
| dental sac, 143, 144 | 156, 157, 158, 159, 160, 161, 162, 163, 165, 167, |
| dentin, 144 | 168, 169, 170, 172, 175, 178, 179 |
| dentinoenamel junction, 144 | lamina subglandularis, 140, 159, 161 |
| duct, 143, 147, 151, 152, 153, 164, 177 | large intestine, 140, 142, 180 |
| duodenum, 140, 141, 162, 163, 164, 165, 166, 180 | light cell, 175, 176 |
| elastic fiber, 152, 157 | light zone, 140, 158 |
| enamel, 144 | lingual salivary gland, 147, 148 |
| enamel epithelium, 143, 144 | lip, 143 |
| enamel space, 144 | liver, 141, 142, 173, 174, 175, 182 |
| eosinophil, 167, 168 | lobule, 148, 173, 182 |
| epidermis, 170 | loose connective tissue, 145 |
| epithelial cell, 160 | lymphatic nodule, 158, 168, 169, 181 |
| epithelium, 139, 144, 145, 151, 175 | lymphatic tissue, 180 |
| erythrocyte, 182 | lymphatic vessel, 155, 157, 173 |
| esophageal gland, 158 | lymphocyte, 161, 163, 165 |
| | lymphoglandular ridge, 181 |
| esophagus, 140, 141, 152, 153, 154, 157, 158, 178 | |
| external anal sphincter, 170, 171 | mammals, 139–141 |
| fiber bundle, 144 | margo plicatus, 158 |
| filiform papilla, 139, 146, 148 | Meissner's plexus, 167 |
| foliate papilla, 139 | mesothelium, 174 |
| forestomach, 140 | mitotic figure, 168 |
| fundic gland, 158, 159, 160 | mixed gland, 140, 145, 151, 152, 153, 157 |
| fundic gland region, 140, 157, 158, 159, 160, 161 | mucosa, 139, 159, 160, 165, 166, 168, 169, 170, |
| fungiform papilla, 139, 146 | 171, 175 |

| mucous acinus, 143, 149, 150, 151, 175 | spine, 146 |
|---|--|
| mucous gland, 140, 145, 148, 152, 153, 154, 165, | spiral colon, 170 |
| 178 | stellate reticulum, 143, 144 |
| mucous neck cell, 159 | stomach, 140, 141, 158, 159, 161, 163 |
| mucous precursor, 160, 161 | stratified squamous epithelium, 143, 145, 147, 148, |
| muscle, 148 | 151, 152, 153, 154, 155, 156, 157, 158, 171, 172, |
| muscle band, 169 | 177, 178, 181 |
| muscularis, 175, 176 | stratum basale, 145 |
| muscularis externa, 139, 152, 153, 154, 155, 156, | stratum compactum, 140, 159, 162, 167 |
| 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, | stratum corneum, 145 |
| 169, 170, 171, 178, 179, 180 | stratum granulosum, 140, 143, 159, 162, 172 |
| muscularis mucosae, 139, 152, 153, 154, 156, 157, | stratum intermedium, 144 |
| 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, | stratum spinosum, 145, 146 |
| 170, 178, 180 | striated border, 140, 163, 165, 169, 170, 175 |
| odontoblast, 144 | striated duct, 149, 150, 151 |
| omasum, 156, 157 | sublingual gland, 150, 151 |
| oropharynx, 140, 151, 152 | submandibular gland, 149, 150 |
| pacinian corpuscle, 141 | submucosa, 139, 145, 152, 153, 154, 155, 156, 158, |
| palatine bone, 145 | 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, |
| pancreas, 141, 142, 176, 177, 182 | 170, 178, 179, 180 |
| paneth cell, 165 | sulcus, 178 |
| papilla, 139, 146, 157 | supporting cell, 146 |
| parietal cell, 140, 158, 159, 160, 161 | surface epithelium, 179 |
| parotid gland, 148, 149 | surface mucous cell, 159, 161, 163 |
| periodontal ligament, 144 | taenia coli, 140, 169, 170 |
| Peyer's patch, 140 | taste buds, 139, 141, 146, 147, 177, 178 |
| plica (folds), 178 | tertiary duct, 178, 179 |
| pore, 178 | tongue, 139, 177 |
| portal tract (area), 141, 174 | tubular gland, 140, 181 |
| portal vein, 173, 174 | tubular mucous unit, 150 |
| precentrum, 144 | urodeum, 181 |
| predentin, 144 | vein, 157, 169 |
| primary duct, 178 | ventriculus (gizzard), 141, 179 |
| proctodeum, 181 | villus, 140, 162, 164, 165, 167, 168, 180, 181 |
| proventriculus, 141, 178, 179 | word roots, 142 |
| pyloric gland, 161, 162 pyloric gland region, 140, 161, 162, 163 | Digital cushion, 111, 112 Digital pad, 106, 107, 111, 112, 126, 136 |
| rectoanal junction, 170, 171, 172 | Dilator, 271 |
| rectum, 140, 170 | Dilator muscle, 279, 280 |
| reticular fiber, 182 | Diplokaryocyte, 258 |
| reticulum, 155, 156 | Disseminate portion (pars disseminata), prostate, 226 |
| rumen, 155 | Distal convoluted tubule, 183, 187, 188, 193, 194 |
| salivary gland, 141, 177 | Distal interphalangeal joint, 40 |
| sebaceous gland, 143, 171, 172 | Distal phalanx, 126, 137 |
| secondary duct, 178, 179 | Dog |
| secretory unit, 149, 164, 165, 166 | blood, 44 |
| sensory cell, 146 | bone, 33, 35, 36, 38, 39 |
| serosa, 139, 154, 159, 161, 164, 166, 168, 169, 173, | cardiovascular system, 79, 82, 84, 85, 86 |
| 178, 180 | cartilage, 29, 30 |
| serous acinus, 143, 149, 150, 175 | connective tissue, 22, 25 |
| serous demilune, 143, 149, 150, 151 | digestive system, 143, 144, 145, 149, 150, 151, 152, |
| serous gland, 148, 165 | 153, 157, 158, 159, 161, 162, 163, 164, 167, 168, |
| simple columnar epithelium, 158, 160, 161, 172, 181 | 169, 170, 171, 172, 175, 176 |
| sinusoid, 141, 173, 174, 182 | ear, 286, 287, 288, 289 |
| skeletal muscle, 143, 146, 148, 151, 153, 154, 172, | endocrine system, 216, 218, 219, 220 |
| 181 | epithelium, 16, 17 |
| small artery, 169 | erythrocyte, average size, 9 |
| small intestine, 140 | eye, 271, 272, 273, 274, 275, 276, 277 |
| smooth muscle, 153, 154, 159, 181 | female reproductive system, 248, 249, 251, 252, 254, |
| soft palate, 145 | 255, 256, 258, 259, 260, 261 |
| space artifact, 144, 148 | histology, 10 |

| Dog, Continued | crista ampullaris, 284, 288 |
|--|--|
| integument, 108, 111, 113, 118, 119, 120, 123, 126 | cupula, 288 |
| lymphatic system, 93, 94, 95, 97, 98, 99, 101, 102 | dark cell, 290 |
| male reproductive system, 230, 233, 235, 236 | duct, 286 |
| muscle, 62 | elastic cartilage, 286 |
| nervous system, 67, 69, 71, 72 | endolymph, 284 |
| respiratory system, 198, 199, 201, 204 | endolymphatic space, 290 |
| urinary system, 186, 187, 189 | epidermis, 286 |
| Dorsal plate, 137 | external auditory meatus, 283, 286, 287 |
| Dorsal root ganglion, 71, 72, 76 | external ear, 283 |
| Down feather, 107 | facial nerve, 287 |
| Drying specimen on warmer, 4, 5 | gelatinous cupula, 284 |
| Duct | guttural pouch, 288 |
| alveolar duct, 203 | hair cell, 290 |
| bile duct, 173, 174 | hair follicle, 286 |
| cochlear duct, 284, 289 | helicotrema, 284 |
| collecting duct, 194, 282 | homogenous cell, 290 |
| connecting duct, 151, 262 connecting duct (excretory canal), 228, 240, 242 | incus, 284, 287 |
| cystic duct, 176 | inner tunnel, 289 |
| digestive system, 143, 147, 151, 152, 153, 164, 177 | |
| | internal ear, 283, 284 |
| ear, 286 | lagena, 285 |
| ejaculatory duct, 228 | lenticular process, 287 |
| of epididymis, 226, 228, 232, 242 | ligament, 287 |
| eye, 277 | light cell, 290 |
| female reproductive system, 259, 260, 265 | maculae, 284, 285 |
| intercalated duct, 141, 149, 150, 151, 176, 177, 199 | malleus, 284, 287 |
| interlobular duct, 122, 141, 148, 151, 176 | mammals, 283–284 |
| intralobular duct, 107, 111, 143, 148, 278 | membranous labyrinth, 284 |
| male reproductive system, 235, 236 | middle ear, 283, 284 |
| papillary duct, 183, 189, 190, 191 | middle-ear ossicles, 283 |
| perilobular collecting duct, 185 | mixed gland, 288 |
| primary duct, 178 | modiolus, 284, 289 |
| respiratory system, 199 | organ of Corti, 284, 289, 290 |
| secondary duct, 178, 179 | osseous spiral lamina, 284, 289 |
| semicircular duct, 284, 290 | otolith (otoconia, statoconia), 284, 288 |
| striated duct, 149, 150, 151, 199 | otolithic membrane, 284, 288 |
| of sweat gland, 110, 111, 112, 118, 120, 121 | perilymph, 284 |
| tertiary duct, 178, 179 | perilymphatic space, 290 |
| Ductus deferens (vas deferens), 226, 228, 233, 234, 242 | pigment granule, 286 |
| Duodenum, 140, 141, 162, 163, 164, 165, 166, 180 | pinna (auricle), 283 |
| Dura mater, 67 | pseudostratified epithelium, 288 |
| , | raphe, 290 |
| Ear, 283–290 | sacculus, 284, 287 |
| adipose tissue, 286 | scala media, 290 |
| ampulla, 284, 288 | scala tympani, 284, 289, 290 |
| annular ligament, 287, 288 | scala vestibuli, 284, 289, 290 |
| articular cartilage, 287 | sebaceous gland, 286 |
| basilar membrane, 284, 289, 290 | semicircular canal, 284, 290 |
| blood vessel, 290 | semicircular duct, 284, 290 |
| bone, 286 | sensory cell, 288 |
| bony labyrinth, 284 | sensory hair cell, 283, 284 |
| | |
| capillary, 290 | spiral ligament, 284, 289 |
| cartilaginous frame, 290 | spiral limbus, 284, 289 |
| ceruminous gland, 286 | spiral limbus, 284, 289 |
| chicken, 284–285 | spiral tunnel, 289 |
| cochlea, 289, 290 | stapes, 284, 287, 288 |
| cochlear canal, 284 | stria vascularis, 284 |
| cochlear duct, 284, 289 | supporting cell, 288, 290 |
| cochlear nerve, 289, 290 | synovial cavity, 287 |
| columella, 284 | tectorial membrane, 284, 289, 290 |
| connective tissue, 288 | tegmentum vasculosum, 285, 290 |

| temporal bone, 287, 288, 289 | diencephalon, 221 |
|--|--|
| tubular ceruminous gland, 283 | ependymal cell, 215 |
| tympanic cavity, 283, 284, 286, 287 | epitheliod cell, 220 |
| tympanic membrane, 283, 286, 287 | fiber of neuroglial cell, 218 |
| Type 1 cell, 288 | follicle, 212, 215, 216, 218, 219, 223 |
| utriculus, 284, 287 | ganglion, 220, 221, 223 |
| vestibular (Reissner's) membrane, 284, 289 | granulocyte, 222 |
| vestibule, 284, 287 word roots, 285 | Herring body, 212 hypophyseal cavity, 212 |
| Efferent arteriole, 189 | infundibular cavity, 212, 215, 216, 217, 221 |
| Efferent ductule, 226, 228, 231, 232, 240, 241, 242 | infundibular stalk, 212, 213, 215, 217, 221 |
| Ejaculatory duct, 228 | light cell, 219 |
| Elastic band, 202, 204 | light principle cell, 212 |
| Elastic cartilage, 27, 29, 30, 118, 199, 276, 286 | mammals, 211–213 |
| Elastic fiber | median eminence (ventral boundary of third |
| cardiovascular system, 81, 84 | ventricle), 212, 213, 215 |
| cartilage, 27, 29, 30 | medulla, 220, 221 |
| connective tissue, 20, 22, 25 | medullary cell, 223 |
| digestive system, 152, 157 | medullary (chromaffin) tissue, 213 |
| eye, 272, 280 | mucous cell, 222 |
| integument, 117, 124 | neurohypophysis, 211, 213 |
| lymphatic system, 100, 101 | parafollicular (C) cell, 212, 218 |
| male reproductive system, 238, 239 | parathyroid gland, 212, 213, 218, 219, 223 |
| respiratory system, 200, 201, 203, 208 | pars distalis, 211, 212, 213, 215, 216, 217, 221, |
| Elastic tendon, 134 Elastic ticsus, 20, 21, 25 | 222 |
| Elastic tissue, 20, 21, 25 Ellipsoid, 90, 99, 100 | pars intermedia, 211, 212, 215, 216 pars nervosa, 212, 213, 215, 216, 221 |
| Embryonal connective tissue, 19 | pars tuberalis, 211, 212, 213, 215, 216, 217, 221, |
| Enamel epithelium, 143, 144 | 222 |
| Enamel space, 144 | pineal gland, 212, 213, 218, 222, 223 |
| Endocardium, 78, 84 | pineal stalk, 218 |
| Endochondral (intracartilaginous) bone, 32, 34, 38 | pinealocyte, 212, 218 |
| Endocrine system, 211–223. See also Female | pituicyte (neuroglial cell), 212, 222 |
| reproductive system; Male reproductive system; | pituitary gland, 211–212, 213, 215, 216, 217, 221, |
| specific endocrine organs | 222 |
| acidophil (alpha cell), 212, 216, 217, 222 | principal cell, 219 |
| adenohypophysis, 211, 213 | principal (chief) cell, 212 |
| adipose tissue, 218, 223 | Rathke's pouch, 212, 215, 216 |
| adrenal cortex, 212 | rosette, 223 |
| adrenal gland, 212, 213, 220, 221, 223 | sinusoid, 217, 220, 221, 222, 223 |
| adrenal medulla, 213 | skull, 221, 222 |
| basophil (beta cell), 212, 217, 222 blood vessel, 216 | space artifact, 219 stroma, 219 |
| capsule, 220, 221, 222, 223 | thyroid gland, 212, 213, 218, 219 |
| caudal region, pars distalis, 213 | ultimobranchial body, 213 |
| cavity of Rathke's pouch, 212, 215, 216 | vestigal cavity, 212 |
| cephalic region, pars distalis, 213 | word roots, 214 |
| cerebellum, 222 | zona fasciculata, 213, 220, 221 |
| chicken, 213–214 | zona glomerulosa (zona multiformis), 213, 220, 221 |
| chief cell, 223 | zona intermedia, 213, 220, 221 |
| chromaffin cell, 221 | zona reticularis, 213, 220, 221 |
| chromophil, 212, 221 | Endolymph, 284 |
| chromophobe, 212, 216, 217, 221 | Endolymphatic space, 290 |
| cilia, 219 | Endometrial gland, 251, 252, 253 |
| ciliated cell, 222 | Endometrium, 257 |
| colloid, 218, 219, 221 | Endonysium, 57, 61, 62 |
| cortical cell, 223 cortical (interrenal) tissue, 213 | Endoneurium, 73 Endosteum, 31, 39 |
| cyst, 219, 221 | Endosteum, 31, 39 Endothelial cell, 79, 80, 97, 98, 99, 119, 134 |
| dark cell, 219 | Endotheliochorial placenta, 254, 255, 256 |
| dark principal cell, 212 | Endothelium, 77, 87, 237 |
| 1 1 · · · · · · · · · · · · · · · · · · | |

| Eosinophil blood, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 | stratified cuboidal epithelium, 12, 13 stratified epithelium, 11, 12 |
|--|---|
| bone marrow, 56 connective tissue, 20, 22, 23, 24 | stratified squamous epithelium, 12, 13, 17 transitional epithelium, 12, 13, 18 |
| digestive system, 167, 168 | word roots, 12 |
| Eosinophilic band cell, 55 | Erectile tissue (corpus cavernosum clitoridis), |
| Eosinophilic myelocyte, 55 | 245 |
| Ependymal cell, 215 | Eroded endometrial epithelium, 255 |
| Epicardium, 78 | Erythrocyte (red blood cell), 41 |
| Epidermal collar, 132 | bone marrow, 55, 56 cardiovascular system, 79, 80, 81, 87 |
| Epidermal lamina, 129, 131 Epidermal peg, 108, 135 | crenated erythrocyte, 42 |
| Epidermal spine, 237 | defined, 43, 44, 45, 46, 47, 48, 49, 50, 51 |
| Epidermis | digestive system, 182 |
| coronary, 128 | female reproductive system, 260, 265 |
| digestive system, 170, 172 | integument, 119, 134 |
| ear, 286 | lymphatic system, 98, 103 |
| eye, 277, 278, 282 | proerythrocyte, 53 |
| histology, 10 | urinary system, 191, 194 |
| integument, 105-106, 109, 110, 112, 114, 115, 117, | Erythroid, 53 |
| 118, 119, 121, 122, 123, 124, 125, 126, 130, 132, | Esophageal gland, 158 |
| 134, 135, 138 | Esophagus |
| keratinized, 172 | digestive system, 140, 141, 152, 153, 154, 157, 158 |
| laminar, 128 | 178 |
| male reproductive system, 239 | respiratory system, 200 |
| nervous system, 76 | Estrous cycle, 244 Fetrus, 244, 245, 252, 259, 260 |
| perioplic, 128, 130 respiratory system, 205 | Estrus, 244, 245, 252, 259, 260 Excretory canal (connecting duct), 228, 240, 242 |
| Epididymis, 226, 228, 233 | Extensor tendon, 40 |
| Epiglottis, 93, 199 | External anal sphincter, 170, 171 |
| Epimysium, 58 | External auditory meatus, 283, 286, 287 |
| Epineurium, 72 | External ear, 283 |
| Epiphyseal disc, 32, 37 | External elastic membrane, 81, 82 |
| Epithelial cell, 160, 220 | External tymponic membrane, 207, 208 |
| Epithelial tuft, 91, 104 | Extracellular matrix, 19, 20, 27 |
| Epitheliochorial placenta, 245, 257, 258 | Extrapulmonary primary bronchus, 197 |
| Epithelioid cell, 83 | Extratesticular rete testis, 226 |
| Epithelium, 11–18. See also Columnar epithelium; | Extrinsic muscle, 282 Eye, 267–282 |
| Cuboidal epithelium; Pseudostratified epithelium; Stratified squamous epithelium; Transitional | acinus, 278 |
| epithelium | annular pad, 269, 279 |
| basement (glassy) membrane, 11, 16 | anterior chamber, 269, 271, 272, 276, 279 |
| bistratified epithelium, 13, 17, 18 | anterior epithelium, 267, 274 |
| ciliated pseudostratified columnar epithelium, 16 | blind spot, 269 |
| columnar epithelium, 11, 13, 16 | blood vessel, 275, 281 |
| connective tissue, 11, 14, 19, 22, 24 | Bowman's membrane, 267, 280 |
| cuboidal epithelium, 11, 13, 15 | bridge, 270, 281 |
| digestive system, 139, 144, 145, 151, 175 | Bruch's membrane, 268 |
| female reproductive system, 264 | Brücke's muscle, 279 |
| goblet cell, 16 | bulbar conjunctiva, 268, 269, 271, 272, 273, 274, |
| hepatocyte, 14 integument, 108, 124, 125 | 276, 279 canal of Schlemm, 270, 279 |
| lamina propria, 16, 17 | capillary, 276, 281 |
| lymphocyte, 16 | capsule, 269, 279 |
| nervous system, 69 | cavity of lens, 279 |
| pseudostratified columnar epithelium, 12, 13, 16 | cavity of vitreous humor, 269, 271, 279 |
| respiratory system, 209 | cellular tapetum lucidum, 268, 276 |
| simple epithelium, 11, 12 | chicken, 269–270 |
| smooth muscle, 14, 16, 17, 18 | choriocapillary layer, 268, 275, 276, 281 |
| squamous epithelium, 11, 13, 14, 15 | choroid, 268, 270, 271, 274, 275, 281, 282 |
| stratified columnar epithelium, 12, 13, 18 | ciliary body, 268, 270, 272, 279, 280 |

ciliary muscle, 268, 270, 271, 272 nonpigmented epithelium, 273, 279, 280 ciliary process, 268, 271, 272, 273, 279 optic disc, 269, 271, 275 collecting duct, 282 optic nerve, 271, 275, 281 cones, 275, 276, 281 ora ciliaris retinae, 268, 271, 274 conjunctiva, 269 outer nuclear layer, 275, 276, 281 cornea, 267, 269, 270, 271, 272, 273, 274, 276. outer plexiform layer, 275, 276, 281 279, 280 palpebral conjunctiva, 269, 271, 272, 273, 277, 278 corneal stoma, 274 pars ciliaris retinae, 268, 271, 272, 274 corneoscleral trabecular meshwork, 268, 272, 273 pars iridica retinae, 268, 271 corpora nigra (iris granule), 268 pecten, 270, 281 pectinate ligament (uveal meshwork), 268, 270, 272, corpus nigrum, 274 273, 276, 279, 280 covering membrane, 281 Crampton's muscle, 279 pigmented columnar cell, 268 cytoplasmic process, 276 pigmented epithelium, 273, 275, 276, 279, 280, 281 pigmented myoepithelial cell, 268 dermis, 282 Descemet's membrane, 267, 273, 274, 276, 280 plasma cell, 282 diffuse lymphatic tissue, 277 posterior chamber, 269, 271, 272, 279 dilator, 271 posterior epithelium, 267, 274, 280 dilator muscle, 279, 280 pupil, 268, 271 duct, 277 retina, 267, 268, 270, 271, 274, 275, 276, 281, 282 elastic cartilage, 276 rods, 275, 276, 281 elastic fiber, 272, 280 sclera, 267, 269, 271, 272, 273, 274, 275, 276, 278, epidermis, 277, 278, 282 extrinsic muscle, 282 scleral cartilage, 269, 279, 280, 281, 282 eyelid, 269, 270, 277, 278, 282 scleral ossicles, 269, 279, 280 scleral trabecular meshwork, 270, 279 feather follicle, 282 fibroblast, 268 scleral venous plexus, 268, 271, 272 sebaceous gland (glands of Zeiss), 269, 278 fibrous tapetum lucidum, 268, 275 fibrous tunic (corneoscleral layer), 267, 269 secretory tubule, 282 filtration angle, 268, 270, 271, 272, 273, 279 serous acinus, 277 fornix of conjunctiva, 269, 278 skeletal muscle, 278 ganglion, 275, 276, 281 smooth muscle, 278 germinal zone, 272 space artifact, 274, 280 spaces of Fontana, 268, 270, 272, 273, 279 goblet cell, 277 hair follicle, 277, 278 sphincter (constrictor) muscle, 268, 271, 273, 279, harderian gland, 269, 270, 278, 282 Herbst corpuscle, 282 squamous epithelium, 280 hyaline cartilage, 277 stratified columnar epithelium, 277 inner limitating membrane, 275 stratified squamous epithelium, 277, 280 inner nuclear layer, 275, 276, 281 stroma (substantia propria), 267, 268, 273, 274, 280 inner plexiform layer, 275, 276, 281 superficial gland of nictitating membrane, 269, 277 interlobular connective tissue, 278, 282 suprachoroid layer, 268 intralobular duct, 278 sweat gland (glands of Moll), 269, 277, 278 iris, 268, 270, 271, 272, 273, 274, 279, 280 tapetal cell, 276 tapetum lucidum, 268, 271, 275, 276 Krause's gland, 278 lacrimal gland, 269, 270, 278, 282 tarsal (Meibomian) gland, 269, 277, 278 lamina cribrosa, 269, 275 tarsus, 269, 277, 278 lens, 267, 269, 271, 272 uveal trabecular meshwork, 268, 273 lens body, 269, 279 vascular layer, 268, 272, 275, 276 lens capsule, 272 vascular tunic (uvea), 267, 268, 270 lens epithelium, 269, 272 word roots, 270 zonular fiber, 268, 269, 271, 273, 279, 280 lens fiber, 269, 272, 279 limbus, 268, 271, 272, 273 Eyelid, 269, 270, 277, 278, 282 lipid vacuole, 280 loose connective tissue, 281 Facial nerve, 287 mammals, 267–269 Fascicles, 58 melanocyte, 268, 273, 274, 275, 276, 281 Fat cell (adipose cell), 20, 24, 25, 165 mucous acinus, 277 Fat vacuole, 187 myoepithelial cell, 271, 273 Feather, 132, 138, 282 nerve fiber, 275, 276, 281 Feather follicle, 132, 138, 282 nictitating membrane, 269, 270, 276 Feather muscle, 132, 133, 134

| Feather pulp, 107, 132, 133 | follicle, 243, 247, 261, 262 |
|---|---|
| Feather sheath, 132, 133 | germinal epithelium, 243, 248 |
| Female reproductive system, 243–265 | germinal vesicle, 246 |
| adipose tissue, 250 | glandular epithelium, 265 |
| adventitia, 259 | glandular groove, 263 |
| albumen, 264 | |
| | glans, 245 |
| allantoic blood vessel, 254, 255, 256, 257, 258 | granulocyte, 261 |
| allantoic epithelium, 254 | granulosa cell, 247, 248, 249 |
| ampulla, 244, 250 | hematoma, 254, 256 |
| anestrus, 244, 245, 259, 260 | hemorrhagic region, 253 |
| antrum, 244, 247, 248 | high columnar cell, 257 |
| apical cell, 265 | hilus cell, 243, 249 |
| artery, 252 | hypertrophied theca cell, 248 |
| artresia, 244 | indeciduate, 244 |
| atretic follicle, 246, 262 | infundibulum, 244, 246, 250, 262, 263 |
| basal cell, 265 | intercotyledonary endometrium, 257 |
| blood vessel, 253, 265 | intermediate cell, 245, 260, 261 |
| caruncle, 244, 245, 253 | interstitial cell, 246, 262 |
| | interstitial gland, 243, 248, 249 |
| cavernous space, 259 | |
| cervix, 244, 258 | isthmus of oviduct, 244, 246, 251, 262, 264 |
| chicken, 246 | keratinized cell, 259 |
| chorioallantoic connective tissue, 255 | labia, 245 |
| chorioallantoic membrane (CAM), 244, 254, 256, | labyrinthine surfaces, 245 |
| 257 | lamina propria, 250, 258, 259, 263 |
| chorioallantoic villus, 256, 257, 258 | longitudinal muscle, 251 |
| chorion laeve, 254, 256 | low columnar walls, 257 |
| cilia, 250, 253, 258 | magnum, oviduct, 246, 262, 263, 264 |
| ciliated cell, 264 | mammals, 243–245 |
| ciliated epithelium, 263 | maternal blood vessel, 255, 256, 257 |
| circular muscle, 251 | medulla, ovary, 243, 246, 261 |
| clitoris, 245 | membrana granulosa, 244, 246, 247, 248, 262 |
| cloaca, 262 | mesometrium, 252 |
| columnar cell, 259, 260 | metestrus, 244, 245, 253 |
| columnar epithelium, 250, 251 | microplacentome, 256 |
| | |
| corona radiata, 244, 247 | mucosa, 263 |
| corpus albicans, 244, 248 | multilaminar (late primary) follicle, 244 |
| corpus luteum, 244, 247, 249 | muscularis, 250, 259, 263, 264, 265 |
| cortical stroma, 243 | myometrium, 244, 251, 252, 253, 254, 257 |
| cotyledonary placenta, 244, 257 | neutrophil, 260, 261 |
| crypt, 256, 257, 258 | oocyte, 247, 261, 262 |
| cryptal epithelium, 258 | ovaries, 243, 246, 247, 248, 249, 261, 262 |
| cuboidal epithelium, 251, 253, 257 | oviduct, 244, 246, 249, 250, 263, 264, 265 |
| cumulus oophorus, 244, 247 | parabasal cell, 245, 260, 261 |
| cytotrophoblast, 255 | perimetrium (serosa), 244 |
| deciduate, 244 | perivitelline membrane, 246, 262 |
| deep glandular layer, 254 | placenta, 244, 254, 255, 256, 257, 258 |
| diestrus, 244, 245, 252, 261 | placental labyrinth, 254, 256 |
| diffuse placenta, 244, 256, 257 | placentome, 257 |
| diplokaryocyte, 258 | prepuce, 245 |
| | primary fold, 258, 263, 264, 265 |
| duct, 259, 260, 265 | |
| endometrial epithelium, 255 | primary follicle, 247 |
| endometrial gland, 251, 252, 253 | primary oocyte, 243 |
| endometrium, 244, 257 | primordial follicle, 243, 247 |
| endotheliochorial placenta, 254, 255, 256 | proestrus, 244, 245, 260 |
| epitheliochorial placenta, 245, 257, 258 | pseudostratified columnar epithelium, 263 |
| epithelium, 264 | pseudostratified epithelium, 253, 264, 265 |
| erectile tissue (corpus cavernosum clitoridis), 245 | regenerating gland, 263, 264 |
| erythrocyte, 260, 265 | resting gland, 263, 264 |
| estrous cycle, 244 | rete ovarii, 243, 249 |
| estrus, 244, 245, 252, 259, 260 | secondary fold, 258, 265 |
| folds, 245, 249, 250, 251 | secondary follicle, 244 |
| | |

| secretory bleb, 250 | Field of view, microscope, 8 |
|---|---|
| secretory cell, 264 | Filiform papilla, 139, 146, 148 |
| secretory gland, 263, 264 | Filoplume, 107 |
| serosa, 250, 263 | Filtration angle, 268, 270, 271, 272, 273, 279 |
| shell gland (uterus), 246, 262, 265 | Fixation, specimen, 3 |
| space artifact, 257 | Fluid-filled space, 75 |
| sperm-host gland, 246, 265 | Fold |
| spongy layer, 254, 256 | claw fold, 126 |
| stratified epithelium, 259 | female reproductive system, 245, 249, 250, 251 |
| stratified squamous epithelium, 258, 259 | follicular fold, 120 |
| stratined squamous epithendin, 250, 257 stratum vasculare, 244, 251, 252 | histology, 10 |
| stroma, 246, 247 | lymphatic system, 91 |
| submucosa, 250 | primary fold, 258, 263, 264, 265 |
| superficial cell, 245, 260, 261 | secondary fold, 258, 265, 265 |
| superficial intermediate cell, 245, 260, 261 | synovial fold, 40 |
| | |
| supraglandular layer, 254 | tertiary fold, 258, 263 |
| syndesmochorial placenta, 245 | vestibular fold, 93 |
| syntrophoblast, 255 | vocal fold, 200 |
| tertiary fold, 258, 263 | Foliate papilla, 139 |
| tertiary (Graafian) follicle, 244, 248 | Follicle |
| theca externa, 244, 246, 247, 248, 262 | compound, 106 |
| theca folliculi, 244, 247 | digestive system, 143, 172 |
| theca interna, 244, 246, 247, 248, 262 | ear, 286 |
| theca lutein cell, 248 | endocrine system, 212, 215, 216, 218, 223 |
| transitional epithelium, 259 | eye, 277, 278, 282 |
| trophoblastic projection, 255 | feather, 132, 138, 282 |
| tubular gland, 264 | female reproductive system, 243, 247, 261, 262 |
| tunica adventitia, 252 | hair, 106, 108, 109, 115, 118, 119, 120, 121, 122, |
| tunica albuginea, 243, 246, 247 | 124, 127, 130, 143, 172, 240, 277, 278, 286 |
| tunica intima, 252 | integument, 106, 108, 109, 115, 118, 119, 120, 121, |
| tunica media, 252 | 122, 124, 127, 130, 132, 134, 138 |
| urethra, 259 | lymphatic system, 90, 91, 104 |
| urethral epithelium, 259 | multilaminar (late primary), 244 |
| urodeum, 246 | primary, 106, 247 |
| uterine (endometrial) gland, 244, 256, 257 | primordial, 243, 247 |
| uterine horn, 251, 252, 253 | secondary, 106, 244 |
| uterus, 244, 253, 256, 265 | tertiary (Graafian), 244, 248 |
| vacuolar cell, 246, 255, 261 | tonsillar, 90 |
| vacuole, 265 | Follicular fold, 120 |
| vagina, 245, 246, 259, 262, 265 | Footpad, 137 |
| vaginal smear, 245, 260, 261 | Forestomach, 140 |
| vein, 252 | Formed elements, 41 |
| vestibular epithelium, 259 | Fornix of conjunctiva, 269, 278 |
| vestibular gland, 259 | Fossa, 93 |
| vestibule, 245, 260 | Fourth ventricle, 69 |
| vilous surfaces, 245 | Foveolae (gastric pit), 140, 158, 159, 160, 161, 162, |
| vulva, 245 | 179 |
| word roots, 246 | Frog, hoof, 128, 129 |
| yolk sphere, 262 | Fuchsin stains, 5 |
| zona pellucida, 244, 247, 248 | Fundic gland, 158, 159, 160 |
| zonary placenta, 244, 254, 255, 256, 257, 258 | Fundic gland region, 140, 157, 158, 159, 160, 161 |
| Fiber bundle, 144 | Fungiform papilla, 139, 146 |
| Fiber of neuroglial cell, 218 | Tungnorm papma, 199, 170 |
| Fibroblast, 19, 20, 22, 23, 24, 25, 135, 268 | Gallbladder, 141, 142, 175 |
| | |
| Fibrocartilaginous cord 239 | Ganglion |
| Fibroelestic membrane 201 | dorsal root ganglion, 71, 72, 76 |
| Fibrous astrogyte 60 | endocrine system, 220, 221, 223 |
| Fibrous astrocyte, 69 | eye, 275, 276, 281 |
| Fibrous capsule, 40 | nervous system, 66 |
| Fibrous tapetum lucidum, 268, 275 | parasympathetic ganglion, 72 |
| Fibrous tunic (corneoscleral layer), 267, 269 | spiral ganglion, 289 |

| 0 (140 163 | 1 1 1 100 100 114 115 117 110 110 |
|---|--|
| Gastric furrow, 140, 162 | sebaceous gland, 106, 109, 114, 115, 117, 118, 119, |
| Gastric gland, 179 | 120, 121, 122, 124, 143, 171, 172, 240, 269, 278, |
| Gastric pit (foveolae), 140, 158, 159, 160, 161, 162, | 286 |
| 179 | secondary, 263 |
| Gelatinous cupula, 284 | secretory, 264 |
| Germinal center, 89, 92 | serous, 148, 165, 198, 200, 202 |
| Germinal epithelium, 243, 248 | shell gland (uterus), 246, 262, 265 |
| Germinal matrix cell, 106 | sperm-host gland, 246, 265 |
| Germinal vesicle, 246 | sublingual gland, 150, 151 |
| Germinal zone, 272 | submandibular gland, 149, 150 |
| | |
| Giemsa's stain, 5 | superficial gland of nictitating membrane, 269, 277 |
| Gingiva, 144 | tarsal (Meibomian) gland, 269, 277, 278 |
| Gizzard (ventriculus), 141, 179 | thyroid gland, 212, 213, 218, 219 |
| Gland. See also Sweat gland; specific glands | tracheal gland, 201 |
| accessory, 226 | tubular, 140, <i>181</i> , <i>264</i> |
| adrenal gland, 212, 213, 220, 221, 223 | uterine (endometrial) gland, 79, 244, 256, 257 |
| anal gland, 140, 170, 171 | vestibular, 259 |
| apocrine tubular gland, 171 | Glands of Moll, 269, 277, 278 |
| Bowman's gland, 198, 206 | Glands of Zeiss, 269, 278 |
| Brünner's gland, 140, 162, 164, 166 | Glandular epithelium, 122, 265 |
| cardiac gland, 157, 158, 160 | Glandular groove, 263 |
| cardiac gland region, 140, 157, 158, 160 | |
| | Glandular stomach, 140 |
| carpal, 121 | Glans, 245 |
| circumanal gland, 141, 170, 171, 172 | Glans penis, 227, 237 |
| digestive system, 178 | Glassy (basement) membrane, 11, 16, 106, 121 |
| esophageal gland, 158 | Glisson's capsule, 141, 142, 174 |
| fundic gland, 158, 159, 160 | Globular leukocyte, 20, 23, 161, 165, 167 |
| fundic gland region, 140, 157, 158, 159, 160, 161 | Glomerulus, 184 |
| gastric gland, 179 | Glomus, 83 |
| gland cell, 178, 179 | Glottis, 199 |
| glands of Moll, 269, 277, 278 | Glycogen body, 75, 76 |
| glands of Zeiss, 269, 278 | Glycogen zone, 107, 137, 138 |
| harderian gland, 269, 270, 278, 282 | Goat |
| hepatoid gland, 141 | blood, 50 |
| integument, 120, 122, 138 | cardiovascular system, 79, 86 |
| | |
| interstitial, 243, 248, 249 | digestive system, 146, 147, 148, 156, 157, 160, 162, |
| Krause's gland, 278 | 166, 170, 174, 175 |
| labial gland, 143 | ear, 286 |
| lacrimal gland, 269, 270, 278, 282 | endocrine system, 219 |
| lingual salivary gland, 147, 148 | epithelium, 17, 18 |
| male reproductive system, 234, 235 | erythrocyte, average size, 9 |
| mammary, 107, 122, 123 | eye, 274, 278 |
| merocrine sweat gland, 106, 110, 111, 126 | integument, 117, 118 |
| mixed, 140, 145, 151, 152, 153, 157, 199, 200, 201, | male reproductive system, 234 |
| 202 | nervous system, 67 |
| mucous, 140, 145, 148, 152, 153, 154, 165, 178, | respiratory system, 199, 200, 201 |
| 205, 206 | urinary system, 190, 191, 192 |
| nasolabial, 110 | Goblet cell |
| | |
| parathyroid gland, 212, 213, 218, 219, 223 | digestive system, 140, 141, 151, 163, 164, 167, 168, |
| parotid gland, 148, 149 | 169, 170, 172, 174, 175, 176, 177 |
| pineal gland, 212, 213, 218, 222, 223 | epithelium, 16 |
| pituitary gland, 211–212, 213, 215, 216, 217, 221, | eye, 277 |
| 222 | respiratory system, 198, 200, 201, 202 |
| preputial, 240 | Gonocyte, 229 |
| prostate, 226, 235 | Graafian (tertiary) follicle, 244, 248 |
| pyloric gland, 161, 162 | Granular layer, 67, 69 |
| pyloric gland region, 140, 161, 162, 163 | Granulocyte |
| regenerating, 263, 264 | agranulocyte, 42 |
| resting, 263, 264 | blood, 42, 43, 51 |
| salivary gland, 93, 94, 141, 177 | bood, 42, 43, 51 bone marrow, 53 |
| sanvary grand, 10, 17, 171, 1// | |
| | endocrine system, 222 |

| female reproductive system, 261 | drying on warmer, 4, 5 |
|---|---|
| lymphatic system, 98, 103 | epidermis, 10 |
| respiratory system, 209 | erythrocyte, average size, 9 |
| Granulosa cell, 247, 248, 249 | field of view, 8 |
| Gray matter, 66, 70, 71, 75, 76 | fixation, 3 |
| Guttural pouch, 288 | fold, 10 |
| | fuchsin stains, 5 |
| H band, 62 | Giemsa's stains, 5 |
| Hair | goat erythrocyte, average size, 9 |
| bulb, 106, 114, 115, 117, 119, 121 | hematoxylin and eosin (H&E) stains, 5 |
| cell, ear, 290 | hints, 5, 8 |
| integument, 106, 113, 114, 115, 117, 119, 120 | histochemical procedure, 5 |
| sensory hairs, 198 | immunohistochemical procedure, 5 |
| sinus (tactile) hair, 106 | infiltration, 3 |
| Hair follicle | interpreting sections, 5, 6–7 |
| digestive system, 143, 172 | knife marks (scratches), 10 |
| ear, 286 | Köhler illumination, 8 |
| eye, 277, 278 | lake, 5 |
| integument, 106, 108, 109, 115, 118, 119, 120, 121, | Mallory stain, 5 |
| 122, 124, 127, 130 | Masson stain, 5 |
| male reproductive system, 240 | methylene blue stains, 5 |
| Hard palate, 145 | microscope use tips, 5–9 |
| Harderian gland, 269, 270, 278, 282 | miscroscopy, 8 |
| Hassall's (thymic) corpuscle, 90, 102 | mordant, 5 |
| Haversian canal, 32, 38, 39 | Orcein stains, 5 |
| Haversian systems, 32, 38 | paraffin procedure, 3, 4 |
| Head of epididymis, 232 | pointer in ocular, 5 |
| Heart, 78 | preparation, 3–5, 4 |
| Helicine artery, 227, 239 | relocating structures, 8 |
| Helicotrema, 284 | resorcin fuchsin, 5 |
| Hemal node, 90, 98 | Romanovsky stains, 5 |
| Hematopoietic tissue, 53 | sectioning with microtome, 3, 4, 5 |
| Hematoxylin and eosin (H&E) stains, 5 | separation artifact, 10 |
| Hemolymph node, 90 | silver stains for reticular fiber and nervous tissue, 5 |
| Hemorrhagic region, 253 | stain precipitate, 10 |
| Hen. See Chicken | staining, 4 |
| Henle's loop, 189, 190, 191, 194 | toluidine blue stains, 5 |
| Hepatic artery, 173 | transferring sections to slide, 4 |
| Hepatocyte, 14, 141, 173, 174, 182 | trichrome stains, 5 |
| Hepatoid gland, 141 | troubleshooting, 9 |
| Herbst corpuscle, 76, 135, 181, 282 | verniers, 8, 8 |
| Herring body, 212 | Weigert's stains, 5 |
| Heterophil, 43, 51, 52, 56 | Wright's stains, 5 |
| High columnar cell, 257 | Homogenous cell, 290 |
| Hilus cell, 243, 249 | Hoof (horse), 128, 132 |
| Histiocyte. See Macrophage | Horn, 127 |
| Histochemical procedure, 5 | Horn tubule, 127, 130, 131, 132 |
| Histology, 3–10 | Horn tubule-like structure, 145 |
| acidic (anionic) stains, 5 | Horse |
| artifact (imperfections), 9, 10 | blood, 46 |
| basic (cationic) stain, 5 | bone, 33, 40 |
| basophilic substances, 5 | cardiovascular system, 83, 87 |
| binocular microscope, 8 | cartilage, 30 |
| centering slides, 5 | connective tissue, 24 |
| chicken erythrocyte, average size, 9 | digestive system, 146, 148, 149, 153, 158, 160, 165, |
| clearing, 3 | 168, 169, 172, 173, 174, 176 |
| coverslipping, 4 | ear, 288 |
| crackling artifact, 10 | endocrine system, 216, 217, 219, 220 |
| dehydration, 3 | eye, 272, 273, 276, 277 |
| dermis, 10 | female reproductive system, 249, 250, 251, 252, 256, |
| dog erythrocyte, average size, 9 | 257, 258 |

| Horse, Continued | collagenous band, 124 |
|---|---|
| histology, 10 | collagenous fiber, 117 |
| integument, 109, 115, 116, 120, 124, 125, 127, 129, | comb, 107, 134, 135 |
| 130, 131, 132 | common follicular opening, 106, 114 |
| lymphatic system, 94, 95, 96, 97, 100 | compound follicle, 106 |
| male reproductive system, 230, 231, 232, 233, 237, | connective tissue, 123 |
| 238, 239, 240 | connective tissue sheath, 106, 116, 118, 119 |
| muscle, 61, 63 | contour feather, 107 |
| nervous system, 74 | corneous cell, 132 |
| respiratory system, 200, 204 | coronary dermis, 128 |
| urinary system, 188, 189, 192 | coronary epidermis, 128 |
| Humerus, 209 | coronary region of hoof, 128, 129, 130 |
| Hyaline cartilage, 27, 29, 34, 35, 177, 200, 201, 202, | corpora amylacea, 122, 123 |
| 277 | cortex, hair, 106, 116, 118 |
| Hypertrophied theca cell, 248 | cuticle, 116 |
| Hypophyseal cavity, 212 | dark cell, 121 |
| Trypophysear earry, 212 | dartos muscle, 117 |
| I band, 62 | dermal lamina, 131 |
| Ileum, 140, 141, 168, 180 | dermal (feather) papilla, 106, 107, 108, 110, 116, |
| Immature bone, 32, 33 | 127, 130, 132, 135 |
| Immunohistochemical procedure, 5 | dermis, 106, 108, 109, 110, 113, 114, 115, 116, |
| Imperfections. See Artifact | 117, 118, 121, 125, 126, 127, 130, 132, 133, 134, |
| Incus, 284, 287 | 135, 136, 137 |
| Indeciduate, 244 | digital cushion, 111, 112 |
| Infiltration, 3 | digital pad, 106, 107, 111, 112, 126, 136 |
| Infraorbital pouch, 121 | distal phalanx, 126, 137 |
| Infundibular cavity, 212, 215, 216, 217, 221 | dorsal plate, 137 |
| Infundibular stalk, 212, 213, 215, 217, 221 Infundibular stalk, 212, 213, 215, 217, 221, 222 | down feather, 107 |
| Infundibulum, oviduct, 244, 246, 250, 262, 263 | duct of sweat gland, 110, 111, 112, 118, 121 |
| Inguinal pouch, 122 | elastic cartilage, 118 |
| Inner core, 75 | elastic fiber, 117, 124 |
| Inner limitating membrane, 275 | elastic tendon, 134 |
| Inner nuclear layer, 275, 276, 281 | endothelial cell, 119, 134 |
| Inner plexiform layer, 275, 276, 281 | epidermal collar, 132 |
| | epidermal lamina, 129, 131 |
| Inner root sheath, 106, 116, 118, 119, 120 Inner tunnel, 289 | epidermal peg, 108, 135 |
| Integument, 105–138 | epidermis, 105–106, 109, 110, 112, 114, 115, 117, |
| adipose tissue, 117, 118, 135, 136, 138 | 118, 119, 121, 122, 123, 124, 125, 126, 130, 132, |
| | 134, 135, 138 |
| air space, 133 annular sinus, 106, 119 | epithelium, 108, 124, 125 |
| arrector pili muscle, 106, 114 | erythrocyte, 119, 134 |
| axial blood vessel, 132 | feather follicle, 132, 138 |
| bar, hoof, 128, 129 | feather muscle, 132, 133, 134 |
| barb, 1001, 128, 129 barb, 107, 132 | feather pulp, 107, 132, 133 |
| barbule, 107, 133 | feather sheath, 132, 133 |
| basal cell, 106, 131, 134, 138 | fibroblast, 135 |
| basement (glassy) membrane, 106, 121 | filoplume, 107 |
| bistratified columnar epithelium, 124, 125 | follicle, 134 |
| bistratified cuboidal epithelium, 124, 125 | follicular folds, 120 |
| | footpad, 137 |
| blood vessel, 118, 124, 125, 131 bone, 130, 131, 135, 137 | |
| | frog, hoof, 128, 129 |
| bulb, hoof, 128, 129 | germinal (matrix) cell, 106 |
| calamus (quill), 107, 133, 134 | gland, 120, 122, 138 |
| capillary, 118, 120 | glandular epithelium, 122 glycogen zone, 107, 137, 138 |
| carpal gland, 121 | |
| cartilage, 130 | hair, 106, 113, 114, 115, 117, 119, 120 |
| central layer, 138 | hair bulb, 106, 114, 115, 117, 119, 121 |
| chestnut, 127 | hair follicle, 106, 108, 109, 115, 118, 119, 120, 121, |
| claw, 126, 137 | 122, 124, 127, 130 Herbet corpuscle, 135 |
| claw fold, 126 | Herbst corpuscle, 135 |
| clear cell, 121 | hoof (horse), 128, 132 |

| horn, 127 | sebaceous gland, 106, 109, 114, 115, 117, 118, 119, |
|---|---|
| horn tubule, 127, 130, 131, 132 | 120, 121, 122, 124 |
| infraorbital pouch, 121 | sebaceous zone, 107, 137 |
| inguinal pouch, 122 | secondary follicle, 106 |
| inner root sheath, 106, 116, 118, 119, 120 | secondary laminae, 128 |
| intercellular bridges, 109, 111 | secretion, 123 |
| interlobular connective tissue, 122 | secretory acinus, 111 |
| interlobular duct, 122 | secretory cell, 121, 123 |
| intermediate cell, 138 | secretory portion, sweat gland, 120 |
| intermediate layer, 134 | secretory unit, 122, 123 |
| intertubular horn, 128 | sinus capillary, 107, 134, 135 |
| intralobular connective tissue, 123 | sinus (tactile) hair, 106 |
| intralobular duct, 107, 111 | sinus hair follicle, 109 |
| isthmus of uropygial gland, 107, 138 | sinus pad, 119 |
| keratin, 136 | skeletal muscle, 111, 114 |
| keratinocyte, 111 | skin, 105–106, 113, 114, 115, 116, 117, 118, 132, |
| lamina propria, 124, 125 | 133, 134 |
| laminae, 126 | smooth muscle, 118, 123, 124, 125 |
| laminae region of hoof, 128, 129, 131 | sole, hoof, 128, 129 |
| laminar dermis, 128, 130, 131 | spur, 136 |
| laminar horn, 128 | stem cell, 120 |
| limiting furrow, 126 | stratified squamous epithelium, 125 |
| lobe, lumen, 107, 137 | stratum basale, 106, 108, 109, 110, 112, 113, 114, |
| lobule, 122 | 117, 125, 127 |
| lymphatic vessel, 122, 124 | stratum corneum, 105, 108, 110, 112, 113, 114, |
| mammals, 105–107 | 117, 125, 127, 132, 133, 134, 135, 136 |
| mammary gland, 107, 122, 123 | stratum germinativum, 132, 133, 134, 135, 136, 137 |
| medulla, hair, 106, 116, 118 | stratum granulosum, 105, 108, 112, 113, 117, 127 |
| melanocyte, 106, 109, 132 | stratum internum, 128 |
| Merkel's cell, 119, 135 | stratum lucidum, 105, 108, 112 |
| merocrine sweat gland, 106, 110, 111, 126 | stratum medium, hoof, 128, 129, 131 |
| middle phalanx, 126 | stratum spinosum, 105, 108, 109, 110, 112, 113, |
| mucous connective tissue, 107, 134, 135 | 114, 117, 125, 127 |
| multilocular fat cell, 137 | stratum tectorium, 128 |
| myoepithelial cell, 106, 110, 120, 121, 123 | subcutis, 105, 108, 114, 136 |
| nasolabial gland, 110 | sulcus, 136 |
| nose, 109 | surface groove, 108, 110 |
| nuclei stack, 132, 135 | sweat gland, 106, 108, 109, 115, 116, 117, 120, |
| oral cavity, 135 | 121, 122, 123, 124 |
| outer-root sheath, 116, 118, 119, 120 | teat, 123 |
| | teat canal, 125 |
| pacinian corpuscle, 111 palatine ridge, 135 | teat sinus, 123, 124, 125 |
| papilla, 107 | tead sinus, 123, 124, 123 tendon, 137 |
| papillary layer, 106, 111, 112 | tomial edge, 135 |
| perioplic dermis, 128, 130 | trabecula, 119 |
| perioplic epidermis, 128, 130 | transitional layer, 134 |
| perioplic region of hoof, 128, 129 | tubular horn, 128 |
| | tubular norm, 128 tubuloacinar gland, 107 |
| periosteum, 135 | |
| pinna, 118 | unguinal scale, 137 |
| planum nasale, 108 | uropygial (preen) gland, 107, 137, 138 |
| planum nasolabiale, 110, 111 | vacuolated cell, 120 |
| planum rostrale, 110 | ventral plate, 137 |
| primary follicle, 106 | venule, 118 |
| primary hair, 118 | wattle, 107 |
| primary lamina, 128 | white line, hoof, 128, 129 |
| pyknotic cell, 116, 120 | word roots, 128 |
| rachis, 107 | Intercalated disc, 58, 63, 64 |
| reticular layer, 106, 111, 112 | Intercalated duct |
| reticulate scale, 136 | digestive system, 141, 149, 150, 151, 176, 177 |
| scrotum, 117, 118 | respiratory system, 199 |
| scutes, 136 | Intercellular bridges, 109, 111 |

| Intercellular space, 178 | Lagena, 285 |
|--|--|
| Intercostal muscle, 205 | Lake, 5 |
| Intercotyledonary endometrium, 257 | Lamellae, 32 |
| Interlobular connective tissue | Lamina cribrosa, 269, 275 |
| digestive system, 148, 149, 150, 151 | Lamina propria |
| eye, 278, 282 | digestive system, 139, 144, 145, 152, 153, 154, 155 |
| integument, 122 | 156, 157, 158, 159, 160, 161, 162, 163, 165, 167 |
| Interlobular duct, 122, 141, 148, 151, 176 | 168, 169, 170, 172, 175, 178, 179 |
| Interlobular septum, 234, 235 | epithelium, 16, 17 |
| Intermediate cell, 138, 245, 260, 261 | female reproductive system, 250, 258, 259, 263 |
| Intermediate layer, 134 | lymphatic system, 104, 124, 125 |
| Intermediate syringeal cartilage, 207, 208 | male reproductive system, 233, 234, 236, 237, 239 |
| Internal ear, 283, 284 | respiratory system, 196, 199, 200, 201, 202, 205 |
| Internal elastic membrane, 77, 80, 81, 82 | urinary system, 191, 192 |
| Internal tymponic membrane, 207, 208 | Lamina subglandularis, 140, 159, 161 |
| Interpreting sections, specimen, 5, 6–7 | Laminae region of hoof, 128, 129, 131 |
| Interrenal (cortical) tissue, 213 | Laminar dermis, 128, 130, 131 |
| Interstitial cell | Laminar epidermis, 128 |
| female reproductive system, 246, 262 | Laminar horn, 128 |
| Leydig cell, 226, 228, 229, 230, 231, 240 | Large intestine, 140, 142, 180 |
| Interstitial gland, 243, 248, 249 | Larynx, 195, 196, 197 |
| Interstitial systems, bone, 38 | Lens, eye, 267, 269, 271, 272 |
| Interterritorial matrix, 27, 29 | Lens body, 269, 279 |
| Intertubular horn, 128 | Lens capsule, 272 |
| Intestinal absorptive cell, 163 | Lens epithelium, 269, 272 |
| Intestinal lumen, 167 | Lens fiber, 269, 272, 279 |
| Intestine, 140, 141–142 | Lenticular process, 287 |
| Intracartilaginous (endochondral) bone, 32, 34, 38 | Leukocyte (white blood cell) |
| Intralobular connective tissue, 123 | blood, 41, 42, 43, 44 |
| Intralobular duct, 107, 111, 143, 148, 278 | connective tissue, 20, 23 |
| Intralobular vein, 193 | digestive system, 161, 165, 167 |
| Intramembranous bone, 32, 33 | globular (globule) leukocyte, 20, 23, 161, 165, |
| Intrapulmonary primary bronchi (mesobronchi), 197 | 167 |
| Iris, 268, 270, 271, 272, 273, 274, 279, 280 | infiltration, lymphatic system, 94 |
| Iris granule (corpora nigra), 268 | polymorphonuclear leukocyte, 43 |
| Islet of Langerhans, 141, 176, 177 | Leydig (interstitial) cell, 226, 228, 229, 230, 231, |
| Isogenous groups, 27, 29 | 240 |
| Isthmus of oviduct, 244, 246, 251, 262, 264 | Ligament, 287 |
| Isthmus of uropygial gland, 107, 138 | Light cell, 175, 176, 219, 290 |
| | Light principle cell, 212 |
| Jejunum, 140, 141, 167, 168 | Light zone, 140, 158 |
| Joint cavity, 40 | Limbus, 268, 271, 272, 273 |
| Juxtaglomerular cell, 184, 188 | Limiting furrow, 126 |
| | Lingual salivary gland, 147, 148 |
| Keratin, 136 | Lingual tonsil, 90 |
| Keratinized cell, 155, 259 | Lip, 143 |
| Keratinized epidermis, 172 | Lipid vacuole, 280 |
| Keratinized epithelium, 171 | Liver, 141, 142, 173, 174, 175, 182 |
| Keratinocyte, 111 | Lobe, lumen, 107, 137 |
| Keratinoid, 141, 179 | Lobule, 122, 148, 173, 182 |
| Kidney, 187 | Longitudinal muscle, 251 |
| Knife marks (scratches), 10 | Loop of Henle, 184 |
| Köhler illumination, 8 | Loose (areolar) connective tissue |
| Krause's gland, 278 | defined, 20, 22, 23, 24 |
| Kupffer cell, 174 | digestive system, 145 |
| T 1: 045 | eye, 281 |
| Labia, 245 | male reproductive system, 232, 233 |
| Labial gland, 143 | Low columnar cell, 257 |
| Labyrinthine surfaces, 245 | Lumen, 230 |
| Lacrimal gland, 269, 270, 278, 282 | Lungs, 196, 209 |
| Lacuna, 27, 29, 30, 32, 33, 38 | Lymph node, 87, 90, 95 |

| Lymphatic nodule | mucous acinus, 93 |
|---|---|
| digestive system, 158, 168, 169, 181 | multinucleate giant cell, 97 |
| lymphatic system, 89, 92, 93, 94, 95, 96, 98, 99, 103 | muscularis, 104 |
| male reproductive system, 240 | muscularis externa, 92, 93, 103 |
| respiratory system, 202 | myoid cell, 91, 104 |
| Lymphatic system, 89–104. See also Lymph node; | nonsinusal spleens, 90 |
| Lymphatic nodule; Lymphatic vessel | palatine tonsil, 90, 93, 94 |
| adipose tissue, 98, 103 | paraepiglottic tonsil, 90, 93 |
| arteriole, 90 | penicillus, 90 |
| blood vessel, 103 | periarterial lymphatic sheath (PALS), 90, 100 |
| bursa of Fabricius, 91, 104 | Peyer's patch, 89, 92 |
| capillary, 90, 100 | postcapillary venule, 90, 97 |
| capillary layer, 91, 104 | primary bronchi, 95 |
| capsule, 90, 94, 95, 96, 98, 99, 100, 101 | pseudostratified epithelium, 104 |
| cartilage plate, 95 | pulp artery, 90 |
| cecal tonsil, 103 | red pulp, 90, 91, 99, 100, 101, 103 |
| central artery, 90, 99, 100 | reticular cell, 92, 97, 98, 103 |
| chicken, 91 | reticular fiber, 96 |
| connective tissue, 103 | |
| | reticular structure, 91, 104 |
| corona, 89, 92 | salivary gland, 93, 94 |
| cortex, lymph node, 90, 95, 96, 97, 101, 102, 104 | septum, 90, 101, 104 |
| cortical sinus, 94, 96 | serosa, 100, 101 |
| cortical tissue, 95 | serous acinus, 93 |
| crypt, 94 | serous demilune, 93 |
| crypt of Lieberkühn, 103 | sheathed artery, 91, 103 |
| deep cortex, 94, 97 | sheathed capillary, 90 |
| diffuse lymphatic tissue, 92, 93, 94, 95, 96, 97, 98 | sinusal spleens, 90 |
| elastic fiber, 100, 101 | skeletal muscle, 93 |
| ellipsoid, 90, 99, 100 | smooth muscle, 96, 97, 100, 101 |
| endothelial cell, 97, 98, 99 | spleen, 90, 91, 99, 100, 101, 103 |
| epiglottis, 93 | splenic artery, 90 |
| epithelial tuft, 91, 104 | stratified squamous epithelium, 93, 94 |
| erythrocyte, 98, 103 | subcapsular sinus, 90, 94, 96, 98 |
| folds (plica), 91 | submucosa, 92, 93 |
| follicle, 91, 104 | thymus, 90, 91, 101, 102, 103, 104 |
| fossa, 93 | tonsil, 90 |
| germinal center, 89, 92 | tonsillar follicle, 90 |
| granulocyte, 98, 103 | trabecula, 90, 94, 96, 97, 99, 100, 101 |
| Hassall's (thymic) corpuscle, 90, 102 | trabecular artery, 90 |
| hemal node, 90, 98 | tracheobronchial lymph node, 95 |
| hemolymph node, 90 | tubal tonsil, 90 |
| lamina propria, 104 | undifferentiated epithelial cell, 91, 104 |
| leukocyte infiltration, 94 | venous sinus, 90, 99 |
| lingual tonsil, 90 | vesicles, 104 |
| lobule of thymus, 90 | vestibular fold, 93 |
| lymph node, 90, 95 | villus, 92, 93 |
| lymphatic nodule, 89, 92, 93, 94, 95, 96, 98, 99, | white pulp, 90, 91, 99, 101 |
| 103 | Lymphatic tissue, 180, 194, 208 |
| lymphatic vessel, 101 | Lymphatic vessel |
| lymphocyte, 92, 97, 98 | cardiovascular system, 78, 87 |
| macrophage, 97, 98 | digestive system, 155, 157, 173 |
| mammals, 89–91 | integument, 122, 124 |
| marginal zone, 90, 99, 100 | lymphatic system, 101 |
| mast cell, 97 | Lymphocyte |
| medulla, lymph node, 95, 97, 98 | blood, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51 |
| medulla, thymus, 90, 101, 102, 104 | connective tissue, 20, 22, 23, 25 |
| medullary cord, 94, 95, 97, 98 | digestive system, 161, 163, 165 |
| medullary sinus, 90, 94, 95, 97, 98 | epithelium, 16 |
| megakaryocyte, 98 | lymphatic system, 92, 97, 98 |
| mesothelium, 99, 103 | male reproductive system, 232 |
| mucosa, 92 | Lymphoglandular ridge, 181 |
| | |

| Macrophage (histiocyte) | muscularis, 233, 234 |
|---|--|
| cardiovascular system, 79 | myoid cell, 226, 230, 231 |
| connective tissue, 20, 24 | os penis, 227, 236, 237 |
| lymphatic system, 97, 98 | parietal prepuce, 240 |
| respiratory system, 203, 204, 209 | pelvis urethra, 227 |
| Macula densa, 184, 187, 188, 189 | penile urethra, 227, 237, 238 |
| Maculae, 284, 285 | penis, 227, 236, 237 |
| Magnum, oviduct, 246, 262, 263, 264 | peritoneum, 225, 227 |
| Male reproductive system, 225-242 | prepuce, 227 |
| accessory glands, 226 | preputial gland, 240 |
| adipose tissue, 238 | primary spermatocyte, 225, 230, 231, 240, 241 |
| adventitia, 234 | prostate gland, 226, 235 |
| ampulla, 226, 234 | pseudostratified epithelium, 232, 233, 234, 235, 242 |
| basal cell, 233, 234, 235 | rete testis, 226, 228, 229, 231, 232, 240, 241 |
| body of penis (corpus penis), 227, 238, 239 | retractor penis muscle, 238 |
| body of prostate gland, 226, 235 | sebaceous gland, 240 |
| bulbourethral (Cowper's) gland, 226, 236 | secondary spermatocyte, 226 |
| capsule, 235 | secretion, 234, 235 |
| cavernous space, 227, 236, 237, 238, 239 | secretory cell, 236 |
| chicken, 227–228 | seminal vesicle, 226, 234, 235 |
| cilia, 241, 242 | seminiferous tubule, 225, 227, 230, 231, 240, 241 |
| columnar epithelium, 232 | serosa, 233, 234 |
| connecting duct (excretory canal), 228, 240, 242 | serous cell, 236 |
| connective tissue, 237, 238 | Sertoli cell, 225, 226, 230, 231, 240, 241 |
| connective-tissue trabecula, 238 | sex cord, 229 |
| corpus cavernosum, 227, 237, 238 | smegma, 239 |
| corpus spongiosum (corpus cavernosum urethra), | smooth muscle, 232, 233, 236, 237, 238, 239, 242 |
| 227, 236, 237 | spermatids, 225, 230, 231, 240, 241 |
| cuboidal epithelium, 232 | spermatogenic cell, 225 |
| dermal papilla, 240 | spermatogonium, 225, 230, 231, 240, 241 |
| dermis, 237 | spermatozoa, 225, 232, 233, 234, 241, 242 |
| disseminate portion (pars disseminata), prostate, 226 | spongy bone, 236 |
| duct, 235, 236 | stereocilia, 233 |
| duct of the epididymis, 226, 228, 232, 242 | straight tubule, 226, 228, 229, 231, 241 |
| efferent ductule, 226, 228, 231, 232, 240, 241, 242 | stratified columnar epithelium, 237 |
| ejaculatory duct, 228 | stratified squamous epithelium, 239, 240 |
| elastic fiber, 238, 239 | stratum cavernosum (vascular stratum), 227, 235 |
| endothelium, 237 | supporting cell, 229 |
| epidermal spine, 237 | testis, 225, 227, 229, 230, 240, 241 |
| epidermis, 239 | trabecula, 235, 239 |
| epididymis, 226, 228, 233 | transitional epithelium, 235, 236, 239 |
| extratesticular rete testis, 226 | transitional zone, 226, 231 |
| fibrocartilaginous cord, 239 | tunica albuginea, 225, 227, 230, 231, 232, 237, 238, |
| gland, 234, 235 | 239, 241 |
| glans penis, 227, 237 | tunica vaginalis, 225, 230, 232 |
| gonocyte, 229 | urethra, 227, 235, 236, 237, 238, 239 |
| hair follicle, 240 | urethral pouch, 239 |
| head of epididymis, 232 | urethral process, 227, 239 |
| helicine artery, 227, 239 | vas deferens, 226, 228, 233, 234, 242 |
| interlobular septum, 234, 235 | villus-like projection, 233 |
| interstitial (Leydig) cell, 226, 228, 229, 230, 231, | visceral prepuce, 240 |
| 240 | word roots, 228 |
| lamina propria, 233, 234, 236, 237, 239 | Malleus, 284, 287 |
| loose connective tissue, 232, 233 | Mallory stain, 5 |
| lumen, 230 | Mammals |
| lymphatic nodule, 240 | blood, 41–43 |
| lymphocyte, 232 | bone marrow, 53 |
| male accessory gland, 226 | digestive system, 139-141 |
| mammals, 225–227 | ear, 283–284 |
| mediastinum testis, 225, 226, 231 | endocrine system, 211–213 |
| mucous cell, 236 | eye, 267–269 |

| female reproductive system, 243–245 | Middle phalanx, 126 |
|--|--|
| integument, 105–107 | Miscroscopy, 8 |
| lymphatic system, 89–91 | Mitotic figure, 55, 168 |
| male reproductive system, 225–227 | Mixed gland |
| urinary system, 183–184 | digestive system, 140, 145, 151, 152, 153, 157 |
| Mammary gland, 107, 122, 123 | ear, 288 |
| Mare. See Horse | respiratory system, 199, 200, 201, 202 |
| Marginal hematoma, 254, 256 | Modiolus, 284, 289 |
| Marginal zone, 90, 99, 100 | Molecular layer, 67, 68, 69 |
| Margo plicatus, 158 | Monocyte, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 |
| Marrow cavity, 31, 34, 35, 36, 37, 39 | Mordant, 5 |
| Masson stain, 5 | Mucosa. See also Muscularis mucosae; Submucosa |
| Mast cell, 20, 22, 23, 79, 97 | digestive system, 139, 159, 160, 165, 166, 168, 169, |
| Maternal blood vessel, 255, 256, 257 | 170, 171, 175 |
| Mature bone, 32, 33 | female reproductive system, 263 |
| Medial bronchial wall, 207 | lymphatic system, 92 |
| Median eminence (ventral boundary of third ventricle), | Mucous acinus |
| 212, 213, 215 | digestive system, 143, 149, 150, 151, 175 |
| Mediastinum testis, 225, 226, 231 | eye, 277 |
| Medium artery, 77, 81, 82 | lymphatic system, 93 |
| Medium vein, 78, 82 | Mucous cell, 209, 222, 236 |
| Medulla | Mucous connective tissue, 20, 22, 107, 134, 135, |
| endocrine system, 220, 221 | 191 |
| hair, 106, 116, 118 | Mucous gland |
| lymph node, 95, 97, 98 | digestive system, 140, 145, 148, 152, 153, 154, 165, |
| lymphatic system, 90, 101, 102, 104 | 178 |
| ovary, 243, 246, 261 | respiratory system, 205, 206 |
| urinary system, 184, 186, 189 | urinary system, 192 |
| Medullary cavity. See Marrow cavity | Mucous neck cell, 159 |
| Medullary cell, 223 | Mucous precursor, 160, 161 |
| Medullary (chromaffin) tissue, 213 | Multilaminar (late primary) follicle, 244 |
| Medullary cone, 185, 193 | Multilocular fat cell, 137 |
| Medullary cord, 94, 95, 97, 98 | Multinucleate giant cell, 97 |
| Medullary sinus, 90, 94, 95, 97, 98 | Multipolar neuron, 65, 70, 71, 75, 76 |
| Medullary tract, 185 | Muscle, 57–64. See also Skeletal muscle; Smooth |
| Megakaryocyte, 36, 39, 54, 55, 98 | muscle |
| Meissner's plexus, 72, 167 | A band, 62 |
| Melanocyte | arteriole, 60 |
| eye, 268, 273, 274, 275, 276, 281 | band, 169 |
| integument, 106, 109, 132 | bifurcation, 63 |
| Membrana granulosa, 244, 246, 247, 248, 262 | cardiac muscle, 57, 58, 64 |
| Membranous labyrinth, 284 | connective tissue, 19 |
| Meninges, 65, 68 | digestive system, 148 |
| Merkel's cell, 119, 135 | endomysium, 57, 61, 62 |
| Merocrine sweat gland, 106, 110, 111, 126 | epimysium, 58 |
| Mesangial cell, 184, 194 | fascicles, 58 |
| Mesenchyme tissue, 20, 22 | H band, 62 |
| Mesenchyme-like tissue, 85 | I band, 62 |
| Mesenteric blood vessel, 75 | intercalated disc, 58, 63, 64 |
| Mesobronchi (intrapulmonary primary bronchi), 197 | muscle fiber, 57 |
| Mesometrium, 252 | myofibrils, 61, 62, 63, 64 |
| | |
| Mesothelium, 67, 99, 103, 174, 204 | myofilaments, 57 |
| Metamyelocyte, 54 | perimysium, 58, 61 |
| Metarubricyte (orthochromatophilic erythroblast), 54, | sarcolemma, 58 |
| 55, 56 Material 244, 245, 253 | sarcoplasm, 57 |
| Metestrus, 244, 245, 253 | skeletal muscle, 57, 58, 61, 63 |
| Methylene blue stains, 5 | smooth muscle, 57, 59 |
| Microplacentome, 256 | venule, 60 |
| Microscope use tips, 5–9 | word roots, 58 |
| Microvilli, 11 | Z band, 62 |
| Middle ear, 283, 284 | Muscle fiber, 57 |

| mucosae digestive system, 175, 176 female reproductive system, 250, 259, 263, 264, 265 lymphatic system, 201, 202, 203 urinary system, 201, 202, 203 urinary system, 219, 192, 194 Muscularis externa digestive system, 139, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180 lymphatic system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myeloblast, 54 Myeloblast, 54 Myocardium, 78, 84, 85, 86 Myopithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, | |
|--|----|
| female reproductive system, 250, 259, 263, 264, 265 lymphatic system, 104 male reproductive system, 233, 234 respiratory system, 201, 202, 203 urinary system, 191, 192, 194 Muscularis externa digestive system, 139, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180 lymphatic system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myeloblast, 54 Myocopithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofilaments, 57 Myoid cell, 91, 104, 226, 230, 231 Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 287, 27, 4, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 | |
| 265 lymphatic system, 104 male reproductive system, 233, 234 respiratory system, 201, 202, 203 urinary system, 191, 192, 194 Muscularis externa digestive system, 139, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180 lymphatic system, 29, 93, 103 respiratory system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myelocyte, 54 Myocardium, 78, 84, 85, 86 Myoepithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 6 | |
| 265 lymphatic system, 104 male reproductive system, 233, 234 respiratory system, 201, 202, 203 urinary system, 191, 192, 194 Muscularis externa digestive system, 139, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180 lymphatic system, 29, 93, 103 respiratory system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myelocyte, 54 Myocardium, 78, 84, 85, 86 Myoepithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 6 | |
| Imphatic system, 104 | |
| respiratory system, 201, 202, 203 urinary system, 191, 192, 194 Muscularis externa digestive system, 139, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180 lymphatic system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myelocyte, 54 Myocardium, 78, 84, 85, 86 Myopithelial cell cye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 164, 166, 167, 168, 169, 170, 170, 170, 170, 170, 170, 170, 170 | |
| respiratory system, 201, 202, 203 urinary system, 191, 192, 194 Muscularis externa digestive system, 139, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180 lymphatic system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myocardium, 78, 84, 85, 86 Myoepithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 dermis, 76 dura mater, 67 endoneurium, 73 epidermis, 76 epineurium, 72 epithelium, 69 fibrous astrocyte, 69 fibrous astrocyte, 69 fibrous astrocyte, 69 fibrous apace, 75 fourth ventricle, 69 ganglia, 65 glycogen body, 75, 76 granular layer, 67, 69 gray matter, 66, 70, 71, 75, 76 Herbst corpuscle, 76 inner core, 75 mesothelium, 67 molecular layer, 67, 69 gray matter, 66, 70, 71, 75, 76 Meissner's plexus, 72 meninges, 65, 68 mesenteric blood vessel, 75 mesothelium, 69 fibrous astrocyte, 69 fibrous astrocyte, 69 fibrous astrocyte, 69 fibrous apace, 75 fourth ventricle, 69 ganglia, 65 glycogen body, 75, 76 granular layer, 67, 69 gray matter, 66, 70, 71, 75, 76 Herbst corpuscle, 76 miner core, 75 mesothelium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, 76 mesothelium, 67 merve, 72, 74, 75 neuroglial cells, 65 neuroglial | |
| urinary system, 191, 192, 194 Muscularis externa digestive system, 139, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180 | |
| Muscularis externa digestive system, 139, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180 | |
| digestive system, 139, 152, 153, 154, 155, 156, 157, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 178, 179, 180 lymphatic system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 minary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myocardium, 78, 84, 85, 86 Myocardium, 78, 84, 85, 86 Myocardium, 78, 84, 85, 86 Myopithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasolabial gland, 110 Nasopharynx, 196, 199 Nasopharynx, 196, 199 Narous system, 72, 74, 75 optic nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 endoneurium, 73 epidermis, 76 epineurium, 72 epithelium, 69 fibrous astrocyte, 69 fibrous atrocyte, 69 fibrous troches, 67 fourth ventricle, 69 ganglia, 65 glycogen body, 75, 76 menerole, 67, 69 mul | |
| 170, 171, 178, 179, 180 lymphatic system, 92, 93, 103 respiratory system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myocardium, 78, 84, 85, 86 Myophithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myometrium, 244, 251, 252, 253, 254, 257 Masal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 fluid-filled space, 75 fourth ventricle, 69 gray matter, 66, 70, 71, 75, 76 Herbst corpuscle, 76 inner core, 75 Meissner's plexus, 72 meninges, 65, 68 mesenteric blood vessel, 75 mesorthelium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, 76 neurons, 65, 69 nissl granular layer, 67, 69 nisellodation of the properties of t | |
| 170, 171, 178, 179, 180 lymphatic system, 92, 93, 103 respiratory system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myocardium, 78, 84, 85, 86 Myophithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myometrium, 244, 251, 252, 253, 254, 257 Masal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 fluid-filled space, 75 fourth ventricle, 69 gray matter, 66, 70, 71, 75, 76 Herbst corpuscle, 76 inner core, 75 Meissner's plexus, 72 meninges, 65, 68 mesenteric blood vessel, 75 mesorthelium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, 76 neurons, 65, 69 nissl granular layer, 67, 69 nisellodation of the properties of t | |
| lymphatic system, 92, 93, 103 respiratory system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myelocyte, 54 Myocardium, 78, 84, 85, 86 Myocardium, 78, 84, 85, 86 Myophthelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofilaments, 57 Myometrium, 244, 251, 252, 253, 254, 257 Masal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 epithelium, 69 fibrous astrocyte, 69 fluid-filled space, 75 fourth ventricle, 69 ganglia, 65 glycogen body, 75, 76 granular layer, 67, 69 molecular layer, 67, 69 molecular layer, 67, 69 molecular layer, 67, 69 monlecular layer, 67 monlecular layer, 67 monlecular layer, 67 monlecular layer, 6 | |
| respiratory system, 200 Muscularis mucosae digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myocardium, 78, 84, 85, 86 Myocardium, 78, 84, 85, 86 Myocpithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 91, 104, 226, 230, 231 Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 fibrous astrocyte, 69 fluid-filled space, 75 fourth ventricle, 69 ganglia, 65 glycogen body, 75, 76 granular layer, 67, 69 granular layer, 67, 69 fourth ventricle, 69 ganglia, 65 glycogen body, 75, 76 granular layer, 67, 69 manuter core, 75 Meissner's plexus, 72 meninges, 65, 68 mesenteric blood vessel, 75 mesothelium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, 76 merve fascicles, 73, 74 nerve fiber, 71, 72, 74 nerve fiber, 71, 72, 74 neuroglial cells, 65 neuromuscular spindle, 75 nuclear chain fiber, 75 nucleous, 71 nucleus, 71 nucleus, 71 nucleus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myelocyte, 54 Myocardium, 78, 84, 85, 86 Myoepithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 fourth ventricle, 69 ganglia, 65 glycogen body, 75, 76 granular layer, 67, 69 gray matter, 66, 70, 71, 75, 76 Herbst corpuscle, 76 Herbst corpuscle, 76 miner core, 75 Meissner's plexus, 72 meninges, 65, 68 mesenteric blood vessel, 75 mesothelium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, 79 myelin sheath, 66, 73, 74 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neuroglial cells, 65 neuromuscular spindle, 75 nuclear bag fiber, 75 nucleolus, 71 nucleus, 71 nucleus, 71 nucleus, 71 nucleus, 71 pacinian corpuscle, 75 pacinian corpuscle, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| digestive system, 139, 152, 153, 154, 156, 157, 158, 159, 160, 161, 162, 164, 166, 167, 168, 169, 170, 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myelocyte, 54 Myocardium, 78, 84, 85, 86 Myoepithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 fourth ventricle, 69 ganglia, 65 glycogen body, 75, 76 granular layer, 67, 69 gray matter, 66, 70, 71, 75, 76 Herbst corpuscle, 76 Herbst corpuscle, 76 miner core, 75 Meissner's plexus, 72 meninges, 65, 68 mesenteric blood vessel, 75 mesothelium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, 79 myelin sheath, 66, 73, 74 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neuroglial cells, 65 neuromuscular spindle, 75 nuclear bag fiber, 75 nucleolus, 71 nucleus, 71 nucleus, 71 nucleus, 71 nucleus, 71 pacinian corpuscle, 75 pacinian corpuscle, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myeloblast, 54 Myocardium, 78, 84, 85, 86 Myocardium, 78, 84, 85, 86 Myocardium, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve Nerve Nerve Nerve North (10, 120, 121, 123) Double (11, 104, 120, 121, 123) Double (12, 124, 125) Double (13, 120, 121, 123) Double (14, 120, 121, 123) Double (15, 67, 68, 69) Double (16, 73, 74) Double (17, 75, 75) Double (18, 65) Double (19, 104, 226, 230, 231) Double (19, 104, 226, | |
| 178, 180 urinary system, 192 Myelin sheath, 66, 73, 74 Myeloblast, 54 Myeloblast, 54 Myocardium, 78, 84, 85, 86 Myocardium, 78, 84, 85, 86 Myocardium, 78, 84, 85, 86 Myocardium, 106, 110, 120, 121, 123 pigmented myoepithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myometrium, 244, 251, 252, 253, 254, 257 Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 parasympathetic ganglion, 72 | |
| Myelin sheath, 66, 73, 74 gray matter, 66, 70, 71, 75, 76 Myeloblast, 54 Herbst corpuscle, 76 Myelocyte, 54 inner core, 75 Myocardium, 78, 84, 85, 86 Meissner's plexus, 72 Myoepithelial cell meninges, 65, 68 eye, 271, 273 mesenteric blood vessel, 75 integument, 106, 110, 120, 121, 123 mesothelium, 67 pigmented myoepithelial cell, 268 molecular layer, 67, 68, 69 Myofibrils, 61, 62, 63, 64, 86 multipolar neuron, 65, 70, 71, 75, Myoid cell, 91, 104, 226, 230, 231 nerve, 72, 74, 75 Myometrium, 244, 251, 252, 253, 254, 257 nerve fascicles, 73, 74 nerve, 72, 74, 75 neuroglial cells, 65 Nasal cavity, 195, 196, 198, 199, 205 neuromuscular spindle, 75 Nasolabial gland, 110 neurons, 65, 69 Nasolabial gland, 110 neurons, 65, 69 Nasolabial gland, 199 nissl granule, 71, 72 Nerve nuclear bag fiber, 75 bone, 39 nuclear bag fiber, 75 cochlear nerve, 289, 290 nucleous, 71 facial nerve, 271, 275, 281 pacinian corpuscle, 75 optic nerve, 271, 275, 281 | |
| Myeloblast, 54 Myelocyte, 54 Myelocyte, 54 Myocardium, 78, 84, 85, 86 Meissner's plexus, 72 Myoepithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofibrils, 61, 62, 63, 64, 86 Myometrium, 244, 251, 252, 253, 254, 257 Myometrium, 244, 251, 252, 253, 254, 257 Masal cavity, 195, 196, 198, 199, 205 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervesystem, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 Meissner's plexus, 72 meinner core, 75 Meissner's plexus, 72 mesingexus, 72 mesingex blood vessel, 75 mesothelium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, myelin sheath, 66, 73, 74 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neurongial cells, 65 neuromscular spindle, 75 nuclear bag fiber, 75 nuclear chain fiber, 75 nucleolus, 71 nucleous, 71 nucleous, 71 neurous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 | |
| Myelocyte, 54 inner core, 75 Myocardium, 78, 84, 85, 86 Meissner's plexus, 72 Myoepithelial cell meninges, 65, 68 eye, 271, 273 mesenteric blood vessel, 75 integument, 106, 110, 120, 121, 123 mesothelium, 67 pigmented myoepithelial cell, 268 molecular layer, 67, 68, 69 Myofibrils, 61, 62, 63, 64, 86 multipolar neuron, 65, 70, 71, 75, Myofilaments, 57 myelin sheath, 66, 73, 74 Myoid cell, 91, 104, 226, 230, 231 nerve, 72, 74, 75 Myometrium, 244, 251, 252, 253, 254, 257 nerve fascicles, 73, 74 Nasal cavity, 195, 196, 198, 199, 205 neuroglial cells, 65 Nasal concha, 198 neuromuscular spindle, 75 Nasolabial gland, 110 neurons, 65, 69 Nasopharynx, 196, 199 nissl granule, 71, 72 Nerve nuclear bag fiber, 75 bone, 39 nuclear chain fiber, 75 cochlear nerve, 289, 290 nucleus, 71 facial nerve, 287 nucleus, 71 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 pacinian corpuscle, 75 respiratory system, 205, 206 pacinian corpuscle, 75 small encapsulated nerve endings, 75 parasympathetic ganglion, 72< | |
| Myocardium, 78, 84, 85, 86 Myoepithelial cell eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofilaments, 57 Myoid cell, 91, 104, 226, 230, 231 Myometrium, 244, 251, 252, 253, 254, 257 Masal cavity, 195, 196, 198, 199, 205 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 Meissner's plexus, 72 meninges, 65, 68 mesenteric blood vessel, 75 mesothelium, 67 mesothelium, 67 melium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, 74 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neuroglial cells, 65 neuromuscular spindle, 75 neuromuscular spindle, 75 nuclear bag fiber, 75 nuclear bag fiber, 75 nucleous, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| Myoepithelial cell meninges, 65, 68 eye, 271, 273 mesenteric blood vessel, 75 integument, 106, 110, 120, 121, 123 mesothelium, 67 pigmented myoepithelial cell, 268 molecular layer, 67, 68, 69 Myofibrils, 61, 62, 63, 64, 86 multipolar neuron, 65, 70, 71, 75, Myofilaments, 57 myelin sheath, 66, 73, 74 Myoid cell, 91, 104, 226, 230, 231 nerve, 72, 74, 75 Myometrium, 244, 251, 252, 253, 254, 257 nerve fascicles, 73, 74 Nasal cavity, 195, 196, 198, 199, 205 neuroglial cells, 65 Nasal concha, 198 neuromuscular spindle, 75 Nasolabial gland, 110 neurons, 65, 69 Nasopharynx, 196, 199 nissl granule, 71, 72 Nerve nuclear bag fiber, 75 bone, 39 nuclear chain fiber, 75 cochlear nerve, 289, 290 nucleolus, 71 facial nerve, 287 nucleolus, 71 nervous system, 72, 74, 75 outer core, 75 optic nerve, 271, 275, 281 pacinian corpuscle, 75 respiratory system, 205, 206 pancreas, 75 small encapsulated nerve endings, 75 parasympathetic ganglion, 72 | |
| eye, 271, 273 integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofilaments, 57 Myoid cell, 91, 104, 226, 230, 231 Myometrium, 244, 251, 252, 253, 254, 257 Masal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 mesothelium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, myelin sheath, 66, 73, 74 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve, 72, 74, 75 neuroglial cells, 65 neuromuscular spindle, 75 neuroaus granule, 71, 72 nuclear bag fiber, 75 nuclear chain fiber, 75 nucleous, 71 nucleous, 71 nucleous, 71 pacinian corpuscle, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| integument, 106, 110, 120, 121, 123 pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofilaments, 57 Myoid cell, 91, 104, 226, 230, 231 Myometrium, 244, 251, 252, 253, 254, 257 Masal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 mesothelium, 67 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, myelin sheath, 66, 73, 74 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neuroglial cells, 65 neuromuscular spindle, 75 neurons, 65, 69 nissl granule, 71, 72 nuclear bag fiber, 75 nucleolus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| pigmented myoepithelial cell, 268 Myofibrils, 61, 62, 63, 64, 86 Myofilaments, 57 Myoid cell, 91, 104, 226, 230, 231 Myometrium, 244, 251, 252, 253, 254, 257 Masal cavity, 195, 196, 198, 199, 205 Nasolabial gland, 110 Nasopharynx, 196, 199 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 molecular layer, 67, 68, 69 multipolar neuron, 65, 70, 71, 75, myelin sheath, 66, 73, 74 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neuroglial cells, 65 neuromuscular spindle, 75 neurons, 65, 69 nissl granule, 71, 72 nuclear bag fiber, 75 nucleolus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| Myofibrils, 61, 62, 63, 64, 86 Myofilaments, 57 Myoid cell, 91, 104, 226, 230, 231 Myometrium, 244, 251, 252, 253, 254, 257 Masal cavity, 195, 196, 198, 199, 205 Nasolabial gland, 110 Nasopharynx, 196, 199 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nerve gascicles, 73, 74 nerve fiber, 71, 72, 74 neuroglial cells, 65 neuromuscular spindle, 75 neurons, 65, 69 nissl granule, 71, 72 nuclear bag fiber, 75 nuclear chain fiber, 75 nucleolus, 71 nucleus, 71 nucleus, 71 outer core, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 myelin sheath, 66, 73, 74 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neuroglial cells, 65 neuromuscular spindle, 75 neurolas granule, 71, 72 nuclear bag fiber, 75 nucleolus, 71 outer core, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| Myofilaments, 57 Myoid cell, 91, 104, 226, 230, 231 Myometrium, 244, 251, 252, 253, 254, 257 Msaal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 myelin sheath, 66, 73, 74 nerve, 72, 74, 75 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neurongial cells, 65 neuromuscular spindle, 75 neurons, 65, 69 nissl granule, 71, 72 nuclear bag fiber, 75 nuclear chain fiber, 75 nucleus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| Myoid cell, 91, 104, 226, 230, 231 Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nerve asystem, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 nerve, 72, 74, 75 nerve, 72, 74, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neurongial cells, 65 neuromuscular spindle, 75 neurons, 65, 69 nissl granule, 71, 72 nuclear bag fiber, 75 nucleolus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | 76 |
| Myometrium, 244, 251, 252, 253, 254, 257 Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 nerve fascicles, 73, 74 nerve fiber, 71, 72, 74 neurongial cells, 65 neuromuscular spindle, 75 neurons, 65, 69 nissl granule, 71, 72 nuclear bag fiber, 75 nucleolus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 parasympathetic ganglion, 72 | |
| nerve fiber, 71, 72, 74 Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 nervous nerve fiber, 71, 72, 74 neuroglial cells, 65 neuromuscular spindle, 75 neurons, 65, 69 nissl granule, 71, 72 neuroglial cells, 65 neu | |
| Nasal cavity, 195, 196, 198, 199, 205 Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 neuroglial cells, 65 neuromuscular spindle, 75 neurons, 65, 69 nissl granule, 71, 72 nuclear bag fiber, 75 nuclear chain fiber, 75 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pancreas, 75 parasympathetic ganglion, 72 | |
| Nasal concha, 198 Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve nuclear bag fiber, 75 nuclear chain fiber, 75 nuclear, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 neuromuscular spindle, 75 neuromuscular spindle, 75 neurons, 65, 69 nissl granule, 71, 72 nuclear bag fiber, 75 nuclear chain fiber, 75 nucleolus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pancreas, 75 pancreas, 75 parasympathetic ganglion, 72 | |
| Nasolabial gland, 110 Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 neurons, 65, 69 nissl granule, 71, 72 nuclear bag fiber, 75 nuclear chain fiber, 75 nucleolus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pancreas, 75 parasympathetic ganglion, 72 | |
| Nasopharynx, 196, 199 Nerve bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 nissl granule, 71, 72 nuclear bag fiber, 75 nuclear chain fiber, 75 nucleolus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pancreas, 75 parasympathetic ganglion, 72 | |
| Nerve nuclear bag fiber, 75 bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 nuclear bag fiber, 75 nuclear chain fiber, 75 nucleolus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pancreas, 75 parasympathetic ganglion, 72 | |
| bone, 39 cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 nuclear chain fiber, 75 nucleous, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pancreas, 75 parasympathetic ganglion, 72 | |
| cochlear nerve, 289, 290 facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 nucleus, 71 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pancreas, 75 parasympathetic ganglion, 72 | |
| facial nerve, 287 nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 nucleus, 71 outer core, 75 pacinian corpuscle, 75 pancreas, 75 parasympathetic ganglion, 72 | |
| nervous system, 72, 74, 75 optic nerve, 271, 275, 281 respiratory system, 205, 206 small encapsulated nerve endings, 75 outer core, 75 pacinian corpuscle, 75 pancreas, 75 parasympathetic ganglion, 72 | |
| optic nerve, 271, 275, 281 pacinian corpuscle, 75 respiratory system, 205, 206 pancreas, 75 small encapsulated nerve endings, 75 parasympathetic ganglion, 72 | |
| respiratory system, 205, 206 pancreas, 75 small encapsulated nerve endings, 75 parasympathetic ganglion, 72 | |
| small encapsulated nerve endings, 75 parasympathetic ganglion, 72 | |
| Nerve fascisles 73 74 parineurium 72 73 74 | |
| | |
| Normal flow 71 72 74 275 277 201 | |
| Nerve fiber, 71, 72, 74, 275, 276, 281 periosteum, 67 | |
| Nervous system, 65–76 perivascular space, 69 adipose tissue, 72 pia mater, 67, 68, 69 | |
| adipose tissue, 72 pia mater, 67, 68, 69 arachnoid layer, 68 plexus, 72 | |
| | |
| | |
| astrocyte, 69 pyramidal cell, 67 Auerbach's plexus, 72 satellite cell, 72 | |
| axon, 65, 73, 74, 76 Schwann cell, 66, 74 | |
| axon, 65, 75, 74, 76 axon hillock, 71 Sharpey's fiber, 76 | |
| blood vessel, 67, 74 skeletal muscle, 75 | |
| bone, 67, 76 small artery, 68 | |
| brain sand, 69 small encapsulated nerve endings, | 75 |
| capillary, 69 space artifact, 73, 76 | |
| capsule, 75 spinal cord, 70 | |
| cell body, 65 subarachnoid space, 67 | |

| submucosa, 72 | Palatine ridge, 135 |
|---|---|
| unipolar neuron, 65, 76 | Palatine tonsil, 90, 93, 94 |
| ventral root, 76 | Palpebral conjunctiva, 269, 271, 272, 273, 277, |
| venule, 69 | 278 |
| white matter, 66, 67, 70, 75, 76 | PALS (periarterial lymphatic sheath), 90, 100 |
| word roots, 66 | Pancreas, 75, 81, 141, 142, 176, 177, 182 |
| Neuroglial cell (pituicyte), 65 | Paneth cell, 165 |
| Neurohypophysis, 211, 213 | Papilla, 107, 139, 146, 157 |
| Neuromuscular spindle, 75, 205 | Papillary duct, 183, 189, 190, 191 |
| Neurons, 65, 69, 86 | Papillary layer, 106, 111, 112 |
| Neutrophil | Parabasal cell, 245, 260, 261 |
| | |
| blood, 42, 44, 45, 46, 47, 48, 49, 50 | Parabronchi, 197, 208, 209 |
| bone marrow, 56 | Paraepiglottic tonsil, 90, 93 |
| connective tissue, 20, 22 | Paraffin procedure, 3, 4 |
| female reproductive system, 260, 261 | Parafollicular (C) cell, 212, 218 |
| Neutrophilic band cell, 55, 56 | Parasympathetic ganglion, 72 |
| Neutrophilic metamyelocyte, 56 | Parathyroid gland, 212, 213, 218, 219, 223 |
| Nictitating membrane, 269, 270, 276 | Parenchyma cell, 86 |
| Nissl granule, 71, 72 | Parietal cell, 140, 158, 159, 160, 161 |
| Nonpigmented epithelium, 273, 279, 280 | Parietal layer, 184, 188 |
| Nonsinusal spleens, 90 | Parietal pleura, 205 |
| Nose, 109 | Parietal prepuce, 240 |
| Nuclear bag fiber, 75 | Parotid gland, 148, 149 |
| Nuclear chain fiber, 75 | Pars ciliaris retinae, 268, 271, 272, 274 |
| Nuclei stack, 132, 135 | Pars convoluta, 183, 186 |
| Nucleolus, 71 | Pars distalis, 211, 212, 213, 215, 216, 217, 221, 222 |
| Nucleus, 71 | Pars intermedia, 211, 212, 215, 216 |
| Trueleus, 7 I | Pars iridica retinae, 268, 271 |
| Odontoblast, 144 | Pars nervosa, 212, 213, 215, 216, 221 |
| Olfactory cell, 198, 206 | Pars radiata, 183, 186, 189 |
| Olfactory epithelium, 198, 206 | Pars tuberalis, 211, 212, 213, 215, 216, 217, 221, |
| Omasum, 156, 157 | 222 |
| | Pecten, 270, 281 |
| Oocyte, 247, 261, 262 | |
| Optic disc, 269, 271, 275 | Pectinate ligament (uveal meshwork), 268, 270, 272, |
| Optic nerve, 271, 275, 281 | 273, 276, 279, 280 |
| Ora ciliaris retinae, 268, 271, 274 | Pelvic urethra, 227 |
| Oral cavity, 135 | Penicillus, 90 |
| Orcein stains, 5 | Penile urethra, 227, 237, 238 |
| Organ of Corti, 284, 289, 290 | Penis, 227, 236, 237 |
| Oropharynx, 140, 151, 152, 196 | Periarterial lymphatic sheath (PALS), 90, 100 |
| Orthochromatophilic erythroblast (metarubricyte), 54, | Pericardial cavity, 86 |
| 55, 56 | Pericardium, 86 |
| Os penis, 227, 236, 237 | Perichondrium, 27, 29, 201 |
| Osseous spiral lamina, 284, 289 | Perilobular collecting duct, 185 |
| Osteoblast, 31, 33, 36, 37, 38, 54, 55 | Perilymph, 284 |
| Osteoclast, 32, 33, 36, 38, 54, 55 | Perilymphatic space, 290 |
| Osteocyte, 32, 33, 36, 37, 38, 39 | Perimetrium (serosa), 244 |
| Osteoid, 32, 33 | Perimysium, 58, 61 |
| Osteon, 32, 38 | Perineurium, 72, 73, 74 |
| Osteoprogenitor (osteogenic) stem cell, 31 | Periodontal ligament, 144 |
| Otolith (otoconia, statoconia), 284, 288 | Perioplic dermis, 128, 130 |
| Otolithic membrane, 284, 288 | Perioplic epidermis, 128, 130 |
| Outer core, 75 | Perioplic region of hoof, 128, 129 |
| Outer nuclear layer, 275, 276, 281 | Periosteum, 31, 34, 35, 37, 39, 67, 135 |
| Outer plexiform layer, 275, 276, 281 | Peritoneum, 225, 227 |
| Outer root sheath, 116, 118, 119, 120 | Perivascular space, 69 |
| Ovaries, 243, 246, 247, 248, 249, 261, 262 | Perivitelline membrane, 246, 262 |
| Oviduct, 244, 246, 249, 250, 263, 264, 265 | |
| Oviduel, 277, 270, 277, 200, 200, 207, 200 | Pessulus, 207, 208 Peyor's patch, 89, 92, 140 |
| Paginian corpusale 75 111 141 | Peyer's patch, 89, 92, 140 |
| Pacinian corpuscle, 75, 111, 141 | Pharynx, 195, 196, 197 |
| Palatine bone, 145 | Pia mater, 67, 68, 69 |

| D' | D: (11:1 406 247 |
|---|---|
| Pig | Primary follicle, 106, 247 |
| blood, 47 | Primary hair, 118 |
| cardiovascular system, 79, 80, 81, 82, 83, 84, 86, 87 | Primary lamina, 128 |
| connective tissue, 24 | Primary oocyte, 243 |
| digestive system, 145, 151, 153, 154, 158, 165, 170, | Primary spermatocyte, 225, 230, 231, 240, 241 |
| 174, 175, 176, 177 | Primordial follicle, 243, 247 |
| | |
| epithelium, 14, 17, 18 | Principal (chief) cell, 212, 219 |
| eye, 273, 274, 275, 277, 278 | Proctodeum, 181 |
| female reproductive system, 248, 257 | Proerythrocyte, 53 |
| histology, 10 | Proestrus, 244, 245, 260 |
| integument, 110, 117, 121 | Promyelocyte, 54, 56 |
| lymphatic system, 92, 95, 100 | Prostate gland, 226, 235 |
| male reproductive system, 229, 230, 234, 236, 238, | Proventriculus, 141, 178, 179 |
| 240 | Proximal convoluted tubule, 183, 187, 188, 189, 193, 194 |
| | |
| muscle, 64 | Pseudostratified columnar epithelium, 12, 13, 16, 196, |
| nervous system, 72, 73, 74 | 263 |
| respiratory system, 203, 204 | Pseudostratified epithelium |
| urinary system, 188, 189, 192 | ear, 288 |
| Pigmented columnar cell, 268 | female reproductive system, 253, 264, 265 |
| Pigmented epithelium, 273, 275, 276, 279, 280, 281 | lymphatic system, 104 |
| Pigmented granule, 286 | male reproductive system, 232, 233, 234, 235, 242 |
| Pigmented myoepithelial cell, 268 | respiratory system, 198, 199, 200, 201, 202, 205, |
| Pineal gland, 212, 213, 218, 222, 223 | 206, 208 |
| | |
| Pineal stalk, 218 | urinary system, 194 |
| Pinealocyte, 212, 218 | Pulmonary artery, 83, 84, 85 |
| Pinna (auricle), 283 | Pulmonary vein, 205 |
| cartilage, 29 | Pulmonic (semilunar) valve, 85 |
| integument, 118 | Pulp artery, 90 |
| Pituicyte (neuroglial cell), 212, 222 | Pupil, 268, 271 |
| Pituitary gland (hypophysis), 211–212, 213, 215, 216, | Puppy. See Dog |
| 217, 221, 222 | Purkinje cell, 67, 69, 86 |
| Placenta, 244, 254, 255, 256, 257, 258 | Pyknotic cell, 116, 120 |
| Placental labyrinth, 254, 256 | Pyknotic nucleus, 205 |
| | |
| Placentome, 257 | Pyloric gland, 161, 162 Pyloric gland region, 140, 161, 162, 163 |
| Planum nasale, 108 | Pyloric gland region, 140, 161, 162, 163 |
| Planum nasolabiale, 110, 111 | Pyramidal cell, 67 |
| Planum rostrale, 110 | |
| Plasma, 41, 44, 80, 87 | Queen. See Cat |
| Plasma cell, 20, 22, 23, 55, 79, 200, 282 | Quill (calamus), 107, 133, 134 |
| Platelet, 41, 43, 44, 45, 46, 47, 49, 50 | |
| Plexus, 72 | Rachis, 107 |
| Plica (folds), 178 | Ram. See Sheep |
| Pluripotent stem cell, 53 | Raphe, 290 |
| Pointer in ocular, 5 | Rathke's pouch, 212, 215, 216 |
| Polychromatophilic erythroblast (rubricyte), 54, 56 | Rectoanal junction, 170, 171, 172 |
| Polymorphonuclear leukocyte, 43 | Rectum, 140, 170 |
| | |
| Pore, 178 | Red blood cell. See Erythrocyte |
| Portal tract (area), 141, 174 | Red pulp, 90, 91, 99, 100, 101, 103 |
| Portal vein, 82, 173, 174 | Regenerating gland, 263, 264 |
| Postcapillary venule, 90, 97 | Reissner's (vestibular) membrane, 284, 289 |
| Posterior chamber, 269, 271, 272, 279 | Relocating structures, 8 |
| Posterior epithelium, 267, 274, 280 | Renal artery, 82 |
| Precementum, 144 | Renal corpuscle, 184, 186, 187, 193 |
| Predentin, 144 | Renal papilla, 183, 189, 190 |
| Preen (uropygial) gland, 107, 137, 138 | Renal pelvis, 183 |
| Preparation, specimen, 3–5, 4 | Resorcin fuchsin stain, 5 |
| Prepuce, 227, 245 | Respiratory bronchiole, 203 |
| Preputial gland, 240 | Respiratory epithelium, 198, 205 |
| Primary bronchi, 95, 201, 208 | Respiratory system, 195–209 |
| Primary duct, 178 | adipose tissue, 199, 200, 201 |
| | |
| Primary fold, 258, 263, 264, 265 | air capillary, 197, 209 |

| air sac, 197 | nerve, 205, 206 |
|--|---|
| alveolar duct, 203 | neuromuscular spindle, 205 |
| alveolar sac, 203 | olfactory cell, 198, 206 |
| alveolar septum, 204 | olfactory epithelium, 198, 206 |
| alveolus, 195, 202, 203, 204, 205 | oropharynx, 196 |
| arytenoid cartilage, 199, 200 | parabronchi (tertiary bronchi), 197, 208, 209 |
| atrium (air vesicle), 197, 208, 209 | parietal pleura, 205 |
| basal cell, 198, 205, 206 | perichondrium, 201 |
| bone, 198, 209 | pessulus, 207, 208 |
| bony tracheal ring, 206, 207 | pharynx, 195, 196, 197 |
| Bowman's gland, 198, 206 | plasma cell, 200 |
| bronchi, 195, 196, 202, 204 | primary bronchi, 201, 208 |
| bronchial cartilage, 208 | pseudostratified columnar epithelium, 196 |
| bronchial ring, 207 | pseudostratified epithelium, 198, 199, 200, 201, 202, |
| bronchiole, 196, 202, 203, 204 | 205, 206, 208 |
| cartilage, 198 | pulmonary vein, 205 |
| cartilaginous tracheal ring, 206, 207 | pyknotic nucleus, 205 |
| cavernous vein, 198 | respiratory bronchiole, 203 |
| cavity of air sac, 209 | respiratory proficilione, 203 |
| | |
| chicken, 196–197 | secondary bronchi, 208, 209 |
| ciliated pseudostratified columnar epithelium, 197 | sensory hairs, 198 |
| columnar epithelium, 202 | septum, 205 |
| connective tissue (lamina of), 209 | serous acinus, 199 |
| cuboidal epithelium, 203 | serous gland, 198, 200, 202 |
| duct, 199 | simple squamous epithelium, 209 |
| elastic band, 202, 204 | skeletal muscle, 199, 200, 206 |
| elastic cartilage, 199 | smooth muscle, 203, 205, 208, 209 |
| elastic fiber, 200, 201, 203, 208 | stratified columnar epithelium, 208 |
| epidermis, 205 | stratified squamous epithelium, 196, 198, 199, 200, |
| epiglottis, 199 | 206, 207, 208 |
| epithelium, 209 | striated duct, 199 |
| esophagus, 200 | submucosa, 199, 200 |
| external tymponic membrane, 207, 208 | supporting cell, 198, 206 |
| extrapulmonary primary bronchi, 197 | syrinx, 197, 207, 208 |
| fibroelastic membrane, 201 | taste buds, 199 |
| glottis, 199 | trachea, 195, 196, 197, 200, 201, 206, 207 |
| goblet cell, 198, 200, 201, 202 | tracheal gland, 201 |
| granulocyte, 209 | trachealis muscle, 200, 201 |
| humerus, 209 | turbinate cartilage, 205 |
| hyaline cartilage, 200, 201, 202 | tympanic membrane, 208 |
| intercalated duct, 199 | Type II alveolar cell, 204 |
| intercostal muscle, 205 | vestibular epithelium, 205 |
| intermediate syringeal cartilage, 207, 208 | vestibule, 195 |
| internal tymponic membrane, 207, 208 | visceral pleura, 196, 204 |
| intrapulmonary primary bronchi (mesobronchi), 197 | vocal fold, 200 |
| lamina propria, 196, 199, 200, 201, 202, 205 | vocal ligament, 200 |
| larynx, 195, 196, 197 | word roots, 197 |
| lungs, 196, 209 | Resting gland, 263, 264 |
| lymphatic nodule, 202 | Rete ovarii, 243, 249 |
| lymphatic tissue, 208 | Rete testis, 226, 228, 229, 231, 232, 240, 241 |
| macrophage (histiocyte), 203, 204, 209 | Reticular cell, 92, 97, 98, 103 |
| medial bronchial wall, 207 | Reticular fiber, 20, 25, 96, 182, 191 |
| mesothelium, 204 | Reticular layer, 106, 111, 112 |
| mixed gland, 199, 200, 201, 202 | Reticular structure, 91, 104 |
| mucous cell, 209 | Reticular tissue, 20, 25 |
| mucous gland, 205, 206 | Reticulate scale, 136 |
| muscularis, 201, 202, 203 | Reticulocyte, 54, 56 |
| muscularis externa, 200 | Reticulum, 155, 156 |
| nasal cavity, 195, 196, 198, 199, 205 | Retina, 267, 268, 270, 271, 274, 275, 276, 281, 282 |
| nasal concha, 198 | Retractor penis muscle, 238 |
| nasopharynx, 196, 199 | Right auricle, 84 |
| | , |

| Rods, 275, 276, 281 | Separation (space) artifact, 10 |
|--|--|
| Romanovsky stains, 5 | Septa, 90 |
| Rooster. See Chicken | Septum, 101, 104, 205 |
| Rosette, 223 | Serosa |
| Rouleaux, 42, 45, 47 | digestive system, 139, 154, 159, 161, 164, 166, 168, |
| Rubricyte (polychromatophilic erythroblast), 54, 56 | 169, 173, 178, 180 |
| Rumen, 155 | female reproductive system, 250, 263 |
| | lymphatic system, 100, 101 |
| Sacculus, 284, 287 | male reproductive system, 233, 234 |
| Salivary gland, 93, 94, 141, 177 | urinary system, 194 |
| Sarcolemma, 58 | Serous acinus |
| Sarcoplasm, 57 | digestive system, 143, 149, 150, 175 |
| Satellite cell, 72 | eye, 277 |
| Scala media, 290 | lymphatic system, 93 |
| Scala tympani, 284, 289, 290 | respiratory system, 199 |
| Scala vestibuli, 284, 289, 290 | Serous cell, 236 |
| Schwann cell, 66, 74 | Serous demilune |
| Sclera, 267, 269, 271, 272, 273, 274, 275, 276, 278, | digestive system, 143, 149, 150, 151 |
| 279 | lymphatic system, 93 |
| Scleral cartilage, 269, 279, 280, 281, 282 | Serous gland |
| Scleral ossicles, 269, 279, 280 | digestive system, 148, 165 |
| Scleral trabecular meshwork, 270, 279 | respiratory system, 198, 200, 202 |
| Scleral venous plexus, 268, 271, 272 | Sertoli cell, 225, 226, 230, 231, 240, 241 |
| Scratches (knife marks), 10 | Serum, 41 |
| Scrotum, 117, 118 | Sex cord, 229 |
| Scutes, 136 | Sharpey's fiber, 39, 40, 76 |
| Sebaceous gland | Sheathed artery, 91, 103 |
| digestive system, 143, 171, 172 | Sheathed capillary, 90 |
| ear, 286 | Sheep |
| eye, 278 | blood, 49 |
| glands of Zeiss, eye, 269 | bone, 40 |
| integument, 106, 109, 114, 115, 117, 118, 119, 120, | cardiovascular system, 79, 80, 84 |
| 121, 122, 124 | connective tissue, 25 |
| male reproductive system, 240 | digestive system, 143, 145, 149, 150, 154, 155, 156, |
| Sebaceous zone, 107, 137 | 157, 160, 166, 177 |
| Secondary bronchi, 208, 209 | endocrine system, 221 |
| Secondary duct, 178, 179 | epithelium, 14, 15, 16, 17 |
| Secondary fold, 258, 263, 265 | eye, 275 |
| Secondary follicle, 106, 244 | integument, 117, 118, 120, 121, 122, 124 |
| Secondary gland, 263 | lymphatic system, 96, 97, 98, 101 |
| Secondary laminae, 128 | male reproductive system, 231, 234, 235, 236, 238, |
| Secondary spermatocyte, 226 | 239 |
| Secretion, 123, 234, 235 | muscle, 59, 63 |
| Secretory acinus, 111 | nervous system, 67, 68, 70, 71 |
| Secretory bleb, 250 | respiratory system, 199, 201, 203 |
| Secretory cell, 121, 123, 236, 264 | urinary system, 187 |
| Secretory gland, 264 | Shell gland (uterus), 246, 262, 265 |
| Secretory portion, sweat gland, 120 | Silver stains, 5, 20 |
| Secretory (striated) duct, 149, 150, 151, 199 | Simple columnar epithelium, 158, 160, 161, 172, 181 |
| Secretory tubule, 282 | Simple epithelia, 11, 12 |
| Secretory unit, 122, 123, 149, 164, 165, 166 | Simple squamous epithelium, 209 |
| Section (specimen) | Single hair follicle, 109 |
| interpreting sections, 5, 6–7 | Sinus capillary, 107, <i>134</i> , <i>135</i> |
| preparation, 3–5, 4 | Sinus pad, 119 |
| sectioning with microtome, 3, 4, 5 | Sinus (tactile) hair, 106 |
| Semicircular canal, 284, 290 | Sinusal spleens, 90 |
| Semicircular duct, 284, 290 | Sinusoid |
| Seminal vesicle, 226, 234, 235 | bone, 36, 38 |
| Seminiferous tubule, 225, 227, 230, 231, 240, 241 | digestive system, 141, 173, 174, 182 |
| Sensory cell, 146, 288 | endocrine system, 217, 220, 221, 222, 223 |
| Sensory hair cell, 283, 284 | vascular sinusoid, 53 |
| | |

| Skeletal muscle | Staining specimen, 4 |
|--|---|
| cardiovascular system, 79, 80 | Stallion. See Horse |
| defined, 57, 58, 61, 63 | Stapes, 284, 287, 288 |
| digestive system, 143, 146, 148, 151, 153, 154, 172, | Statoconia (otolith), 284, 288 |
| 181 | Stellate reticulum, 143, 144 |
| eye, 278 | Stem cell, 120 |
| integument, 111, 114 | Stereocilia, 11, 233 |
| lymphatic system, 93 | Stomach, 140, 141, 158, 159, 161, 163 |
| nervous system, 75 | Straight tubule, 226, 228, 229, 231, 241 |
| respiratory system, 199, 200, 206 | Stratified columnar epithelium, 12, 13, 18, 208, 237, |
| Skin and skin derivatives. See Integument | 277 |
| Skull, 221, 222 | Stratified cuboidal epithelium, 12, 13 |
| Small artery, 68, 77, 79, 80, 81, 169 | Stratified epithelium, 11, 12, 259 |
| Small encapsulated nerve endings, 75 | Stratified squamous epithelium |
| Small intestine, 140 | defined, 12, 13, 17 |
| Small vein, 77, 79, 80 | digestive system, 143, 145, 147, 148, 151, 152, 153, |
| Smegma, 239 | 154, 155, 156, 157, 158, 171, 172, 177, 178, 181 |
| Smooth muscle | eye, 277, 280 |
| cardiovascular system, 79, 84 | female reproductive system, 258, 259 |
| digestive system, 153, 154, 159, 181 | integument, 125 |
| epithelium, 14, 16, 17, 18 | lymphatic system, 93, 94 |
| eye, 278 | male reproductive system, 239, 240 |
| integument, 118, 123, 124, 125 | respiratory system, 196, 198, 199, 200, 206, 207, |
| lymphatic system, 96, 97, 100, 101 | 208 |
| male reproductive system, 232, 233, 236, 237, 238, | Stratum basale |
| 239, 242 | digestive system, 145 |
| muscle, 57, 59 | integument, 106, 108, 109, 110, 112, 113, 114, 117, |
| respiratory system, 203, 205, 208, 209 | 125, 127 |
| urinary system, 187 | Stratum cavernosum (vascular stratum), 227, 235 |
| Smudged cell, 44, 45, 47, 52, 55 | Stratum compactum, 140, 159, 162, 167 |
| Soft palate, 145 | Stratum compactum, 170, 137, 102, 107 |
| Sole, hoof, 128, 129 | digestive system, 145 |
| Sow. See Pig | integument, 105, 108, 110, 112, 113, 114, 117, 125, |
| Space artifact | 127, 132, 133, 134, 135, 136 |
| digestive system, 144, 148 | Stratum germinativum, 132, 133, 134, 135, 136, 137 |
| | Stratum granulosum |
| endocrine system, 219 | |
| eye, 274, 280 female reproductive system, 257 | digestive system, 140, 143, 159, 162, 172 |
| | integument, 105, 108, 112, 113, 117, 127 |
| nervous system, 73, 76 | Stratum intermedium, 144 |
| Spaces of Fontana, 268, 270, 272, 273, 279 | Stratum internum, 128 |
| Spermatids, 225, 230, 231, 240, 241 Spermatogenic cell, 225 | Stratum lucidum, 105, 108, 112 |
| | Stratum medium, hoof, 128, 129, 131 Stratum spinosum |
| Spermatogonium, 225, 230, 231, 240, 241 | digestive system, 145, 146 |
| Spermatozoa, 225, 232, 233, 234, 241, 242 | • • • |
| Sperm-host gland, 246, 265 | integument, 105, 108, 109, 110, 112, 113, 114, 117, |
| Sphincter (constrictor) muscle, 268, 271, 273, 279, 280 | 125, 127 Structure textorium 128 |
| Spinal cord, 70 | Stratum tectorium, 128 |
| Spine, 146 | Stratum vasculare, 244, 251, 252 |
| Spiral colon, 170 | Stria vascularis, 284 |
| Spiral ganglion, 289 | Striated border, 140, 163, 165, 169, 170, 175 |
| Spiral ligament, 284, 289 | Striated duct, 149, 150, 151, 199 |
| Spiral limbus, 284, 289 | Stroma 210 |
| Spiral tunnel, 289 | endocrine system, 219 |
| Spleen, 90, 91, 99, 100, 101, 103 | eye, 273, 274, 280 |
| Splenic artery, 90 | female reproductive system, 246, 247 |
| Spongy bone, 32, 33, 37, 236 | Stroma (substantia propria), 267, 268 |
| Spongy layer, 254, 256 | Subarachnoid space, 67 |
| Spur, 136 | Subcapsular sinus, 90, 94, 96, 98 |
| Squamous cell, 280 | Subcapsular vein, 187 |
| Squamous epithelium, 11, 13, 14, 15 | Subcutis, 105, 108, 114, 136 |
| Stain precipitate, 10 | Sublingual gland, 150, 151 |

| Submandibular gland, 149, 150 | Territorial matrix, 27, 29 |
|--|--|
| Submucosa | Tertiary bronchi (parabronchi), 197 |
| digestive system, 139, 145, 152, 153, 154, 155, 156, | Tertiary duct, 178, 179 |
| 158, 159, 160, 161, 162, 164, 165, 166, 167, 168, | Tertiary fold, 258, 263 |
| 169, 170, 178, 179, 180 | Tertiary (Graafian) follicle, 244, 248 |
| female reproductive system, 250 | Testis, 225, 227, 229, 230, 240, 241 |
| lymphatic system, 92, 93 | Theca externa, 244, 246, 247, 248, 262 |
| nervous system, 72 | Theca folliculi, 244, 247 |
| respiratory system, 199, 200 | Theca interna, 244, 246, 247, 248, 262 |
| urinary system, 192 | Theca lutein cell, 248 |
| Sulcus, 136, 178 | Thrombocyte, 43, 51 |
| Superficial cell, 245, 260, 261 | Thymic (Hassall's) corpuscle, 90, 102 |
| Superficial cortex, 187 | Thymus, 90, 91, 101, 102, 103, 104 |
| Superficial gland of nictitating membrane, 269, | Thyroid gland, 212, 213, 218, 219 |
| 277 | Toluidine blue stains, 5 |
| Superficial intermediate cell, 245, 260, 261 | Tomcat. See Cat |
| Supporting cell | Tomial edge, 135 |
| digestive system, 146 | Tongue, 139, 177 |
| ear, 288, 290 | Tonsil, 90 |
| male reproductive system, 229 | Tonsillar follicle, 90 |
| respiratory system, 198, 206 | Trabecula |
| Suprachoroid layer, 268 | integument, 119 |
| Supraglandular layer, 254 | lymphatic system, 90, 94, 96, 97, 99, 100, |
| Surface epithelium, 179 | 101 |
| Surface groove, 108, 110 | male reproductive system, 235, 239 |
| Surface mucous cell, 159, 161, 163 | Trabecular artery, 90 |
| Sweat gland | Trachea, 195, 196, 197, 200, 201, 206, 207 |
| cardiovascular system, 80 | Tracheal gland, 201 |
| duct of, 110, 111, 112, 118, 120, 121 | Trachealis muscle, 200, 201 |
| eye, 277, 278 | Tracheobronchial lymph node, 95 |
| glands of Moll, eye, 269 | Transferring sections to slide, 4 |
| integument, 108, 109, 115, 116, 117, 120, 121, 122, | Transitional epithelium |
| 123, 124 | defined, 12, 13, 18 |
| merocrine sweat gland, 106, 110, 111, 126 | female reproductive system, 259 |
| secretory portion, sweat gland, 120 | male reproductive system, 235, 236, 239 |
| Syndesmochorial placenta, 245 | urinary system, 191, 192 |
| Synovial cavity, 287 | Transitional layer, 134 |
| Synovial fold, 40 | Transitional zone, 226, 231 |
| Syntrophoblast, 255 | Trichrome stains, 5 |
| Syrinx, 197, 207, 208 | Trophoblastic projection, 255 |
| | Troubleshooting, microscopy, 9 |
| Taenia coli, 140, 169, 170 | Tubal tonsil, 90 |
| Tapetal cell, 276 | Tubular ceruminous gland, 283 |
| Tapetum lucidum, 268, 271, 275, 276 | Tubular gland, 140, 181, 264 |
| Tarsal (Meibomian) gland, 269, 277, 278 | Tubular horn, 128 |
| Tarsus, 269, 277, 278 | Tubular mucous unit, 150 |
| Taste buds | Tubuloacinar gland, 107 |
| digestive system, 139, 141, 146, 147, 177, 178 | Tunica adventitia |
| respiratory system, 199 | cardiovascular system, 77, 80, 81, 82, 83, 85 |
| Teat, 123 | female reproductive system, 252 |
| Teat canal, 125 | Tunica albuginea |
| Teat sinus, 123, 124, 125 | female reproductive system, 243, 246, 247 |
| Tectorial membrane, 284, 289, 290 | male reproductive system, 225, 227, 230, 231, 232, |
| Tegmentum vasculosum, 285, 290 | 237, 238, 239, 241 |
| Temporal bone, 287, 288, 289 | Tunica intima |
| Tendon | cardiovascular system, 77, 81, 82, 83, 84, 85 |
| bone, 39 | female reproductive system, 252 |
| connective tissue, 24, 25 | Tunica media |
| elastic tendon, 134 | cardiovascular system, 77, 80, 81, 82, 83, 84, 85 |
| extensor tendon, 40 | female reproductive system, 252 |
| integument, 137 | Tunica vaginalis, 225, 230, 232 |
| | |

| Turbinate cartilage, 205 | macula densa, 184, 187, 188, 189 |
|---|---|
| Tympanic cavity, 283, 284, 286, 287 | mammals, 183–184 |
| Tympanic membrane | medulla, kidney, 184, 186, 189 |
| ear, 283, 286, 287 | medullary cone, 185, 193 |
| respiratory system, 208 | medullary tracts, 185 |
| Type 1 cell (of macula), 288 | medullary type, 184 |
| Type I (glomus) cell, 87 | mesangial cell, 184, 194 |
| Type II (alveolar) cell, 204 | mucous connective tissue, 191 |
| Type II (sustentacular) cell, 87 | mucous gland, 192 |
| Type if (Susteintaeurar) cent, or | muscularis, 191, 192, 194 |
| Ultimobranchial body, 213 | muscularis muscosae, 192 |
| | |
| Umbilical artery, 83 | papillary duct, 183, 189, 190, 191 |
| Undifferentiated epithelial cell, 91, 104 | parietal layer, 184, 188 |
| Unguinal scale, 137 | pars convoluta, 183, 186 |
| Unipolar neuron, 65, 76 | pars radiata, 183, 186, 189 |
| Urachus, 191 | perilobular collecting duct, 185 |
| Ureter, 184, 185, 191, 192, 194 | proximal convoluted tubule, 183, 187, 188, 189, |
| Urethra | 193, 194 |
| corpus spongiosum (corpus cavernosum urethra), | pseudostratified epithelium, 194 |
| 227, 236, 237 | renal corpuscle, 184, 186, 187, 193 |
| female reproductive system, 259 | renal papilla, 183, 189, 190 |
| male reproductive system, 227, 235, 236, 237, 238, | renal pelvis, 183 |
| 239 | reticular fiber, 191 |
| pelvic urethra, 227 | serosa, 194 |
| penile urethra, 227, 237, 238 | smooth muscle, 187 |
| Urethral epithelium, 259 | subcapsular vein, 187 |
| Urethral pouch, 239 | submucosa, 192 |
| Urethral process, 227, 239 | superficial cortex, 187 |
| Urinary bladder, 184, 192 | transitional epithelium, 191, 192 |
| Urinary space, 184, 187, 188, 189, 194 | urachus, 191 |
| Urinary system, 183–194 | ureter, 184, 185, 191, 192, 194 |
| | urinary bladder, 184, 192 |
| adipose tissue, 189 | |
| adventitia, 191 | urinary space, 184, 187, 188, 189, 194 |
| afferent arteriole, 188, 189 | vasa recta, 184, 189, 190, 191 |
| Bowman's capsule, 184, 187, 188, 194 | visceral layer, 184 |
| brush border, 183, 188 | word roots, 185 |
| capillary, 192 | Urodeum, 181, 246 |
| capsule, 183, 186, 187 | Uropygial (preen) gland, 107, 137, 138 |
| cavity of renal pelvis, 189, 190 | Uterine (endometrial) gland, 79, 244, 256, 257 |
| central vein, 184 | Uterine horn, 251, 252, 253 |
| chicken, 184–185 | Uterus, 244, 253, 256, 265 |
| collecting duct, 194 | Utriculus, 284, 287 |
| collecting tubule, 184, 185, 187, 188, 189, 190, 191, | Uvea (vascular tunic), 267, 268, 270 |
| 194 | Uveal meshwork (pectinate ligament), 268, 270, 272, |
| connective tissue, 187 | 273, 276, 279, 280 |
| cortex, kidney, 183, 186, 188, 193, 194 | Uveal trabecular meshwork, 268, 273 |
| cortical nephrons, 184 | • |
| cortical type, 184 | Vacuolar cell, 246, 261 |
| distal convoluted tubule, 183, 187, 188, 193, 194 | Vacuolated cell, 120, 255 |
| efferent arteriole, 189 | Vacuole, 43, 51, 265 |
| erythrocyte, 191, 194 | Vagina, 245, 246, 259, 262, 265 |
| fat vacuole, 187 | Vaginal smear, 245, 260, 261 |
| glomerulus, 184 | |
| | Valves, 78, 80, 82, 85, 87 |
| Henle's loop, 189, 190, 191, 194 | Vas deferens (ductus deferens), 226, 228, 233, 234, |
| intralobular vein, 193 | 242 V |
| juxtaglomerular apparatus, 184 | Vasa recta, 184, 189, 190, 191 |
| juxtaglomerular cell, 184, 188 | Vasa vasorum, 78, 84 |
| kidney, 187 | Vascular layer, 268, 272, 275, 276 |
| lamina propria, 191, 192 | Vascular sinusoid, 53 |
| loop of Henle, 184 | Vascular stratum (stratum cavernosum), 227, 235 |
| lymphatic tissue, 194 | Vascular tunic (uvea), 267, 268, 270 |

| Vein | Vocal fold, 200 |
|--|--|
| bone, 39 | Vocal ligament, 200 |
| cardiovascular system, 81, 87 | Volkmann's canal, 32, 38, 39 |
| cavernous vein, 198 | Vulva, 245 |
| central vein, digestive system, 173, 174, 182 | • |
| central vein, urinary system, 184 | Wattle, 107 |
| digestive system, 157, 169 | Weigert stain (resorcin fuchsin), 5 |
| female reproductive system, 252 | White blood cell. See Leukocyte |
| intralobular vein, 193 | White line, hoof, 128, 129 |
| medium vein, 78, 82 | White matter, 66, 67, 70, 75, 76 |
| portal vein, 82, 173, 174 | White pulp, 90, 91, 99, 101 |
| pulmonary vein, 205 | Word roots |
| small vein, 77, 79, 80 | blood, 43 |
| subcapsular vein, 187 | bone, 32 |
| Vena cava, 84 | bone marrow, 54 |
| Venous sinus, 90, 99 | cardiovascular system, 78 |
| Ventral boundary of third ventricle (median eminence), | cartilage, 28 |
| 212, 213, 215 | connective tissue, 21 |
| Ventral plate, 137 | digestive system, 142 |
| Ventral root, 76 | ear, 285 |
| Ventriculus (gizzard), 141, 179 | endocrine system, 214 |
| Venule | epithelium, 12 |
| cardiovascular system, 77, 79, 80, 87 | eye, 270 |
| integument, 118 | female reproductive system, 246 |
| muscle, 60 | integument, 128 |
| nervous system, 69 | male reproductive system, 228 |
| Verniers, for relocating structures, 8, 8 | muscle, 58 |
| Vesicles, 104 | nervous system, 66 |
| Vestibular epithelium, 205, 259 | respiratory system, 197 |
| Vestibular fold, 93 | urinary system, 185 |
| Vestibular gland, 259 | Wright stain (Romanovsky), 5 |
| Vestibular (Reissner's) membrane, 284, 289 | Wight stain (Romano voley), s |
| Vestibule | Yolk sphere, 262 |
| ear, 284, 287 | Tolk opinere, 202 |
| female reproductive system, 260 | Z band, 62 |
| male reproductive system, 245 | Zona fasciculata, 213, 220, 221 |
| respiratory system, 195 | Zona glomerulosa (zone multiformis), 213, 220, 221 |
| Vestigal cavity, 212 | Zona intermedia, 213, 220, 221 |
| Villous surfaces, 245 | Zona pellucida, 244, 247, 248 |
| Villus | Zona reticularis, 213, 220, 221 |
| chorioallantoic villus, 256, 257, 258 | Zonary placenta, 244, 254, 255, 256, 257, 258 |
| digestive system, 140, 162, 164, 165, 167, 168, 180, | Zone multiformis (zona glomerulosa), 213, 220, 221 |
| 181 | Zone of calcification, 36, 37, 38 |
| female reproductive system, 256, 257, 258 | Zone of hypertrophy, <i>35</i> , <i>36</i> , <i>37</i> , <i>38</i> |
| lymphatic system, 92, 93 | Zone of multiplication, 35, 37, 38 |
| Visceral layer, 184 | Zone of ossification, 35, 36, 37, 38 |
| Visceral pleura, 196, 204 | Zone of reverse cartilage, 35, 37 |
| Visceral prepuce, 240 | Zonular fiber, 268, 269, 271, 273, 279, 280 |